

LOCAL BEHAVIOR OF PLANAR ANALYTIC VECTOR FIELDS VIA INTEGRABILITY

MARÍA JESÚS ÁLVAREZ¹ AND ANTONI FERRAGUT²

AF wants to dedicate this paper to the memory of his father

ABSTRACT. We present an algorithm to study the local behavior of singular points of planar analytic vector fields having a first integral which is a quotient of analytic functions. The algorithm is based on the blow up method. It emphasizes the curves passing through the singular points and avoids the computation of the desingularized systems. Vector fields having a rational first integral are a particular case.

1. INTRODUCTION

A *real planar analytic vector field* is a vector fields defined on \mathbb{R}^2 of the form

$$X(x, y) = P(x, y) \frac{\partial}{\partial x} + Q(x, y) \frac{\partial}{\partial y}, \quad (1)$$

where P and Q are coprime analytic functions. We refer to the vector field (1) or equivalently to its associated planar analytic differential system

$$\dot{x} = P(x, y), \quad \dot{y} = Q(x, y). \quad (2)$$

Let $m = \min\{m_P, m_Q\}$ be the *multiplicity of the vector field* (1) *at the origin*, where m_P and m_Q are the multiplicities of P and Q at the origin, respectively.

The study of the topological behavior of the solutions of a planar differential system in a neighborhood of a singular point is one of the main unsolved problems in the qualitative theory of differential systems. Concerning the singular points having at least one eigenvalue different from zero, the problem is solved except for the center-focus case. Regarding the degenerate singular points, with both eigenvalues of the jacobian matrix at the point equal to zero, the situation is more complicated. The topology around a non-monodromic singular point can be much richer. The Andreev Theorem (see [2]) classifies the nilpotent singular points (degenerate singular points whose associated jacobian matrix is not identically zero) except the center-focus case. If the jacobian matrix is identically null the problem is open. In this case, the only possibility is studying each degenerate point case by case. The main technique which is used to perform the study of this kind of points is the blow up technique, which is explained in subsection 2.2. Roughly speaking the idea behind this method is to explode, through a change of variables that is not a diffeomorphism, the singularity to a line or to a circle. Then the study of the original singular point can be reduced to the study of the new singular points that appear on this line or circle and that will be, probably, simpler. If these new singular points are again degenerate the process is repeated.

Dumortier showed in [7] that, for a given singular point of a C^∞ -Lojasiewicz system (which includes the analytic case), this chain of changes of variable is finite. However, the process of desingularizing a singular point is very long and it involves a big number of computations. There are several generalizations of this technique that consist in doing

1991 *Mathematics Subject Classification.* 34C05, 34A34, 34C14.

Key words and phrases. planar analytic vector field, generalized rational first integral, blow up.

* The authors are partially supported by MTM2008-03437. AF is partially supported by a *Juan de la Cierva* grant and by 2009SGR410.

several blow ups at the same time, see for instance [8, 1]. But although they shorten the blow up process, a previous study of the system has to be done to apply these generalizations and consequently the study of the point is still very tedious and long.

In this work we study the relationship between some integrability objects and the topological behavior of the singular points. Concretely we develop a simple algorithm which allows to completely characterize the topological behavior of the orbits of an analytic system in a neighborhood of a degenerate singular point at the origin, no matter its degeneracy, under the assumption that a generalized rational first integral is defined. This characterization is given in terms of the curves passing through the origin and of their multiplicity. As far as we know, this is the first work in which the first integral is applied to characterize the local behavior of degenerate critical points. In some sense, we blow up the first integral. As a particular case we apply the method when the system is polynomial and has a rational first integral.

The paper is structured as follows. In section 2 we give the main definitions on analytic functions and integrability, and we explain how the blow up technique works. In section 3 we provide some preliminary results that will be necessary for stating the algorithm, which is presented in section 4. Finally, in section 5 we show some examples of application. One of them considers the inverse problem of constructing differential system having a first integral and having a given set of curves as separatrices and a given distribution of canonical regions in a previously determined way. A natural question is afterwards raised.

2. BASIC DEFINITIONS AND RESULTS

2.1. Analytic functions and Integrability. We first briefly introduce the notions of formal power series and analytic functions. For more information we refer the reader to the work of Seidenberg (see [13]), see also [14, 5]. Let

$$\mathbb{C}\llbracket x, y \rrbracket = \left\{ \varphi(x, y) = \sum_{i,j} \varphi_{i,j} x^i y^j : \varphi_{i,j} \in \mathbb{C} \right\}$$

be the ring of formal power series in two variables with coefficients in \mathbb{C} . With the usual operations of addition and multiplication, this ring is factorial. The elements of the subring $\mathbb{C}\{x, y\}$ of convergent power series are said to be *analytic* functions.

Let $\varphi(x, y) \in \mathbb{C}\llbracket x, y \rrbracket \setminus \{0\}$ be an irreducible non-unit element, i.e. $\varphi(0, 0) = 0$. An *analytic branch* centered at $(0, 0)$ is the equivalence class of φ under the equivalence relation $\varphi \sim \psi$ if $\varphi = \nu\psi$, where ν is a unit element, i.e. $\nu(0, 0) \neq 0$.

A *solution* of a formal differential equation $\dot{x} = P(x, y)$, $\dot{y} = Q(x, y)$ is an analytic branch $\varphi(x, y)$ centered at the origin such that there exists $k(x, y) \in \mathbb{C}\{x, y\}$ satisfying $P\varphi_x + Q\varphi_y = k\varphi$. k is the *cofactor* of φ .

In the following we introduce some notions of integrability. If $U \subseteq \mathbb{R}^2$ is an open set, a non-constant \mathcal{C}^1 function $H : U \rightarrow \mathbb{R}$, eventually multi-valued, which is constant on all the solutions of X contained in U is a *first integral* of X on U . Moreover we have $XH = 0$ on U . The importance of the first integral is on its level sets: the existence of such a function H determines the phase portrait of the system on U , because the level sets $H = h \in H(U)$ provide the expression of the curves laying on U . Consequently, given a differential system (2), it is important to know whether it has a first integral.

Two analytic functions $f(x, y)$ and $g(x, y)$ defined on a subset $U \subset \mathbb{R}^2$ are said to be *coprime* if the set of points $\{(x, y) \in U : f(x, y) = g(x, y) = 0\}$ is isolated. We call the ratio of two coprime analytic functions a *generalized rational function*. A generalized rational function $H = f/g$ defined on U is a first integral of system (2) if $\Sigma = \{(x, y) \in U : g(x, y) = 0\}$ is a set of integral curves of system (2) and H is a first integral on $U \setminus \Sigma$. Obviously, $H = f/g$ is a first integral of system (2) if and only if $(Xf)g - (Xg)f = 0$ on U .

The following theorem (see [11]) is an extension of the Poincaré Theorem (see [12]) for generalized rational first integrals.

Theorem 1. *Assume that the origin is an elementary singular point of the analytic differential system*

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = A \begin{pmatrix} x \\ y \end{pmatrix} + \dots \quad (3)$$

with eigenvalues $\lambda_1 \neq 0$ and λ_2 . Then system (3) has a generalized rational first integral in some neighborhood of the origin if and only if one of the following conditions holds:

- (i) $\lambda_1 \neq 0 = \lambda_2$ and the origin is not an isolated singular point;
- (ii) $\lambda_1/\lambda_2, \lambda_2/\lambda_1 \in \mathbb{Q}^+ \setminus \mathbb{N}$;
- (iii) $\lambda_1 = \lambda_2 \neq 0$, $A = \text{diag}(\lambda_1, \lambda_2)$;
- (iv) $\lambda_1/\lambda_2 \in \mathbb{N} \setminus \{1\}$ or $\lambda_2/\lambda_1 \in \mathbb{N} \setminus \{1\}$ and the germ (3) is analytically equivalent to its linear part;
- (v) $\lambda_1/\lambda_2 \in \mathbb{Q}^-$ and the germ (3) is analytically orbitally equivalent to its linear part.

From this theorem we know that if an analytic system has a generalized rational first integral then any elementary singular point must be a saddle, a center or a node; it cannot be neither a focus nor a saddle-node.

The notion of remarkable curve of a rational first integral was introduced by Poincaré (see [12]). It is proved in [4] that there are finitely many remarkable values for a given rational first integral. As far as we know, since Poincaré's ones very few results have been published about the remarkable values with the exception of these last years (see [4] and [10]).

We next introduce the notions of remarkable values and remarkable curves for generalized rational first integrals. We say that $c \in \mathbb{C} \cup \{\infty\}$ is a *remarkable* value of a generalized rational first integral $H = f/g$ if $f + cg = f_1^{n_1} \cdots f_r^{n_r}$, where $r, n_i \in \mathbb{N}$ and $f_i \in \mathbb{C}\{x, y\}$ are non-constant solutions of system (2), for all $i \in \{1, \dots, r\}$. The curves $f_i = 0$ are called *remarkable* curves and the n_i are their exponents. Here if $c = \infty$ then $f + cg$ denotes g . If some exponent n_i is bigger than one, then c and f_i are said to be *critical*. Finally we define the *remarkable factor* R to be the product of all the critical remarkable curves of H powered to their corresponding exponent minus one. We note that the function R is the greatest common divisor of $g^2 H_x$ and $g^2 H_y$.

Next proposition improves a result of [10] about the relationship between the exponents of two curves passing through an elementary singular point and its behavior.

Proposition 2. *Assume that the differential system (2) has a generalized rational first integral $H = f/g$. Suppose that the origin is an elementary singular point and that exactly two branches (real or complex) of the curve $fg = 0$ cross it. Suppose that the two branches correspond to irreducible solutions $f_1 = 0$ and $f_2 = 0$, and let $n_1, n_2 \in \mathbb{Z} \setminus \{0\}$ be their respective exponents in the expression of H . Then:*

- (i) *If $n_1 n_2 < 0$ then the origin is a node.*
- (ii) *If $n_1 n_2 > 0$ and both branches are real, then it is a saddle.*
- (iii) *If $n_1 n_2 > 0$ and both branches are complex conjugate, then it is a center.*

Moreover, the quotient of the eigenvalues at the origin is a positive rational multiple of $-n_1/n_2$.

Proof. Set $H = \prod_{i=0}^p f_i^{n_i}$ and let k_i be the cofactor of f_i , for all i . Applying $Xf_i = k_i f_i$ at the origin, as $f_i(0, 0) \neq 0$ for $i > 2$ and $(0, 0)$ is a singular point of the system, one obtains $k_i(0, 0) = 0$ for all $i > 2$.

As $XH = 0$, applying this expression at the origin and after straightforward computations, we have $\sum n_i k_i(0, 0) = 0$, and then $n_1 k_1(0, 0) + n_2 k_2(0, 0) = 0$. Applying similar

arguments to those of [5], both cofactors at the origin are multiple of the two eigenvalues of the vector field at $(0,0)$, say $k_1(0,0) = s_1\lambda$ and $k_2(0,0) = s_2\mu$, for $s_1, s_2 \in \mathbb{N}$ and λ, μ the eigenvalues. Hence

$$\frac{s_1 \lambda}{s_2 \mu} = \frac{k_1(0,0)}{k_2(0,0)} = -\frac{n_2}{n_1},$$

and therefore the proposition follows. \square

2.2. The technique of the blow up. Consider the real planar analytic differential system

$$\dot{x}_i = P(x_i, y_i), \quad \dot{y}_i = Q(x_i, y_i) \quad (4)$$

in the variables (x_i, y_i) and assume that the origin is a degenerate singular point of this system. The *directional blow up in the x_i direction* (resp. *y_i*) is the mapping $(x_i, y_i) \mapsto (x_{i+1}, x_{i+1}y_{i+1})$ (resp. $(x_i, y_i) \mapsto (x_{i+1}y_{i+1}, y_{i+1})$), where $x_{i+1} = x_i$ (resp. $y_{i+1} = y_i$) and y_{i+1} (resp. x_{i+1}) is a new variable. This transformation converts the origin of the (x_i, y_i) -plane into the line $x_{i+1} = 0$ (resp. $y_{i+1} = 0$). The expression of system (4) after the blow up, for instance in the x_i direction, is

$$\begin{aligned} \dot{x}_{i+1} &= P(x_{i+1}, x_{i+1}y_{i+1}), \\ \dot{y}_{i+1} &= \frac{Q(x_{i+1}, x_{i+1}y_{i+1}) - y_{i+1}P(x_{i+1}, x_{i+1}y_{i+1})}{x_{i+1}}, \end{aligned} \quad (5)$$

that is always well-defined since we are assuming that the origin is a singularity.

Observe that, after the blow up, x_{i+1}^{m-1} is a common factor of \dot{x}_{i+1} and \dot{y}_{i+1} . Thus we scale the independent variable to remove it. Along all this paper, when working with system (5) we will assume that such a reparametrization has been done.

We reproduce in the following two well-known results (see [3]) that provide the relationship between the original singular point of system (4) and the new singularities of system (5). We recall that a *characteristic direction* of system (4) is a solution of the equation

$$\mathcal{F}_m(x_i, y_i) := x_i Q_m(x_i, y_i) - y_i P_m(x_i, y_i) = 0, \quad (6)$$

provided that this polynomial is not identically zero. We call \mathcal{F}_m the *characteristic polynomial*. If $\mathcal{F}_m \not\equiv 0$ and φ_t is a solution of system (4) tending to the origin in forward or backward time, then it must do it tangent to one of the characteristic directions.

Proposition 3. *Let $\varphi_t = (x_i(t), y_i(t))$ be a trajectory tending to the origin of system (4), in forward or backward time. Suppose that $\mathcal{F}_m \not\equiv 0$. Assume that φ_t is tangent to one of the two angle directions $\tan \theta = v$, $v \neq \infty$. Then the following statements hold.*

- (i) *The two angle directions $\theta = \arctan v$ (in $[0, 2\pi)$) are characteristic directions.*
- (ii) *The point $(0, v)$ on the (x_{i+1}, y_{i+1}) -plane is an isolated equilibrium point of system (5).*
- (iii) *The trajectory φ_t corresponds to a solution of system (5) tending to the singular point $(0, v)$.*
- (iv) *Conversely, any solution of system (5) tending to the singular point $(0, v)$ on the (x_{i+1}, y_{i+1}) -plane corresponds to a solution of system (4) tending to the origin in one of the two angle directions $\tan \theta = v$.*

The conclusion of the previous proposition is that in order to study the behavior of the solutions around the origin of system (4) it is enough to study the singular points of the form $(0, v)$ of system (5), that will be simpler. But, as we said before, it is possible that, despite they are simpler, some of them are still quite complicated. If this is the case, then we have to study these degenerate singularities by blowing them up and repeating the process.

If $\mathcal{F}_m \equiv 0$ we apply the following result.

Proposition 4. *If $\mathcal{F}_m \equiv 0$ then system (4) assumes the form $\dot{x} = xW_{m-1} + \dots$, $\dot{y} = yW_{m-1} + \dots$, with $W_{m-1} \not\equiv 0$ a homogeneous polynomial of degree $m-1$. In this case, for every nonsingular direction θ (i.e. the direction not satisfying the equation $W_{m-1}(x, y) = 0$) there exists exactly one semipath tending to the origin in the direction θ . If θ^* is a singular direction, there may be either no semipaths tending to the origin in the direction θ^* , or a finite number, or infinitely many.*

3. PRELIMINARY RESULTS

In this section we state and prove several results that will be useful in order to show that the algorithm that we state in section 4 works. All along this section we work with an analytic system of type (4) and its corresponding blown up system (5) in the x_i direction (the case where the y_i directional blow up is applied follows in a similar way). We assume that system (4) has a generalized rational first integral $H = f/g$.

We denote by m_h the multiplicity of an analytic function h at the origin. We also assume that for every system only one directional blow up is needed. This means the following: if we want to do the blow up for instance in the x_i direction to system (4), then there is no curve tending to the origin in this direction. This can be easily ensured by a convenient rotation of the system.

We use the notation f , g and R also for the corresponding numerator, denominator and remarkable factor of the blown up first integrals associated to H . In a similar way we use \mathcal{F}_m and we use m to refer to the multiplicity at a considered singular point.

In the whole process of desingularization we denote the variables of the systems as (x_i, y_i) , $i \in \mathbb{N} \cup \{0\}$. We start with $(x_0, y_0) = (x, y)$; the $(i+1)$ -th blow up is denoted by $(x_i, y_i) \rightarrow (x_{i+1}, y_{i+1})$, for $x_i = x_{i+1}$ and $y_i = x_{i+1}y_{i+1}$, and it goes from the i -th desingularization of the initial system (2) to the $(i+1)$ -th one. The results in this section are stated for system (4).

First we remark how the integrability objects are transformed after the $(i+1)$ -th blow up.

Lemma 5. *If $h(x_i, y_i) = 0$ is a solution of system (4) with cofactor k , then the function $h(x_{i+1}, x_{i+1}y_{i+1})/x_{i+1}^{m_h} = 0$ is a solution of system (5) with $k(x_{i+1}, x_{i+1}y_{i+1})/x_{i+1}^{m_k}$ as cofactor. Moreover the functions $H(x_{i+1}, x_{i+1}y_{i+1})$ and $x_{i+1}^\omega R|_{y_i=x_{i+1}y_{i+1}}$, where $\omega = |m_f - m_g| - 1 - m_R$, are respectively a first integral and the remarkable factor of system (5).*

From lemma 10 below we obtain the multiplicity of R for system (2) in terms of the multiplicities of the system and of the curves f and g . The multiplicities of the remarkable factors of the blown up systems can be computed using the multiplicity of R and lemma 5. We note that we do not need the expression of R but its multiplicity, which is computable using by lemma 10.

In the case where H is a rational first integral, all the remarkable curves of H can be computed, as there are in the literature several methods to compute them, for instance the one concerning the extactic curves (see [6, 10]) and a new one provided in [9].

The proof of lemma 5 follows from straightforward computations. Obviously they also follow when the blow up $x_i = x_{i+1}y_{i+1}$ is applied instead of $y_i = x_{i+1}y_{i+1}$.

The following proposition allows to control whether the characteristic polynomial \mathcal{F}_m of a blown up system is identically zero without computing the differential system explicitly. We denote by \hat{h} the homogeneous polynomial of lowest degree of an analytic function h .

Proposition 6. *We have $\mathcal{F}_m \equiv 0$ if and only if $m_{f+cg} = m_g$ for all $c \in \mathbb{C}$.*

Proof. Suppose that there is no $s \in \mathbb{C}$ such that $\hat{f} + s\hat{g} \equiv 0$. Then $f + cg$ has always the same multiplicity and therefore $(\hat{f} + c\hat{g})|_{y_i=x_{i+1}y_{i+1}}/x_{i+1}^{m_g}$ is a polynomial in y_{i+1} with c as

a parameter that has, varying c , infinitely many solutions. Therefore $\mathcal{F}_m = 0$ has infinitely many roots. As \mathcal{F}_m is to be a polynomial, we have $\mathcal{F}_m \equiv 0$.

On the other side, if $\mathcal{F}_m \equiv 0$ then the origin is crossed by the solutions of the system with infinitely many slopes. Suppose that there exists $s \in \mathbb{C}$ such that $m_{f+sg} \neq m_g$. Assume, without loss of generality, that $s = 0$. Then $\widehat{f + cg} = 0$ is equivalent either to $\hat{g} = 0$ or to $\hat{f} = 0$ for all $c \in \mathbb{C}$ and therefore the number of different slopes is finite, a contradiction. \square

Remark 1. *If $\mathcal{F}_m \not\equiv 0$ then there exists $s \in \mathbb{C} \cup \{\infty\}$ such that $m_{f+sg} > m_{f+cg}$ for all $c \in \mathbb{C} \cup \{\infty\}$, $c \neq s$.*

3.1. The dicritical case. The case $\mathcal{F}_m \equiv 0$ is called the *dicritical case*. It is known from proposition 4 that in the dicritical case we can write $P_m = xW_{m-1}$ and $Q_m = yW_{m-1}$, where W_{m-1} is a homogeneous polynomial of degree $m-1$. Moreover, the blown up system (5) has a line of singularities, all of them semi-hyperbolic except a finite number. This finite set of singular points corresponds to the singular directions of system (4) and is obtained from the equation $W_{m-1} = 0$. Once they are known, they can be studied separately.

The following proposition, due to Maria Alberich and AF, allows to compute the singular directions in terms of the first integral.

Proposition 7. *Let w be a homogeneous polynomial. Let $e_3 \in \mathbb{N} \cup \{0\}$ be the exponent of w in the factorization of \hat{R} . Consider the following property:*

(H1) *There exists $c \in \mathbb{C} \cup \{\infty\}$ such that w is a multiple factor of $\hat{f} + c\hat{g}$ with multiplicity $e_1 \in \mathbb{N} \setminus \{1\}$ and $w^{e_1} \nmid \gcd(\hat{f}, \hat{g})$.*

Then a divisor w of W_{m-1} either satisfies (H1) or $w \mid \gcd(\hat{f}, \hat{g})$. Conversely, let w be a homogeneous polynomial such that either (H1) holds or w divides $\gcd(\hat{f}, \hat{g})$ with multiplicity $e_2 \in \mathbb{N}$. Then $w^e \mid W_{m-1}$ and $w^{e+1} \nmid W_{m-1}$, where $e = e_1 - 1 + e_2 - e_3$ if (H1) holds (here $e_2 = 0$ if $w \nmid \gcd(\hat{f}, \hat{g})$) and $e = e_2 - e_3$ otherwise.

Proof. Let $S^{x_i} := g^2 H_{x_i} = f_{x_i} g - f g_{x_i}$ and $S^{y_i} := -g^2 H_{y_i} = -f_{y_i} g + f g_{y_i}$. As H is a first integral of system (2) we have $(P, Q) = (S^{y_i}, S^{x_i})/R$. From the equalities $\widehat{S^{y_i}} = -\hat{f}_{y_i} \hat{g} + \hat{f} \hat{g}_{y_i}$ and $\widehat{S^{x_i}} = \hat{f}_{x_i} \hat{g} - \hat{f} \hat{g}_{x_i}$, it is clear that $\gcd(\hat{f}, \hat{g})$ divides $\widehat{S^{x_i}}$ and $\widehat{S^{y_i}}$.

Now if $c \in \mathbb{C} \cup \{\infty\}$ is such that $\hat{f} + c\hat{g} = 0$ has a multiple factor w , then both $(\hat{f} + c\hat{g})_{x_i}$ and $(\hat{f} + c\hat{g})_{y_i}$ vanish on $w = 0$. Hence on $w = 0$ we have

$$\frac{\hat{f}}{\hat{g}} = \frac{\hat{f}_{y_i}}{\hat{g}_{y_i}} = \frac{\hat{f}_{x_i}}{\hat{g}_{x_i}} = -c,$$

and therefore w divides both $\widehat{S^{x_i}}$ and $\widehat{S^{y_i}}$.

On the other hand, let w be a common factor of $\widehat{S^{x_i}}$ and $\widehat{S^{y_i}}$. Then on $w = 0$ we have

$$\hat{f}_{y_i} \hat{g} = \hat{f} \hat{g}_{y_i}, \quad \hat{f}_{x_i} \hat{g} = \hat{f} \hat{g}_{x_i}. \quad (7)$$

If w divides $\gcd(\hat{f}, \hat{g})$ with multiplicity $e_2 \in \mathbb{N}$ then these equalities hold on $w = 0$. Moreover we can write (7) as

$$\frac{\hat{f}}{\hat{g}} = \frac{\hat{f}_{y_i}}{\hat{g}_{y_i}} = \frac{\hat{f}_{x_i}}{\hat{g}_{x_i}}.$$

All the polynomials in these equalities are homogeneous and the numerators and denominators have the same degree two by two, hence all the quotients are equal to a constant, say $-c$. Then there exists $e_1 \in \mathbb{N}$, $e_1 > e_2$, such that $w^{e_1} \mid (\hat{f} + c\hat{g})$.

The expression of e follows from the explanation above and $(P, Q) = (S^{y_i}, S^{x_i})/R$. \square

Remark 2. Proposition 7 allows to compute the singular directions without computing the differential system explicitly. Moreover as a consequence of the computation, the value of m appears naturally.

Proposition 8. Suppose that the origin is a singular point corresponding to a singular direction. Then:

- (i) If $m > 1$ and the origin is dicritical, then another blow up is required.
- (ii) If $m = 1$ and the origin is dicritical, then it is a star-node.

The study of the case where the origin is not dicritical is done in the next subsection.

3.2. The non-dicritical case. The dicritical case leads either to the non-dicritical one or to a star-node, hence it remains to study the non-dicritical case. From now on in this section we assume that $\mathcal{F}_m \neq 0$.

If $m_f = m_g$ then there exists $s \in \mathbb{C} \cup \{\infty\}$ such that $m_{f+sg} > m_g$ (see remark 1). This curve factorizes after the blow up (for instance $y_i = x_{i+1}y_{i+1}$ and after removing $x_{i+1}^{m_g}$) as a positive power of x_{i+1} and another polynomial, say W . Therefore from the intersection of the curves $x_{i+1} = 0$ and $W = 0$ new singular points may appear.

From now on and until the end of section 4 we assume that the curve $f + sg = 0$ defined in remark 1 is in the numerator of H , that is $s = 0$. This transformation can be easily done taking $H + s = (f + sg)/g$ as first integral instead of H .

Remark 3. From lemma 5 we know that $x_{i+1} = 0$ is a remarkable curve of (5) and that the first integral of system (5) has the same remarkable values as H and also $c = 0$, as we are assuming that $m_f > m_g$.

The following proposition ensures that all the orbits that are needed in the desingularization process are contained into the curves appearing in the expression of H .

Proposition 9. The whole set of characteristic directions of the differential system (4) at the origin is obtained from the tangents at the origin of $fg = 0$. I.e., it is the set of solutions y_i/x_i of the equation $\widehat{fg} = 0$.

Proof. As $m_f > m_g$, $\widehat{f + cg}$ is equal to \hat{g} for all $c \neq 0$ and to \hat{f} for $c = 0$. Hence all the characteristic directions of all the solutions of the system at the origin are found either in $\hat{f} = 0$ or in $\hat{g} = 0$. \square

Remark 4. By proposition 9 we can compute the singular points on $x_{i+1} = 0$ after the blow up $y_i = x_{i+1}y_{i+1}$ without computing the differential system explicitly. From proposition 2, if the singular points are elementary then we can characterize them. Otherwise a new blow up is required.

Next result allows to compute the multiplicity m of system (4) at the origin. We shall see in section 4 that the knowledge of m is a key point in the application of our algorithm.

Lemma 10. We have

$$m_f + m_g - m_R = m + 1. \quad (8)$$

Proof. We write P and Q in terms of f and g :

$$P = -\frac{f_{y_i}g - fg_{y_i}}{R}, \quad Q = \frac{f_{x_i}g - fg_{x_i}}{R},$$

where R is the remarkable factor. We have

$$R(x_iQ - y_iP) = \left(x_i \frac{\partial f}{\partial x_i} + y_i \frac{\partial f}{\partial y_i}\right)g - \left(x_i \frac{\partial g}{\partial x_i} + y_i \frac{\partial g}{\partial y_i}\right)f = (m_f - m_g)\hat{f}\hat{g} + \dots$$

Therefore the lemma follows directly taking multiplicities in the equalities above, as we are assuming $\mathcal{F}_m \neq 0$ and $m_f \neq m_g$. \square

Remark 5. *When applying a blow up to a differential system (4), by means of a change of time we cancel a factor x_{i+1}^{m-1} appearing in both \dot{x}_{i+1} and \dot{y}_{i+1} . Depending on m , this change of time implies a change in the orientation of the orbits contained into the half-plane $x_{i+1} < 0$. This happens only if m is even. Thus from proposition 10 in the non-dicritical case and also from proposition 7 in the dicritical case we can compute m to know if such a change of time is to be done, altogether without computing the system explicitly.*

Proposition 11. *Assume that a blow up $y_i = x_{i+1}y_{i+1}$ is applied to system (4). Suppose that the origin is a singular point of system (5) with multiplicity m . Then:*

- (i) *If $m > 1$ then the origin is degenerate.*
- (ii) *If $m = 1$ and both $f = 0$ and $g = 0$ pass through the origin and $g = 0$ does it not transversally to $x_{i+1} = 0$, then it is nilpotent.*
- (iii) *If $m = 1$ and both $f = 0$ and $g = 0$ pass through the origin but $g = 0$ does it transversally to $x_{i+1} = 0$, then it is a node. Moreover $f + cg = 0$ crosses the origin of system (4) with the slope of $g = 0$, for all $c \in \mathbb{C} \setminus \{0\}$.*
- (iv) *If $m = 1$ and only $f = 0$ passes through the origin, then it is a saddle. Moreover the separatrices of the origin of system (4) are contained in $f = 0$.*

Proof. We prove each subcase separately.

- (i) The origin of system (5) is not elementary, so a new blow up is required.
- (ii) As two curves belonging to two different level sets of H meet the origin in the same direction, it is neither a saddle (which would require exactly one level) nor a node (which would require at least two directions). Thus as $m = 1$ and the origin cannot be semi-hyperbolic (see theorem 1), it is nilpotent.
- (iii) The singular point is a node as it is elementary and $f + cg = 0$ crosses it for all $c \in \mathbb{C} \cup \{\infty\}$.
- (iv) The singular point is a saddle as both curves $x_{i+1} = 0$ and $f = 0$ belong to the same level set $c = 0$ of H .

□

Remark 6. *We remind here a well-known result due to Seidenberg (see [13]): if a differential system has a node at the origin, then there is exactly one branch crossing the origin with a determined slope and there are infinitely many branches crossing the origin with another determined slope. In our case, $f = 0$ contains the first branch and $f + cg = 0$, with $c \neq 0$, contain the rest of the branches.*

4. THE ALGORITHM

We explain in this section how our algorithm works. First we assume that the systems we deal with have some properties that are stated in the next subsection.

4.1. Assumptions.

- (i) If $\mathcal{F}_m \neq 0$ then we take $H = f/g$ with $m_f > m_g$.
- (ii) We suppose that only one of the directional blow ups is to be done.

We make these assumption for all the systems appearing after the different blow ups. We note that they are not restrictive, they are done for a better understanding of the explanation and the process.

4.2. Statement of the algorithm.

We describe each step of the algorithm.

- (a) We check whether $\mathcal{F}_m \neq 0$ and $m_f = m_g$. If this is the case, there exists $s \in \mathbb{C}$ such that $m_{f+sg} > m_g$ and we take $f + sg$ as the numerator of H instead of f .
- (b) If $\mathcal{F}_m \neq 0$ then we compute the singular points on $x_{i+1} = 0$ (or $y_{i+1} = 0$, depending on the direction of the blow up) from the curves $f = 0$ (after dropping the factor $x_{i+1}^{m_f - m_g}$) and $g = 0$ (see proposition 9). If $\mathcal{F}_m \equiv 0$ then we follow proposition 7.

- (c) For each singular point of step (b) we compute the multiplicity m of the blown up system at this point and check whether another blow up is required. This can be done using propositions 11 and 7.
- (d) No new desingularization is to be done for elementary singular points (meaning saddles, nodes or centers). For the degenerate singular points a new blow up is required. In this case we check the initial assumptions for the new system and go back to step (a).

The algorithm ends as the chain of blow ups is finite.

The construction of a table is very useful to follow the desingularization process. In this table each row corresponds to a step of the algorithm, i.e. to a blow up. Each change of variables is written in the first column. Two columns named SP_f and SP_g show the singular points that we obtain from f and g , respectively. In the dicritical case we write the dicritical points in the cells of both SP_f and SP_g . Three more columns \hat{f} , \hat{g} and \hat{R} show the lowest terms of f , g and R after the singular point is moved to the origin. We shall write a \star in the cells where no new (relevant) information is to be added.

When the table is done all the singular points appearing from all the necessary blow ups have been computed and studied. It is clear in the non-dicritical case that the singular points come from the intersection of $\hat{f} = 0$ and/or $\hat{g} = 0$ with $x_{i+1} = 0$ on each step. We can also know their behavior from the multiplicity m of the system at the points and from f and g , as we stated in proposition 11. The dicritical case reduces to the non-dicritical one or to the star-node from proposition 8.

4.3. Construction of the local phase portraits. Once the desingularization process is finished, we need to *go back* to the initial system. We start at the last system of the desingularization, say (x_n, y_n) , for some $n \in \mathbb{N}$, which corresponds to the last row of the table. We situate on the (x_n, y_n) -plane all the singular points (say on $x_n = 0$) and the lines crossing the y_n axis at these points corresponding to the curves belonging to f or g that provide these singular points. The local behavior of the system at all these singular points is known: they are saddles, nodes or centers. The plane is then divided into several canonical regions and we know the behavior of the system at all of them.

Next we check if some half-plane must change the orientation (see remark 5). We also notice that the quadrants either II and IV or III and IV are to be swapped as it is usual in the blow up process.

Now we change the variables into the previous ones in the desingularization process, (x_{n-1}, y_{n-1}) . The curves we drew are transformed into new curves by the change of variables; for instance, the curve $y_n = a + \dots$, $a \in \mathbb{R}$, would be transformed into $y_{n-1} = ax_{n-1} + \dots$, as the change of variables was $y_{n-1} = x_n y_n$. All the singular points of the (x_n, y_n) -plane on $x_n = 0$ meet now at the origin. The shapes and situation of the canonical regions can also be modified. The axis remain invariant if they appear in the corresponding expression of the first integral.

We repeat this procedure until we obtain the local phase portrait of the initial system at the origin with the initial variables $(x_0, y_0) = (x, y)$, and then we are finished.

Note that, in the dicritical case, all the points on $x_{i+1} = 0$ are singular, and all of them except a finite number are semi-hyperbolic. Thus for each one of them, say $(0, v)$, there exists exactly one curve on the (x_i, y_i) -plane crossing the origin with slope v (see [3]).

Remark 7. *The algorithm we have shown allows to completely study the local behavior around a singular point, no matter how degenerate it is, without needing to use the blow up technique explicitly. We have presented an alternative method to this technique for planar differential systems having a generalized rational first integral which uses the information that is provided by some specific curves crossing the singular points, that are also computed.*

5. EXAMPLES

We present in this last section some examples in order to illustrate how the algorithm must be applied. In all cases we show the corresponding table of desingularization and a figure with all the different phase portraits that we need to obtain the phase portrait of the initial system.

As we said in the introduction, a particular case of analytic curves are the polynomial ones. The first example deals with a polynomial system having a rational first integral.

Example 1. Consider the rational function $H = f/g$, where $f(x, y) = -(x^{10} + 8x^{18} + 12x^{14}y + 6x^{10}y^2 + x^6y^3 - 24x^{12}y^4 - 24x^8y^5 - 6x^4y^6 + 24x^6y^8 + 11x^2y^9 - y^{10} - 8y^{12})(-x^{10} + 8x^{18} + 12x^{14}y + 6x^{10}y^2 + x^6y^3 - 24x^{12}y^4 - 24x^8y^5 - 6x^4y^6 + 24x^6y^8 + 13x^2y^9 + y^{10} - 8y^{12})$ and $g(x, y) = (x^2y - 2y^4 + 2x^6)^6$. We want to study the local behavior of the singular point at the origin of the polynomial differential system associated to H . As f and g have both multiplicity eighteen at the origin, we rename $f + g$ (which has multiplicity twenty) as f . Hence we set $f(x, y) = (x^{10} - y^{10} - x^2y^9)^2$. Moreover as $g = 0$ has a vertical tangent at the origin, we apply the change $x \rightarrow x + 3y$ to both functions. We set $x_0 := x$ and $y_0 := y$.

	SP_f	SP_g	\hat{f}	\hat{g}	\hat{R}
$y_0 = x_1y_1$	$y_1 = -\frac{1}{2}$ $y_1 = -\frac{1}{4}$ ★ ★	★ ★ $y_1 = 0$ $y_1 = -\frac{1}{3}$	$x_1^2l_1^2$ $x_1^2l_2^2$ x_1^2 x_1^2	★ ★ y_1^6 x_1^6	x_1l_1 x_1l_2 $x_1y_1^5$ x_1^6
$y_1 \rightarrow y_1 - \frac{1}{3}$ $x_1 = x_2y_2$	$x_2 = 0$	$x_2 = 0$	x_2^2	$y_2^4l_3^6$	$x_2y_2^3l_3^5$
$x_2 = x_3y_3$	$x_3 = 0$ ★	★ $x_3 = -\frac{243}{2}$	x_3^2 ★	y_3^8 $y_3^8l_4^6$	$x_3y_3^7$ $y_3^7l_4^5$

TABLE 1. Application of the algorithm in example 1. The l_i are straight lines crossing the corresponding singular point with neither horizontal nor vertical tangency. In particular, $l_3 = 2x_2 + 243y_2$.

We construct table 1 as it is explained in section 4. Three blow up are needed to completely desingularize the singular point at the origin. From table 1 we can study all the singular points appearing in the whole blow up process:

- (1) First blow up, $x_0 = x_1$, $y_0 = x_1y_1$:
 - $(0, 0)$: as $m_R = 6$ and $m_f + m_g = 8$ we have $m = 1$. Moreover both $f = 0$ and $g = 0$ pass through this point and $g = 0$ does it transversally, hence it is a node.
 - $(0, -1/2)$ and $(0, -1/4)$: as $m_R = 2$ and $m_f + m_g = 4$ we have $m = 1$ in both cases. Moreover, only $f = 0$ passes through these points, hence they are saddles.
 - $(0, -1/3)$: as $m_R = 6$ and $m_f + m_g = 8$ we have $m = 1$. Moreover, both $f = 0$ and $g = 0$ pass through this point and $g = 0$ does it not transversally, hence the point is nilpotent and a new blow up is required.

Before the second blow up we move the point $(0, -1/3)$ to the origin.

- (2) Second blow up, $x_1 = x_2y_2$, $y_1 = y_2$:
 - $(0, 0)$: as $m_R = 9$ and $m_f + m_g = 12$ we have $m = 2$, hence a new blow up is required.
- (3) Third blow up, $x_2 = x_3y_3$, $y_2 = y_3$ (where we take into account that now $m_f < m_g$ and the roles of f and g are swapped):




FIGURE 1. The desingularization of example 1. The numbers indicate the canonical regions of the systems to show how they change during the desingularization process. The biggest point represents the origin of each system.

- $(0, 0)$: as $m_R = 8$ and $m_f + m_g = 10$ we have $m = 1$. Moreover, both $f = 0$ and $g = 0$ pass through this point and $f = 0$ does it transversally, hence it is a node.
- $(-243/2, 0)$: as $m_R = 12$ and $m_f + m_g = 14$ we have $m = 1$. Moreover, only $g = 0$ passes through this point, hence it is a saddle.

Now the desingularization process is done. Next we explain how to get the phase portrait of the initial system to end the process.

- (1) After the third blow up we obtain two singular points on the (x_3, y_3) -plane coming from the intersection of $y_3 = 0$ and the curves $x_3 = 0$ and $x_3 = -243/2 + \mathcal{O}(y_3)$.
- (2) Back to the (x_2, y_2) -plane we study the origin. The canonical regions of the (x_3, y_3) -system are modified and we have swapped the third and fourth quadrants of the (x_3, y_3) -plane. The curve $y_2 = 0$ remains invariant, and the others become $x_2 = 0$ and $x_2 = -243y_2/2 + \mathcal{O}(y_2^2)$.
- (3) Back to the (x_1, y_1) -plane and after swapping again the third and fourth quadrants, $y_2 = 0$ disappears as solution, $x_2 = 0$ becomes $x_1 = 0$ and $x_2 = -243y_2/2 + \mathcal{O}(y_2^2)$ becomes $x_1 = -243y_1^2/2 + \mathcal{O}(y_1^3)$. After this update we undo the change $y_1 \rightarrow y_1 - 1/3$ and the singular point is now at $y_1 = -1/3$. There are three more singular points, as table 1 showed.
- (4) Back to the initial system on the (x_0, y_0) -plane and after swapping the second and third quadrants, $x_1 = 0$ disappears as solution and only some branches of $f = 0$ and $g = 0$ remain as separatrices. $f = 0$ provides an elliptic sector and $g = 0$ a hyperbolic sector.

A diagram of the whole process is shown in figure 1. \square

In the next example we deal with an analytic system having a generalized rational first integral.

Example 2. Let

$$\begin{aligned} f(x, y) &= 3y^4 + x^2y^3 - 2x^3y^2 + 3x^5y - x^4y^2 + 2x^7 - 2x^5y^2 + xy^6 + \\ &\quad 3x^4y^4 + y^9 + x^{12}y^8 + \dots, \\ g(x, y) &= x^2 - 2y^2 + x^2y - 3x^3y + xy^3 + 5x^5 - 4x^3y^2 + y^5 + \dots, \end{aligned}$$

where the dots mean higher order terms, be two analytic functions and let $H = f/g$. Consider the analytic differential system associated to H . We apply our algorithm in order to study the local behavior around the singular point at the origin of this system. We construct table 2, from which we study all the singular points appearing in the whole blow up process. As $m_f = 4$, $m_g = 2$ and $m = 5$, we have $m_R = 0$ from proposition 10. Therefore $R = 1$. After the first blow up $y_0 = x_1y_1$ the remarkable factor is x_1 .

	SP_f	SP_g	\hat{f}	\hat{g}	\hat{R}
$y_0 = x_1y_1$	$y_1 = 0$ ★ ★	★ $y_1 = -\frac{\sqrt{2}}{2}$ $y_1 = \frac{\sqrt{2}}{2}$	$x_1^3l_1l_2$ x_1^2 x_1^2	★ l_3 l_3	x_1 x_1 x_1
$x_1 = x_2y_2$	$x_2 = 0$ $x_2 = \frac{1}{2}$ $x_2 = -2$	★ ★ ★	$x_2^2y_2^5l_4$ $y_2^5l_5$ $y_2^5l_6$	★ ★ ★	$x_2y_2^4$ y_2^4 y_2^4
$x_2 = x_3y_3$	$x_3 = 0$ $x_3 = \frac{3}{2}$	★ ★	$x_3^2y_3^8$ $y_3^8l_7$	★ ★	$x_3y_3^7$ y_3^7

TABLE 2. Application of the algorithm in example 2. The l_i are straight lines crossing the corresponding singular point with neither horizontal nor vertical tangency. In particular $l_1 = 2x_1 - y_1$, $l_2 = x_1 + 2y_1$, $l_3 = x_1 - 4y_1$ and $l_4 = 2x_2 - 3y_2$.

- (1) First blow up, $x_0 = x_1$, $y_0 = x_1y_1$:
 - $(0, 0)$: as $m_R = 1$ and $m_f + m_g = 5$ we have $m = 3$, hence a new blow up is required.
 - $(0, \sqrt{2}/2)$ and $(0, -\sqrt{2}/2)$: as $m_R = 1$ and $m_f + m_g = 3$ we have $m = 1$ in both cases. Moreover, both $f = 0$ and $g = 0$ pass through these points and $g = 0$ does it transversally, hence they are nodes.
- (2) Second blow up, $x_1 = x_2y_2$, $y_1 = y_2$:
 - $(0, 0)$: as $m_R = 5$ and $m_f + m_g = 8$ we have $m = 2$, hence a new blow up is required.
 - $(1/2, 0)$ and $(-2, 0)$: as $m_R = 4$ and $m_f + m_g = 6$ we have $m = 1$ in both cases. Moreover, only $f = 0$ passes through this point, hence they are saddles.
- (3) Third blow up, $x_2 = x_3y_3$, $y_2 = y_3$:
 - $(0, 0)$: as $m_R = 8$ and $m_f + m_g = 10$ we have $m = 1$. Moreover, only $f = 0$ passes through this point, hence it is a saddle.
 - $(3/2, 0)$: as $m_R = 7$ and $m_f + m_g = 9$ we have $m = 1$. Moreover, only $f = 0$ passes through this point, hence it is a saddle.

A diagram of the whole process is shown in figure 2. \square

The following example appears in [3]. It deals with the dicritical case.




FIGURE 2. The desingularization of example 2.

	SP_f	SP_g	\hat{f}	\hat{g}	\hat{R}
$\mathcal{F}_m \equiv 0$	$(0, 0)$		y_0^2	$(x_0 + y_0)^2$	$x_0 + y_0$
$y_0 = x_1 y_1$	$y_1 = 0$	*	$x_1^2 + y_1^2$	*	1

TABLE 3. Application of the algorithm in example 3.

Example 3. Let $f(x, y) = y^2 + (x + y)^4$ and $g(x, y) = (x + y)^2$ be two polynomials and let $H = f/g$. Consider the polynomial system associated to H . The polynomials f and g have the same multiplicity at the origin, but there is no $s \in \mathbb{C}$ such that $m_{f+sg} > m_g$, hence we are in the dicritical case.

It is clear that $\gcd(\hat{f}, \hat{g}) = 1$. On the other hand, $\widehat{f + cg}$ has the multiple factors y^2 for $c = 0$ and $(x + y)^2$ for $c = \infty$. Moreover $\hat{R} = 2(x + y)$. Thus from proposition 7 we obtain $W_{m-1} = W_1 = y$. From this computation we know that the differential system associated to H has multiplicity $m = 2$.

We construct as usual a table of desingularization. One only blow up is to be done in order to completely know the behavior of the singular point at the origin of the initial system. After the blow up $x = x_0 = x_1, y = y_0 = x_1 y_1$, the multiplicity at the origin is $m = 1$. Only $f = 0$, which is formed by two complex curves, crosses the origin. Therefore we have a center. The desingularization process is finished. Figure 3 shows how we get the phase portrait of the initial system.

□

To end this section we consider the following problem: given a finite set of analytic curves crossing the origin, $f_1 = 0, \dots, f_p = 0$, we want to construct a planar differential system having a generalized rational first integral and having these curves as solutions. Moreover




FIGURE 3. The desingularization of example 3.

we want to be able to fix *a priori* the behavior (elliptic, hyperbolic or parabolic) of the canonical regions defined by the curves when they meet at the origin.

To get this differential system we need to choose convenient integers n_1, \dots, n_p and to build a function $H = \prod_{i=1}^p f_i^{n_i}$ in such a way that the desired behavior between each pair of curves is obtained. We illustrate this idea with an example.

Example 4. Consider the algebraic curves $f_1(x, y) = y^2 - x^3 = 0$, $f_2(x, y) = -x + y - 2xy^2 = 0$, $f_3(x, y) = y^3 + x^5 = 0$ and $f_4(x, y) = 3x^2 + y^3 - 4x^3y^4 = 0$. We want to construct a system having a rational first integral such that these four curves determine a local behavior around the origin as in figure 4. We assume that the singular point is not dicritical. From this figure we know that f_2 and f_4 must be in a different level set than f_1 and f_3 (see the parabolic sectors), therefore we take $H = f/g = (f_1^{n_1} f_3^{n_3}) / (f_2^{n_2} f_4^{n_4})$, with $n_i \in \mathbb{N}$. We choose these numerator and denominator because the separatrices of the hyperbolic sectors must belong to the same level set, while those of a parabolic sector must belong to different level sets.




FIGURE 4. The local behavior around the origin of the differential system in example 4.

The polynomial differential system having this function as (rational) first integral has multiplicity 7 at the origin no matter the values of the $n_i \in \mathbb{N}$. In order to begin the application of our algorithm, first of all we do the change of variables $x \rightarrow x + y, y \rightarrow x - 2y$, as there are curves approaching the origin tangent to both axis. Let $n_0 := m_f - m_g = 2n_1 - n_2 + 3n_3 - 2n_4 \in \mathbb{Z}$. We take $n_0 \neq 0$ as the singular point is not dicritical. We construct table 4 as usual.

From the construction of table 4 we can study all the singular points appearing in the whole blow up process:

- (1) First blow up, $x_0 = x_1, y_0 = x_1 y_1$. As we want the singular point $(0, 0)$ in the (x_1, y_1) -plane to be a node in order to obtain an elliptic sector in region 16 (see figure 5), we take $n_0 > 0$.
 - $(0, 1/2)$: as $m = 2$, a new blow up is required.

- $(0, -1)$: as $m = 1$, both $f = 0$ and $g = 0$ pass through this point and $g = 0$ does it not transversally (because $n_0 > 0$), a new blow up is required.
- $(0, 0)$: as $m = 1$, both $f = 0$ and $g = 0$ pass through this point and $g = 0$ does it transversally (because $n_0 > 0$), it is a node.

The second and third blow ups concern the point $(0, -1)$. We move this point to the origin of the (x_1, y_1) -plane.

- (2) Second blow up, $x_1 = x_2y_2$, $y_1 = y_2$.
 - $(0, 0)$: as $m = 2$, a new blow up is required.
- (3) Third blow up, $x_2 = x_3y_3$, $y_2 = y_3$. Because of the configuration that we want to obtain we must take $m_f > m_g$. This implies that we must take $2n_0 > n_4$.
 - $(0, 0)$: as $m = 1$ and only $f = 0$ passes through this point, it is a saddle.
 - $(-1/9, 0)$: as $m = 1$, both $f = 0$ and $g = 0$ pass through this point and $g = 0$ does it transversally, it is a node.

The rest of blow ups concern the point $(0, 1/2)$ in the (x_1, y_1) -plane. We move this point to the origin.

- (4) Fourth blow up, $x_1 = \bar{x}_2\bar{y}_2$, $y_1 = \bar{y}_2$:
 - $(0, 0)$: as $m = 2$, a new blow up is required.

Before the next blow up, and as there are curves with vertical and horizontal tangent, we do the change of variable $\bar{x}_2 \rightarrow \bar{x}_2 + \bar{y}_2$.
- (5) Fifth blow up, $\bar{x}_2 = \bar{x}_3$, $\bar{y}_2 = \bar{x}_3\bar{y}_3$:
 - $(0, 0)$: as $m = 2$, a new blow up is required.
 - $(0, -1)$: as $m = 1$ and only $f = 0$ passes through this point, it is a saddle.
 - $(0, 27/5)$: as $m = 1$ and only $f = 0$ passes through this point, it is a saddle.
- (6) Sixth blow up, $\bar{x}_3 = \bar{x}_4\bar{y}_4$, $\bar{y}_3 = \bar{y}_4$:
 - $(0, 0)$: as $m = 1$ and only $f = 0$ passes through this point, it is a saddle.

	SP_f	SP_g	\hat{f}	\hat{g}
$y_0 = x_1y_1$	$y_1 = \frac{1}{2}$ ★ ★	★ $y_1 = -1$ $y_1 = 0$	$x_1^{n_0+n_1+2n_3}$ $x_1^{n_0}$ $x_1^{n_0}$	★ $x_1^{n_4}$ $y_1^{n_2}$
$y_1 \rightarrow y_1 - 1$ $x_1 = x_2y_2$	$x_2 = 0$	$x_2 = 0$	$x_2^{n_0}y_2^{n_0-n_4}$	$l_1^{n_4}$
$x_2 = x_3y_3$	$x_3 = 0$ ★	★ $x_3 = -\frac{1}{9}$	$x_3^{n_0}y_3^{2n_0-2n_4}$ $y_3^{2n_0-2n_4}$	★ $l_2^{n_4}$
$y_1 \rightarrow y_1 + \frac{1}{2}$ $x_1 = \bar{x}_2\bar{y}_2$	$\bar{x}_2 = 0$	★	$\bar{x}_2^*\bar{y}_2^*l_3^{n_1}$	★
$\bar{x}_2 \rightarrow \bar{x}_2 + \bar{y}_2$ $\bar{y}_2 = \bar{x}_3\bar{y}_3$	$\bar{y}_3 = 0$ $\bar{y}_3 = -1$ $\bar{y}_3 = \frac{27}{5}$	★ ★ ★	$\bar{x}_3^*\bar{y}_3^*l_4^{n_3}$ $\bar{x}_3^*\bar{y}_3^*$ $\bar{x}_3^*l_5^{n_1}$	★ ★ ★
$\bar{x}_3 = \bar{x}_4\bar{y}_4$	$\bar{y}_4 = 0$ $\bar{y}_4 = \frac{256}{243}$	★ ★	$\bar{x}_4^*\bar{y}_4^*$ $\bar{y}_4^*l_6^{n_3}$	★ ★

TABLE 4. Application of the algorithm in example 4. The l_i are straight lines crossing the corresponding singular point with neither horizontal nor vertical tangency. In particular $l_1 = 9x_2 + y_2$, $l_3 = 27\bar{x}_2 - 32\bar{y}_2$ and $l_4 = 243\bar{x}_3 - 256\bar{y}_3$. The powers of the factors of \hat{g} marked with a ★ are not relevant for the explanation. The polynomial \hat{R} in the first row is $x^{n_0+n_1+2n_3-3}$; in the other cases it can be obtained directly from the factors of \hat{f} and \hat{g} powered to their respective exponent minus one.




FIGURE 5. The desingularization of example 4.

- $(256/243, 0)$: as $m = 1$ and only $f = 0$ passes through this point, it is a saddle.

Now the desingularization process is done. To obtain the phase portrait of the initial system we must undo the changes of variables. As a conclusion, in order to ensure that we obtain the desired configuration, we must take $2n_0 > n_4$. A rational first integral is the function $H = (f_1^2 f_3^4)/(f_2 f_4^4)$. \square

From example 4 a natural question arises:

Open question. *Given a finite set of analytic curves crossing the origin and a local topological configuration around it, is it possible to find an analytic system having a generalized rational first integral and having a singular point at the origin with the given local topological behavior?*

Acknowledgement. The second author wants to thank Maria Alberich many comments that helped us to write this paper.

REFERENCES

- [1] M.J. ÁLVAREZ AND X. JARQUE, *A k-blow up algorithm and the Nilpotent Theorem*, to appear in Int. J. Bif. and Chaos.
- [2] A. ANDREEV, *Investigation on the behaviour of the integral curves of a system of two differential equations in the neighborhood of a singular point*, Trans. Amer. Math. Soc. **8**, (1958), 187–207.
- [3] A. A. ANDRONOV, E.A. LEONTOVICH, I.I. GORDON AND A.G. MAIER, *Qualitative theory of second order differential equations*, John Wiley and Sons, New York, 1973.
- [4] J. CHAVARRIGA, H. GIACOMINI, J. GINÉ AND J. LLIBRE, *Darboux integrability and the inverse integrating factor*, J. Diff. Eq. **194** (2003), 116–139.
- [5] J. CHAVARRIGA, H. GIACOMINI AND M. GRAU, *Necessary conditions for the existence of invariant algebraic curves for planar polynomial systems*, Bull. Sci. Math. **129** (2005) 99–126.
- [6] C. CHRISTOPHER, J. LLIBRE AND J.V. PEREIRA, *Multiplicity of invariant algebraic curves in polynomial vector fields*, Pac. J. Math. **229** (2007), 63–117.
- [7] F. DUMORTIER, *Singularities of vector fields on the plane*, J. Diff. Eq., **23** (1977), 53–106.
- [8] F. DUMORTIER AND C. HERSSENS, *Tracing phase portraits of planar polynomial vector fields with detailed analysis of the singularities*, Qual. Theory of Dyn. Systems, **1** (1999), 97–130.
- [9] A. FERRAGUT AND H. GIACOMINI, *A new algorithm for finding rational first integrals of polynomial vector fields*, preprint 2009.
- [10] A. FERRAGUT AND J. LLIBRE, *On the remarkable values of the rational first integrals of polynomial vector fields*, J. Diff. Eq. **241**, (2007), 399–417.
- [11] WEIGU LI, J. LLIBRE AND XIANG ZHANG, *Planar analytic vector fields with generalized rational first integrals*. Bull. Sci. Math. **125** (2001), 341–361.
- [12] H. POINCARÉ, *Sur l'intégration des équations différentielles du premier ordre et du premier degré I and II*, Rend. Circolo Matematico di Palermo **5** (1891), 161–191; **11** (1897), 193–239.
- [13] A. SEIDENBERG, *Reduction of singularities of the differential equation $Ady = Bdx$* , Amer. J. Math. **90** (1968), 248–269.
- [14] S. WALCHER, *On the Poincaré problem*, J. Diff. Eq. **166** (2000) 51–78.

¹ DEPARTAMENT DE MATEMÀTIQUES I INFORMÀTICA, UNIVERSITAT DE LES ILLES BALEARS, EDIFICI ANSELM TURMEDA, CTRA. DE VALLDEMOSSA, KM 7.5, 07122 PALMA, ILLES BALEARS-SPAIN

² DEPARTAMENT DE MATEMÀTICA APLICADA I, UNIVERSITAT POLITÈCNICA DE CATALUNYA, AV. DIAGONAL, 647, 08028 BARCELONA, CATALONIA-SPAIN
E-mail address: chus.alvarez@uib.es, antoni.ferragut@upc.edu