Dear Author

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style.
- Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

<http://dx.doi.org/10.1007/s10113-018-1443-2>

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

1	Article Title	Evaluation of the representativeness and the causes of variability of atmospheric carbon dioxide observations at two sites of the Aigüestortes Long-Term Ecological Research node in Central Pyrenees, Spain	
2	Article Sub- Title		
3	Article Copyright - Year	Springer-Verlag GmbH Germany, part of Springer Nature 2018 (This will be the copyright line in the final PDF)	
4	Journal Name	Regional Environmental Change	
5		Family Name	Curcoll
6		Particle	
7		Given Name	Roger
8		Suffix	
9		Organization	Institut de Ciència i Tecnologia Ambientals, ICTA-UAB
10		Division	
11		Address	ICTA-ICP Building Z Campus UAB, Bellaterra (Cerdanyola), Barcelona 08193, Spain
12		Organization	Institut Català de Ciències del Clima, IC3
13	Corresponding Author	Division	
14		Address	Barcelona, Spain
15		Organization	Associació LTER-Aigüestortes
16		Division	
17		Address	Barcelona, Spain
18		Organization	Universitat Politècnica de Catalunya (UPC)
19		Division	Departament d'Enginyeria Química
20		Address	Barcelona, Spain
21		e-mail	roger.curcoll@uab.cat
22	Author	Family Name	Camarero
23		Particle	
24		Given Name	Lluís
25		Suffix	
26		Organization	Associació LTER-Aigüestortes
27		Division	
28		Address	Barcelona, Spain
29		Organization	Centre d'Estudis Avançats de Blanes, CEAB-CSIC

30		Division	
31		Address	Girona, Spain
32		e-mail	camarero@ceab.csic.es
33	Author	Family Name	Bacardit
34		Particle	
35		Given Name	Montse
36		Suffix	
37		Organization	Centre de Lauegi, Conselh Generau d’ Aran
38		Division	
39		Address	Lleida, Spain
40		e-mail	m.bacardit@aran.org
41	Author	Family Name	Àgueda
42		Particle	
43		Given Name	Alba
44		Suffix	
45		Organization	Institut Català de Ciències del Clima, IC3
46		Division	
47		Address	Barcelona, Spain
48		Organization	Universitat Politècnica de Catalunya (UPC)
49		Division	Departament d’Enginyeria Química
50		Address	Barcelona, Spain
51		e-mail	alba.agueda@upc.edu
52	Author	Family Name	Grossi
53		Particle	
54		Given Name	Claudia
55		Suffix	
56		Organization	Institut Català de Ciències del Clima, IC3
57		Division	
58		Address	Barcelona, Spain
59		Organization	Institut de Tècniques Energètiques (INTE), UPC
60		Division	
61		Address	Barcelona, Spain
62		Organization	Departament de Física Nuclear, UPC
63		Division	
64		Address	Barcelona, Spain
65		e-mail	claudia.grossi@upc.edu
66	Author	Family Name	Gacia
67		Particle	

68		Given Name	Esperança
69		Suffix	
70		Organization	Associació LTER-Aigüestortes
71		Division	
72		Address	Barcelona, Spain
73		Organization	Centre d’Estudis Avançats de Blanes, CEAB-CSIC
74		Division	
75		Address	Girona, Spain
76		e-mail	gacia@ceab.csic.es
77		Family Name	Font
78		Particle	
79		Given Name	Anna
80		Suffix	
81	Author	Organization	King’s College London
82		Division	Environmental Research Group, MRC PHE Centre for Environment and Health
83		Address	London, UK
84		e-mail	anna.font_font@kcl.ac.uk
85		Family Name	Morguí
86		Particle	
87		Given Name	Josep-Anton
88		Suffix	
89		Organization	Institut Català de Ciències del Clima, IC3
90		Division	
91		Address	Barcelona, Spain
92		Organization	Institut de Ciència i Tecnologia Ambientals, ICTA-UAB
93		Division	
94	Author	Address	ICTA-ICP Building Z Campus UAB, Bellaterra (Cerdanyola), Barcelona 08193, Spain
95		Organization	Associació LTER-Aigüestortes
96		Division	
97		Address	Barcelona, Spain
98		Organization	Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals (BEECA) UB
99		Division	
100		Address	Barcelona, Spain
101		e-mail	JosepAnton.Morgui@uab.cat
102	Schedule	Received	25 September 2017
103		Revised	

104		Accepted	20 November 2018
105	Abstract	<p>In order to improve the understanding of the carbon cycle in the Pyrenean region, two atmospheric monitoring mountain stations were set up within the Long-Term Ecological Research node of Aigüestortes i Estany de Sant Maurici at Central Pyrenees, Spain. The atmospheric concentration of carbon dioxide (CO₂) was measured over 2008–2014 and 2010–2014 at Estany Llong (ELL) site and Centre de Recerca d’Alta Muntanya (CRAM), respectively. Measurements were carried out fortnightly off-line with high precision instrumentation at ELL and every minute online with a lower precision sensor at CRAM in conjunction with meteorological variables. The two datasets were analyzed in this study, quantifying whenever possible annual growth rates (AGR), seasonal variability, and diurnal amplitudes. Results were also compared with the NOAA Marine Boundary Layer (MBL) reference product and CO₂ data from other background monitoring stations. Four-harmonics adjusted CO₂ data from ELL showed a high correlation with the NOAA MBL reference product for the same latitude (Spearman’s rho $\rho = 0.96$). In addition, AGRs of CO₂ at ELL correlated well with those observed at Mace Head (MHD) station ($\rho = 0.94$), suggesting that ELL can be considered a background station. Winter CRAM CO₂ data was not statistically different from ELL data, while in summer, it was 5.5 ppm lower on average, suggesting a higher photosynthesis uptake. The amplitude of the CO₂ diurnal cycle at CRAM was found to be exponentially related to the local mean daily temperature and dependent on forthcoming wind sector (N-NW or E-SE-S-SW). An increase in CRAM CO₂ concentrations was observed under N-NW winds during daytime, which could be related to traffic emissions. This study demonstrates that the use of CO₂ sensors with low precision but continuously corrected and periodically calibrated can be used for the study of local and regional CO₂ sources and sinks.</p>	
106	Keywords separated by ' - '	CO ₂ - Mountain site - LTER - Long-Term Ecological Research - Atmospheric measurements - Pyrenees	
107	Foot note information	<p>The online version of this article (https://doi.org/10.1007/s10113-018-1443-2) contains supplementary material, which is available to authorized users.</p> <p>Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.</p>	

Electronic supplementary material

Figure S1
General location of Aigüestortes Long Term Ecological Research site in the Pyrenees chain (left). Location of Centre de Recerca d’Alta Muntanya (CRAM) and Estany Llong (ELL) stations in the park (top-center). Corine land cover maps from the cartography of habitats in Catalonia (Carreras and Diego 2007) around CRAM (top left) and ELL (bottom left). (PNG 14442 kb)

Figure S2

Schematic for the CO₂ analysis of flask samples using two Licor7000 analyzers. (PDF 15 kb)

Figure S3

Diurnal and annual CO₂ variability at Centre de Recerca d'Alta Muntanya (CRAM). Trend level plot for CO₂ at CRAM for 2010, 2011, 2012 and 2013. (PDF 14 kb)

Evaluation of the representativeness and the causes of variability of atmospheric carbon dioxide observations at two sites of the Aigüestortes Long-Term Ecological Research node in Central Pyrenees, Spain

Roger Curcoll^{1,2,3,4} · Lluís Camarero^{3,5} · Montse Bacardit⁶ · Alba Àgueda^{1,4} · Claudia Grossi^{1,7,8} · Esperança Gacia^{3,5} · Anna Font⁹ · Josep-Anton Morguí^{1,2,3,10}

Received: 25 September 2017 / Accepted: 20 November 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

In order to improve the understanding of the carbon cycle in the Pyrenean region, two atmospheric monitoring mountain stations were set up within the Long-Term Ecological Research node of Aigüestortes i Estany de Sant Maurici at Central Pyrenees, Spain. The atmospheric concentration of carbon dioxide (CO₂) was measured over 2008–2014 and 2010–2014 at Estany Llong (ELL) site and Centre de Recerca d'Alta Muntanya (CRAM), respectively. Measurements were carried out fortnightly off-line with high precision instrumentation at ELL and every minute online with a lower precision sensor at CRAM in conjunction with meteorological variables. The two datasets were analyzed in this study, quantifying whenever possible annual growth rates (AGR), seasonal variability, and diurnal amplitudes. Results were also compared with the NOAA Marine Boundary Layer (MBL) reference product and CO₂ data from other background monitoring stations. Four-harmonics adjusted CO₂ data from ELL showed a high correlation with the NOAA MBL reference product for the same latitude (Spearman's rho $\rho = 0.96$). In addition, AGRs of CO₂ at ELL correlated well with those observed at Mace Head (MHD) station ($\rho = 0.94$), suggesting that ELL can be considered a background station. Winter CRAM CO₂ data was not statistically different from ELL data, while in summer, it was 5.5 ppm lower on average, suggesting a higher photosynthesis uptake. The amplitude of the CO₂ diurnal cycle at CRAM was

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10113-018-1443-2>) contains supplementary material, which is available to authorized users.

✉ Roger Curcoll
roger.curcoll@uab.cat

Lluís Camarero
camarero@ceab.csic.es

Montse Bacardit
m.bacardit@aran.org

Alba Àgueda
alba.agueda@upc.edu

Claudia Grossi
claudia.grossi@upc.edu

Esperança Gacia
gacia@ceab.csic.es

Anna Font
anna.font_font@kcl.ac.uk

José-Anton Morguí
JosepAnton.Morgui@uab.cat

¹ Institut Català de Ciències del Clima, IC3, Barcelona, Spain

² Institut de Ciència i Tecnologia Ambientals, ICTA-UAB, ICTA-ICP Building Z Campus UAB, Bellaterra (Cerdanyola), 08193 Barcelona, Spain

³ Associació LTER-Aigüestortes, Barcelona, Spain

⁴ Departament d'Enginyeria Química, Universitat Politècnica de Catalunya (UPC), Barcelona, Spain

⁵ Centre d'Estudis Avançats de Blanes, CEAB-CSIC, Girona, Spain

⁶ Centre de Lauegi, Conselh Generau d'Aran, Lleida, Spain

⁷ Institut de Tècniques Energètiques (INTE), UPC, Barcelona, Spain

⁸ Departament de Física Nuclear, UPC, Barcelona, Spain

⁹ Environmental Research Group, MRC PHE Centre for Environment and Health, King's College London, London, UK

¹⁰ Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals (BEECA) UB, Barcelona, Spain

27 found to be exponentially related to the local mean daily temperature and dependent on forthcoming wind sector (N-NW or E-SE-
 28 S-SW). An increase in CRAM CO₂ concentrations was observed under N-NW winds during daytime, which could be related to
 29 traffic emissions. This study demonstrates that the use of CO₂ sensors with low precision but continuously corrected and
 30 periodically calibrated can be used for the study of local and regional CO₂ sources and sinks.

31 **Keywords** CO₂ · Mountain site · LTER · Long-Term Ecological Research · Atmospheric measurements · Pyrenees

32 Introduction 33

34 Accurate assessment of CO₂ emissions and their redistribution
 35 among the atmosphere, ocean, and terrestrial biosphere are
 36 important to better understand the global carbon cycle, to sup-
 37 port the development of climate policies, and to mitigate and
 38 tackle climate change (Bouwman 1989; Post et al. 1990; Tans
 39 et al. 1990a; Le Quéré et al. 2016).

40 The distribution of CO₂ sources and sinks is poorly
 41 constrained at regional (10²–10⁶ km²) and seasonal scales
 42 (Ciais et al. 2000; Geels et al. 2007; Bergamaschi et al.
 43 2010; Hase et al. 2015; Hu et al. 2018). Top-down techniques
 44 are useful to estimate carbon fluxes from atmospheric CO₂
 45 observations (Ciais et al. 2010). These top-down inversion
 46 models are more consistent for regions where the atmospheric
 47 network is dense (Peylin et al. 2013). Unfortunately, the num-
 48 ber of measurements and their spatial coverage are still insuf-
 49 ficient in some European regions such as the NW of the
 50 Mediterranean (Marquis and Tans 2008).

51 Remote mountain stations, located far away from anthropo-
 52 genic sources, can provide useful information to investigate
 53 trends of large-scale background levels (Ciattaglia et al. 1987;
 54 Colombo et al. 2000; Schmidt 2003; Schibig et al. 2015;
 55 Bamberger et al. 2017). When local influences are filtered,
 56 mountain greenhouse gas (GHG) time series can be considered
 57 suitable for global inversion models (Levin 1987; Bamberger
 58 et al. 2017). Monitoring GHGs in mountainous remote sites is
 59 also useful to understand the variability of the response of the
 60 mountain ecosystems to global warming (e.g., Randerson et al.
 61 1999; Bakwin et al. 2004; Angert et al. 2005). Moreover, when
 62 the inversion models focus on small scales and utilize continental
 63 measurements of CO₂ they can better resolve the responses of
 64 various vegetation types and the impact of human interventions
 65 on land-atmosphere fluxes (Pillai et al. 2011).

66 In order to supply to the lack of CO₂ data in the NW
 67 Mediterranean region, atmospheric measurements were car-
 68 ried out since 2008 at Estany Llong (ELL), a mountain site
 69 in Central Pyrenees (north Spain). In 2010, continuous CO₂
 70 measurements started at the nearby site of Centre de Recerca
 71 d'Alta Muntanya (CRAM). The two sites are part of the Long-
 72 Term Ecological Research (LTER) node of the “Aigüestortes i
 73 Estany de Sant Maurici National Park” (ANP). The ANP has
 74 been a node of scientific research for hydrology (Catalan
 75 1989), atmospheric chemistry (Bacardit and Camarero 2009;

Curcoll et al. 2010; Camarero et al. 2017), atmospheric biol-
 ogy (Hervàs et al. 2009), biogeochemistry of surface waters
 (Camarero and Catalan 1993), aquatic organisms (Chappuis
 et al. 2011, 2014), and terrestrial flora and fauna (Carrillo and
 Ninot 1992).

The present study aims to assess the suitability of two
 Central Pyrenees GHG atmospheric stations for regional scale
 CO₂ monitoring and to evaluate the influencing factors driv-
 ing the observed CO₂ variability at different temporal scales.
 More specifically, in this study, ELL and CRAM CO₂ time
 series are presented for the first time and they are statistically
 analyzed with two specific goals: (i) to offer the data to the
 scientific community and discuss their spatial representative-
 ness; (ii) to determine the sources of variability and to differ-
 entiate anthropogenic from natural influences.

After a short description of the sites, the sampling and
 analyzing methodology is addressed, remarking the precision
 and accuracy of the analytical procedure. Daily, monthly, and
 annual variabilities of the two datasets are examined whenever
 possible, depending on their temporal resolution. To better
 interpret our results, we compare the obtained annual growth
 rates and seasonal variabilities with those from other
 European sites (Monte Cimone, Italy; Mace Head, UK; and
 CIBA, Spain). We also examine the influence of wind direc-
 tion and ambient temperature. Finally, we use the results of
 these two sites (ELL and CRAM) to discuss whether the use
 of low-precision sensors can be justified in areas with low
 maintenance requirements.

Methods

Site description

In 2008, LTER-Aigüestortes node was established as a member
 of the LTER-Spain network (Camarero and Morgui 2009). In the
 same year, a continuous atmospheric CO₂ monitoring program
 was established at the eastern extreme of Estany Llong Lake
 (ELL). The remote sampling site of ELL (42.57 N, 0.95 E,
 2003 m above sea level [m asl]) is located inside the ANP. The
 ANP (14,119 ha) and its peripheral protection area (26,733 ha)
 are located in the southern axial chain of the Central Pyrenees, in
 northeastern Spain. It is a high-mountain area and its altitude
 ranges between 1383 and 3029 m asl. The climate is Atlantic

mountain-type (Albentosa Sánchez 1973), with strong winters and snow present from November to May. Precipitation presents annual accumulated values of 1200–1300 mm. The annual mean temperature is 5.2 °C with the highest monthly average in July (19.3 °C) and the lowest in February (− 4.3 °C) (Ninyerola et al. 2003). The location is actually a mountain meadow of *Festuca* sp. and *Nardus* sp. surrounded by granite mountains and coniferous forest (Carreras and Diego 2007). Valley and mountain breezes are common at this site with upslope winds during the day and downslope breezes at night (Camarero et al. 2018).

In 2010, continuous atmospheric CO₂ measurements started at CRAM (42.62 N, 0.77 E, 1584 m asl). The CRAM is located at the top of the Noguera Ribagorçana valley that has a North-South orientation. The valley is located 16 km west of the ELL remote site, in the peripheral area of the ANP. The CRAM station is located 200 m south of the Vielha road tunnel entrance (5230 m long). It is surrounded by grassland in the North and mixed, deciduous, and coniferous forests in the South (Carreras and Diego 2007). A map with the location of both sites with Corine land cover cartography can be found in the support material (Fig. S1).

Estany Llong sampling and analytical method

Fortnightly samplings were carried out at ELL since April 2008. Samples were collected in two 1-L cylindrical borosilicate glass flasks (Normag Labor und Prozesstechnik GmbH, Germany) with Kel-F PCTFE valves fitted at both ends. This material shows the lowest permeation of gases compared to other sealing materials (Sturm et al. 2004). Flasks were sampled using a new developed self-made sampling device suitable for remote sites. Air was taken at 3.5 m above ground level (m agl) and was dried by passing through a desiccant (magnesium perchlorate). Then, the sampled air was filtered to remove particles and pumped to the flasks. Air was flushed through a pair of flasks for at least 10 min at a flow rate between 2.0 and 3.5 L min^{−1}. After, the exit valve was closed and flasks were pressurized with sampled air up to 1.5 bars. Air samples were collected between 10 and 12 GMT, when the mixed boundary layer was developed and the valley breeze was established. These previous conditions were selected to sample the widest possible influence area. Air temperature, relative humidity, atmospheric pressure, and wind direction and speed were measured using a portable Kestrel 5500 weather device during ELL flask sampling. According to factory specifications, accuracy of the instrument is 0.5 °C (temperature), 2% (relative humidity), 1.5 mbar (pressure), 5° (wind direction), and 3% (wind speed).

Flask samples from ELL site were analyzed for CO₂ using two differential, non-dispersive infrared gas analyzers (Licor7000) in series. Each Licor7000 had two cells and two detectors for a simultaneous measurement of CO₂ and H₂O vapor on a reference and a sample stream. A vacuum pump was

installed to empty the backspace between the flask valve and the analyzer inlet. A sketch of the analyzing instrumental configuration can be found in the support material (Fig. S2). Both sample and reference were flushed through the cells at a flow rate of 0.1 L min^{−1} controlled by two Bronkhorst mass flow controllers. On the sample line, a Valco multivalve was used to switch between calibration standards, samples, and reference gas. Reference gas passed through the sample cell before and after each flask analysis. The value of the reference gas in the sample cell was used as a baseline to cope with instrument drift.

The Licor7000 has a strong dependency on cell pressure and temperature (Gomez-Pelaez and Ramos 2011). To minimize the pressure influence, both Licor7000 were modified and the pressure sensors were removed and two additional Bronkhorst backpressure controllers at the end of the reference and sample lines were installed to keep the pressure stable inside the cells. Since 2010, the room temperature in the laboratory was controlled with a specific air-conditioning equipment keeping the temperature stable at 23 ± 0.3 °C. In 2013, both Licor7000 analyzers were installed in a Boxcult box and the cell temperature of instruments was stabilized at 30 ± 0.1 °C. Improvements made on temperature control reduced the error and the drift of both analyzers. Average absolute difference between CO₂ calibrated measures of both analyzers decreased through the years, being 0.104 ppm in the 2008–2009 period, 0.041 in 2010–2012, and 0.029 in 2013–2015. The CO₂ values used in this study are the mean calculated from both analyzers. The use of the two Licor7000 analyzers in series improved the precision of the measurement and was useful for tracking instrument drift.

Each day that a set of flasks had to be analyzed in the lab, the two Licor7000 were calibrated every 4 h with four secondary standard cylinders. These secondary standards were recalibrated every 6 months with 6 NOAA standards (WMO X2007 scale; Zhao and Tans 2006). Each flask was analyzed for 7 min, time that best compromised the stability of the measurements and the lack of flow and pressure drop. For each measurement, only the last 30 s was considered for calculation. The two flasks of each sampling were analyzed on different days to cope with the instrument variability and differences due to calibrations. Uncertainties in the measured concentrations stemmed from both the sampling method and the analysis (Tans et al. 1990b). Pairs of flasks that had a difference higher than 0.7 ppm of CO₂ between both calibrated values were discarded. The average difference between the two flasks of each pair was − 0.03 (1σ ± 0.3) ppm. This deviation was consistent with other CO₂ flask measurements (e.g., Conway et al. 1994; Andrews et al. 2014).

Continuous sampling at CRAM

A Vaisala GMP343 carbon dioxide probe was installed for continuous CO₂ measurements in 2010 at CRAM station. Air inlet

218 was at 3 m agl and the probe was protected from direct radiation
 219 and rain. The GMP343 sensor is an infrared absorbance CO₂
 220 detector that uses a micromachined electrically tunable Fabry-
 221 Perot Interferometer (FPI). The FPI enables a reference measure-
 222 ment at a wavelength where no absorption occurs to improve the
 223 measurement (Vaisala 2017). The GMP343 has been used in
 224 atmospheric and biospheric studies where very high precision
 225 is not required (Pumpanen et al. 2008; Rigby et al. 2008).
 226 Factory precision of the instrument was 1% with a minimum
 227 long-term drift of ± 10 ppm CO₂ year⁻¹. A Vantage Pro2 Davis
 228 weather station that delivered temperature, pressure, relative hu-
 229 midity, and wind direction and wind speed data was also installed
 230 at CRAM. The GMP343 has an internal correction algorithm
 231 that considers temperature, humidity, and pressure. Pressure
 232 and relative humidity from the meteorological station were sent
 233 to the instrument via RS232 every 10 min to correct CO₂ data.
 234 Correction for temperature was made using the internal GMP343
 235 temperature sensor.

236 The instrument was calibrated at least twice a year by com-
 237 paring concentrations with a collocated Picarro G2301 Cavity
 238 RingDown Spectroscopy Analyzer (CRDS), this latter with a
 239 precision better than 0.03 ppm for CO₂ (Crosson 2008;
 240 Richardson et al. 2012). In the inter-comparison events, the
 241 G2301 analyzer was collocated for at least 1 week. G2301
 242 results were corrected for water vapor (Rella et al. 2013) and
 243 further calibrated in the laboratory using six NOAA reference
 244 gases before and after each inter-comparison. Residuals of the
 245 linear fit between CO₂ values measured by GMP343 and CO₂
 246 values measured by G2301 showed a strong dependence on
 247 temperature. Therefore, an additional correction for tempera-
 248 ture was implemented. The final CO₂ concentrations mea-
 249 sured by GMP343 were obtained by linear regression of mea-
 250 sured CO₂ and temperature as expressed in Eq. 1, where T_{GMP}
 251 is the temperature measured by the GMP343, $\text{CO}_{2(\text{GMP})}$ is the
 252 value of the GMP343, and $\text{CO}_{2(\text{corr})}$ the corrected CO₂ value,
 253 measured by the G2301 analyzer. After applying the addition-
 254 al temperature correction, residual standard error (RSE) for
 255 CO₂ calibration events was 1.5 ppm (1 week).

$$\text{CO}_{2(\text{corr})} = A + B \cdot \text{CO}_{2(\text{GMP})} + C \cdot T_{\text{GMP}} \quad (1)$$

258
 259 A strong linear drift was observed on GMP343 data be-
 260 tween calibrations (~ 15 ppm year⁻¹), probably caused by
 261 the soiling of the mirrors of the sensor. A linear scaling was
 262 applied to CO₂ time series to correct for drift.

263 Data analysis

264 CO₂ time series curve fitting

265 The Carbon Cycle Group CuRve (CCGCRV) is a digital time
 266 series filtering algorithm developed by Kirk Thoning within
 267 the Carbon Cycle Group, Earth System Research Laboratory

at NOAA (USA) in the late 1980s (Thoning et al. 1989).
 Approximations of the seasonal cycle and the long-term trend
 are made by fitting a polynomial equation combined with a
 harmonic function. CCGCRV fit is recommended for analysis
 of time series where it is important to accurately represent the
 magnitude of the seasonal cycle or the timing of the seasonal
 inflection points and for analyses where the year-to-year var-
 iation in the seasonal and trend components is retained
 (Pickers and Manning 2015). We have applied the
 CCGCRV to the ELL CO₂ series using the recommended
 parameters by Pickers and Manning (2015): number of yearly
 harmonics = 4, number of polynomial terms = 3, short-term
 cutoff = 50 days, long-term cutoff = 667 days.

Bivariate polar plots for wind sector analysis

The location of potential local sources and sinks at both ELL and
 CRAM was explored through bivariate polar plots (Carslaw and
 Ropkins 2012). The ELL bivariate polar plot was composed with
 the residuals of the ELL CO₂ series from the four-harmonics
 CCGCRV fit curve instead of using the actual concentrations.
 By doing so, we remove the seasonal and the inter-annual vari-
 abilities. For the CRAM dataset, bivariate polar plots for absolute
 CO₂ concentrations were generated.

Ambient temperature influence on the diurnal amplitude

Due to the measurement error of the GMP343 sensors being
 greater than 1 ppm, the time series at CRAM was only used
 for the analysis of large CO₂ variations (diurnal and seasonal
 cycles) but not for the study of long-term trends.
 Meteorological variables from the Davis station were also used.
 Hourly means were calculated from minute measurements. For
 all the measurement days at CRAM, the amplitude of the CO₂
 diurnal cycle was calculated as the difference between the daily
 maximum concentration and the daily minimum.

To evaluate the influence of air temperature on the diurnal
 CO₂ cycle, the Lloyd and Taylor formulation was used. It is a
 semi-empirical formulation that relates the temperature depen-
 dency of soil respiration, reflecting the decrease in the respi-
 ration activation energy with increasing temperature (Lloyd
 and Taylor 1994) (Eq. 2). Although this equation was formu-
 lated for soil respiration (R , $\mu\text{mol C m}^{-2} \text{ s}^{-1}$), we consider that
 the amplitude of the CO₂ diurnal cycle at CRAM is dependent
 on the soil respiration and ecosystem photosynthesis, both
 controlled by temperature (Berry and Bjorkman 1980; Smith
 and Dukes 2013). Therefore, in our approximation, the vari-
 able R refers to the amplitude of the diurnal cycle (ppm CO₂).

$$R = Ae^{\frac{-E_0}{T-T_0}} \quad (2)$$

While the regression parameter T_0 is kept constant at $-$
 46.02 °C (Lloyd and Taylor 1994), the activation energy kind

of parameter (E_0) is a free parameter that sets the temperature sensitivity (Reichstein et al. 2005). T is the average diurnal temperature and A is a dataset-dependent variable.

Data retrieval from NOAA Marine Boundary Layer, background monitoring stations, and global flux models for data comparison

ELL CO₂ time series were compared with the NOAA CO₂ Marine Boundary Layer (MBL) reference product for the same latitude (Dlugokencky et al. 2017b). The NOAA MBL reference product is derived directly from measurements of weekly air samples and gives low-noise representation of the annual variability and the ongoing global increase of CO₂ (Masarie and Tans 1995). The zonal product of the NOAA MBL gives the marine background concentration for a range of latitudes (42.57–42.58° north in our case). Each value has an associated uncertainty assigned by the density of measurements on a given period. The average associated error for the 2008–2014 period was 0.61 ppm. The CCGCRV curve was also applied to the MBL data series for a better statistical analysis.

To evaluate ELL as a mountain site capturing long-term changes, we have retrieved data from nearby stations that measure CO₂ with weekly flask measurements. Selected stations were as follows: (1) Monte Cimone (CMN; 44.18 N, 10.70 E, 2165 m asl), a mountain station at similar latitude and altitude (Ciattaglia 1983; Ciattaglia et al. 1987; Cundari et al. 1990; Cundari et al. 1995; Cristofanelli et al. 2018); (2) Mace Head (MHD; 53.33 N, 9.90 W, 5 m asl), located in the west Atlantic coast of Ireland and considered in many studies as a reference for Europe (Conway et al. 1994; Dlugokencky et al. 2017a); (3) CIBA (CIB; 41.81 N, 4.93 W, 856 m asl), a northern-Spain station located in an agricultural area at a similar latitude as ELL (Sánchez et al. 2010; Dlugokencky et al. 2017a). Data from these stations have been obtained via NOAA ObsPack products (Cooperative Global Atmospheric Data Integration Project 2013, 2017). The growth rate for these stations was also obtained using the CCGCRV algorithm.

Modeled values of annual terrestrial flux from two global models, MACC-V2 (Chevallier et al. 2010) and JENA-MPI (Rödenbeck et al. 2003), were obtained from the Global Carbon Atlas (<http://www.globalcarbonatlas.org/en/flux-time-series>).

Results

Estany Llong data series results

Estany Llong annual growth rates

The CCGCRV analysis was applied at ELL CO₂ data series with a good agreement (residual standard error = 1.71 ppm)

(Fig. 1). Signals of strong systematic anthropogenic or natural influences of samples seem not to be present in the ELL CO₂ data series. When comparing the harmonic function of ELL with the MBL reference, annual cycle of both series had a similar pattern and amplitude, with a slightly earlier spring decrease of the CO₂ concentration at ELL. The non-parametric Spearman's rank coefficient was used to analyze the correlation between series. Spearman's rho (ρ) correlation between the two whole series was 0.96, and ELL CO₂ concentrations were 0.89 ± 1.57 (1 σ) ppm lower than the MBL reference over the period 2008–2013. Thus, the two time series do not differ. Moreover, we must take into account the average associated error for the NOAA MBL series, 0.61 ppm (Dlugokencky et al. 2017b). This result suggests that ELL is mainly capturing a background signal.

From the four-harmonic function, the annual growth was obtained. The variability in the annual growth rate of both ELL and NOAA MBL CO₂ series followed a similar tendency ($\rho = 0.77$), with the highest value in 2010 (2.76 and 3.22 ppm CO₂/year, respectively) and the lowest in 2009 for ELL (1.11 ppm) and in 2011 for NOAA MBL (1.48 ppm CO₂/year). Average annual growth rate for the period 2008–2013 was 2.18 ± 0.74 (1 σ) ppm CO₂/year for ELL and 2.20 ± 0.65 (1 σ) ppm CO₂/year for NOAA MBL, showing that ELL CO₂ concentrations were growing with a similar trend as for the marine background concentrations (Fig. 2).

The highest correlation between annual growth rate at ELL against the other monitoring stations was with Mace Head ($\rho = 0.94$) (Atlantic coastal site) (Fig. 2). The other mountain station, Monte Cimone, showed a different behavior, with higher annual variability.

Fig. 1 Observed CO₂ time series from 2008 to 2014 based on Estany Llong (ELL) fortnightly CO₂ data series (red points), ELL CO₂ four-harmonics fit (straight line), NOAA Marine Boundary Layer (MBL) reference (dashed line), and the Centre de Recerca d'Alta Muntanya (CRAM) hourly CO₂ data series (green small dots)

Fig. 2 Observed CO₂ annual growth rate of Estany Llong (ELL), NOAA Marine Boundary Layer (MBL) reference, Mace Head (MHD), CIBA (CIB), and Monte Cimone (CMN) data series. Annual global CO₂ terrestrial flux as MACC V2 and JENA MPI models is shown for years 2008–2011 on the right axis. Vertical lines are the standard deviations of annual growth rates

The non-parametric Mann-Whitney-Wilcoxon test was used to statistically find the differences between the CO₂ concentrations from the different sectors. An average difference of 4.7 ppm was found between N-NW and E-SE-S-SW winds for the period 2011–2013 (95% confidence interval (CI) 4.1 to 5.3, $p < 10^{-15}$) (Fig. 4a). The observed differences were magnified in daytime, when the photosynthetic activity is higher, the up-valley wind is more frequent, and the transit of vehicles is higher (4.9 ppm, 4.3 to 5.3 CI, $p < 10^{-15}$) (Fig. 4b). In nighttime, when traffic was significantly reduced, the CO₂ concentrations were mostly driven by respiration processes and the accumulation in the lower boundary layer and there was no significant difference between the concentrations of the two directions ($p > 0.5$) (Fig. 4c). These differences in concentration between N-NW and E-SE-S-SW winds were also greater in summer (5.0 ppm, 4.5 to 5.5 CI, $p < 10^{-15}$) (Fig. 4d), when photosynthetic processes decreased CO₂ concentration, and were less significant in winter (1.2 ppm, 0.8 to 1.5 CI, $p < 10^{-9}$) (Fig. 4e).

Estany Llong wind direction influence

The wind rose for ELL (Fig. 3a) shows that main wind direction followed the valley orientation (NE-E and SW-W). The bivariate polar plot generated from the residuals of the four-harmonics fitting curve (Fig. 3b) shows that on average, there was no difference in CO₂ concentration when the wind blew from the valley (SW-W) or the mountain top (NE-E). Other forthcoming directions showed higher (SE) and lower (NW) signals, but the amount of data was too scarce to generate consistent conclusions.

Centre de Recerca d'Alta Muntanya data series results

Centre de Recerca d'Alta Muntanya diurnal cycle

The diurnal cycle of CO₂ concentration at CRAM shows a significant variability depending on the season (Fig. S3). The highest daily maximum concentration was measured in February and the lowest in August. In summer, when ecosystem respiration and photosynthesis at northern mid-latitudes get the maximum rates (Potter and Randerson 1993), the diurnal amplitude was higher, with average values of 31.4 ± 10.5 (1 σ) ppm. In winter months, the average diurnal amplitude was reduced to 7.5 ± 4.5 (1 σ) ppm. The average amplitude of the annual cycle of daily means in CRAM was 9.1 ppm.

Centre de Recerca d'Alta Muntanya wind direction influence

The influence of the air masses origin in the CRAM CO₂ concentration is shown in a bivariate polar plot in Fig. 4a.

Centre de Recerca d'Alta Muntanya temperature influence

The influence of temperature on the diurnal cycle amplitude in the CRAM CO₂ was analyzed. Since up to 20 ppm difference was observed depending on the wind forthcoming direction, data series was split depending on the predominant wind direction over a day: North to North-West sector (N-NW) and East to South-Westerly (E-SE-S-SW). The natural logarithm of the diurnal CO₂ amplitude was plotted against the ambient temperature for both wind sectors, and the Lloyd and Taylor equation was used to fit the values (Fig. 5). Resulting parameters of Eq. 2 are shown in Table 1. The diurnal CO₂ amplitude at the average daily temperature at CRAM (8.1 °C) for both N-NW and E-SE-S-SW wind sectors was 13.3 and 9.0 ppm, respectively. The non-parametric Kruskal-Wallis test was used to evaluate if the wind direction was a good explanatory variable of the amplitude of the diurnal cycle. Test result indicated that the diurnal amplitudes for air coming from both sectors were statistically different ($p < 10^{-15}$). A non-parametric analysis of covariance (Young and Bowman 1995; Bowman and Azzalini 1997) showed that the effect of the temperature on the diurnal cycle was not different for the two wind sectors considered ($p > 0.01$). Despite the two wind sectors had different mean amplitudes and were differently influenced by anthropogenic emissions, the effect of temperature on driving the amplitude of the diurnal cycle was the same.

Seasonal variability of Estany Llong and CRAM CO₂ concentration

The mean seasonal variation over 2010–2013 was calculated for ELL and CRAM and also for the other sites (CMN, MHD,

Fig. 3 **a** Wind rose for flask sampling events at Estany Llong (ELL). **b** Polar plot of CO₂ residuals from the four-harmonics fit curve at ELL. WSpeed wind speed (m/s)

CIB, and MBL) (Fig. 6). Annual maximum at ELL was found in March, and annual minimum in August. ELL annual cycle amplitude was maximum in 2010 (16.2 ppm CO₂) and minimum in 2011 (11.2 ppm CO₂), with an average of 13.2 ± 2.2 (1 σ) ppm CO₂.

For CRAM series, only the hours 10–12 GMT were selected for comparison with the ELL CO₂ series. Annual maximum was found in February, a month earlier than ELL, but the annual minimum was also observed in August. Annual cycle amplitude was maximum in 2011 (22.2 ppm CO₂) and minimum in 2013 (19.6 ppm CO₂), with an average of 20.6 ± 1.7 (1 σ) ppm.

The Mann-Whitney-Wilcoxon test was used to assess if the mean concentrations at ELL and CRAM were statistically different or not. The test was done for winter (DJF) and summer months (JJA). While in summer, the mean concentrations at both

locations were statistically different ($p < 10^{-4}$); winter concentrations were not distinct ($p > 0.1$). ELL monthly means were inside the first interquartile range of CRAM data from October to May. In summer, both series diverge, being CRAM measurements lower than ELL (5.5 ppm, 3.5 to 7.4 CI).

The highest correlation of the seasonal CO₂ cycle at ELL was observed with the other mountain site, Monte Cimone ($\rho = 0.84$), indicating that both mountain sites behave similarly under seasonal changes (Fig. 6). During spring, CRAM and ELL concentrations were similar to the CIBA, located at the same latitude ($p > 0.05$). However, CIBA measurements at summertime were extremely variable ($\sigma = 4.6$ ppm), causing higher average values (8.9 ppm, 7.6 to 10.3 CI, $p < 10^{-15}$). Differences in concentrations for ELL and MHD were not significant for any season ($p > 0.05$).

Fig. 4 Influence of wind direction on the observed CO₂ data measured at Centre de Recerca d'Alta Muntanya (CRAM). Bivariate polar plots of CO₂ data: **a** average 2010–2014; **b** diurnal (08 to 20 GMT); **c** nocturnal

(22 to 05 GMT); **d** summer (May to October); **e** winter (November to April). WSpeed wind speed (m/s)

Fig. 5 Influence of temperature on the observed CO₂ data measured at Centre de Recerca d'Alta Muntanya (CRAM). Natural logarithm of CO₂ diurnal amplitude against daily mean temperature for N-NW coming winds and for E-SE-S-SW winds and their corresponding Lloyd and Taylor fit (Lloyd and Taylor 1994)

CO₂ differences of ELL and CRAM with MBL were not statistically significant in winter and fall ($p > 0.1$). In summer, ELL also had similar concentrations as the MBL reference, but CRAM was 7.0 ppm higher (5.6 to 8.5 CI, $p < 10^{-15}$). The major differences between ELL and the MBL reference were encountered in spring, being ELL data 2.1 ppm lower on average than MBL (0.7 to 3.6 CI, $p < 0.05$). This might indicate that ELL data was more influenced by a regional or continental processes (with strong spring uptake) rather than very local, because the site is usually covered with snow in spring months (Albentosa Sánchez 1973).

Discussion

Annual cycle and growth rate comparison

The comparison of ELL data with other sites showed that ELL mountain station was mostly measuring background

Table 1 Parameters of the Lloyd and Taylor curve depending on forthcoming wind direction

Forthcoming wind direction	A	E_0	R^2
N-NW	2469	282.6	0.65
E-SE-S-SW	3699	289.5	0.72

Fig. 6 Average annual CO₂ cycle measured from the observed data at Centre de Recerca d'Alta Muntanya (CRAM), Estany Llong (ELL), Mace Head (MHD), Monte Cimone (CMN), CIBA (CIB), and NOAA Marine Boundary Layer (MBL) reference for the period 2010–2013. Shaded regions are the interquartile ranges for CRAM (light red) and ELL (gray)

concentrations and captured the annual fluctuations driven by global changes. The statistical difference between ELL and the MBL reference was only significant in spring, probably influenced by a major continental uptake by the former. These results suggest that ELL is a suitable site to help constraining background concentration in European top-down inversion models.

The high-mountain sites appeared to have similar behaviors when comparing annual cycles. ELL and CMN annual cycles showed high levels of correlation despite the maximum concentration at ELL was observed in March, a month earlier than at CMN and the MBL reference. In ELL, a large decrease of the measured CO₂ concentration was observed in April, probably caused by the influence of the southern Mediterranean vegetation from central Spain that is becoming photosynthetically more active earlier in late winter to early spring (Verger et al. 2016).

The lower summer concentrations observed at CRAM compared to ELL could be explained by higher photosynthesis rates near CRAM station. The surrounding vegetation differs between the two sites, with more mixed and deciduous forests in CRAM compared with meadows, coniferous forests, and rocky granite areas in ELL (Fig. S1). Winter similarities are coherent with low average temperatures and the presence of snow at both sites, reducing all kind of CO₂ uptake or release from vegetation or soils (Mast et al. 1998; Merbold et al. 2013). That might also explain the lower differences between ELL and CRAM with the NOAA MBL during winter months.

The strength of the biospheric and oceanic CO₂ exchanges strongly varies in space and time. This variability is, in turn, closely linked back to climatic influences (Goulden 1996; Piao et al. 2013). The annual terrestrial flux indicates the terrestrial CO₂ annual natural budget, i.e., the difference between the natural terrestrial sources and the natural terrestrial sinks (Houghton 2003). The magnitude of the annual growth from the different monitoring sites is correlated with the annual terrestrial flux from MACC-V2 and JENA-MPI biospheric models for the period 2008–2011 (Fig. 2). The increase in annual growth in 2010 was coincidental with an increase in the terrestrial flux. Longer time series are however needed to establish whether this relation is maintained over time.

Influence of wind direction and atmospheric temperature on CRAM CO₂ concentrations

The amplitude of the CO₂ diurnal cycle at CRAM was highly correlated with temperature following an exponential relation, probably driven by the exponential increase of soil respiration with temperature and the general increase of photosynthesis with temperature at low and mid temperatures (Berry and Bjorkman 1980; Smith and Dukes 2013). Seasonal changes in boundary layer height caused by temperature annual cycle may also affect the variability of the measured CO₂ (Pino et al. 2013). The amplitude of the diurnal cycle was also found to be dependent on wind direction. At CRAM annual average temperature (8.1 °C), the CO₂ diurnal amplitude for days with a predominant wind from N-NW sector was 4.3 ppm lower on average than the diurnal amplitude for days with a predominance of E-SE-S-SW winds. However, the logarithm of the CO₂ diurnal amplitude for both wind sectors co-varied symmetrically with the average daily temperature. A higher CO₂ diurnal signal produced by anthropogenic emissions from vehicles on the N-NW direction, coupled with a similar natural diurnal cycle for both wind sectors, could explain this behavior. Future measurements of CO, which has been widely used as a proxy tracer for fossil fuel-derived CO₂ (Meijer et al. 1996; Vogel et al. 2010; Turnbull et al. 2015), could help determine if the increase in CO₂ concentration from northerly winds was only caused by traffic emissions or was also due to lower vegetation influence.

Conclusions

The analysis of the ELL CO₂ series showed that this site behaves as a remote background station that is scarcely influenced by local micrometeorology and local potential sources. The fortnightly sample series are useful to understand the seasonal cycle and the annual growth rate, but poorly serves to know short-term influences. Annual growth rates at ELL were comparable to those observed in remote background

locations and correlated with CO₂ terrestrial flux models for 2008–2011.

Continuous data from CRAM showed that CO₂ concentrations were dependent on the wind direction according to the location of local sources and sinks. The near presence of the Vielha tunnel north of the monitoring site influenced the measured concentrations and on average CO₂ concentrations from this direction were ~ 5 ppm higher on average during daytime. A large amplitude of the CO₂ diurnal cycle (up to 31.4 ppm) was observed in summer. The variability of the diurnal amplitude was proved to be related to the average diurnal temperature due to photosynthesis and soil respiration. The use of a Lloyd and Taylor curve revealed a similar influence of the temperature in the variability of the CO₂ diurnal amplitude for N-NW and E-SE-S-SW wind sectors. CO₂ traffic emissions could be responsible for an increase of CO₂ concentrations in daytime, lowering the observable amplitude of the CO₂ diurnal cycle when wind was from the N-NW independently of temperature. Future traffic-related tracer measurements such as CO could help resolve the influence of traffic emissions.

The combination of two temporal CO₂ series, one at a high-mountain background environment with low time resolution but high precision measurements, and the second in a site more influenced by local emissions, with lower precision but higher time resolution, has been demonstrated to be useful for the study of the sources and sinks controlling the CO₂ budget in mountain ecosystems. Although the precision and accuracy of the GMP343 sensor are low, with a periodic calibration, it can improve up to 1.5 ppm. These CO₂ sensors can be useful to study the diurnal variability in remote stations and easy to deploy due to its low maintenance and low energy consumption.

Despite concentrations at the two locations were statistically similar during winter months, summer concentrations showed to be different indicating the presence of more photosynthetic active vegetation near CRAM than at ELL. A future installation of a continuous measurement station near ELL, not directly influenced by anthropogenic emissions, could help resolve the factors driving the diurnal and seasonal CO₂ cycle at ELL. Additional CO₂ flux measurements and soil decomposition studies (tea-bags ILTER global decomposition project, 2016–2019) are already being carried out in grassland areas in the National Park. These measurements will be used to better understand the local influences and the response of ecosystems to temperature changes and to reduce the uncertainties in the regional carbon balance.

Acknowledgments The authors would like to acknowledge the people from the Aiguestortes i Estany de Sant Maurici National Park, especially Mercè Aniz, for their support on this research. We would also like to thank the University of Barcelona (UB) for their actions and help in installing a continuous measurement station at the Centre de Recerca d'Alta Muntanya (CRAM). We finally thank all the people who

participated in the samplings, particularly Helena Parga, Laura Agraz, Rober Sanchez, Cristina Cereza, Montserrat Recolons, Manel Nofuentes, Silvia Borrás, Lidia Cañas, Eusebi Vazquez, and Paola Occhipinti. We used the Marine Boundary Layer reference product kindly provided by E. Dlugokencky, K. Masarie, P. Lang, and P. Tans from NOAA. We used the ObsPack CO₂ flask data from Mace Head, CIBA, and Monte Cimone stations. We thank all measurement groups for submitting CO₂ concentration data to the ObsPack-GLOBALVIEW product.

Funding information This work has been mainly funded by own resources of the LAO-Climadat Lab at IC3 and by the Fluxpyr project (POCTEFA Program EFA 34/08 - Interreg IV-A). Data is available through virtual access provided and funded in the frame of eLTER H2020 - European Long-Term Ecosystem and Socio-Ecological Research Infrastructure (EC project GA: 654359; INFRAIA; 2015–2019). **Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

Albentosa Sánchez LM (1973) Los climas de Catalunya. Estudio de climatología dinámica. Phd Thesis. University of Barcelona

Andrews AE, Kofler JD, Trudeau ME, Williams JC, Neff DH, Masarie KA, Chao DY, Kitzis DR, Novelli PC, Zhao CL, Dlugokencky EJ, Lang PM, Crotwell MJ, Fischer ML, Parker MJ, Lee JT, Baumann DD, Desai AR, Stanier CO, De Wekker SFJ, Wolfe DE, Munger JW, Tans PP (2014) CO₂, CO, and CH₄ measurements from tall towers in the NOAA earth system research laboratory's global greenhouse gas reference network: instrumentation, uncertainty analysis, and recommendations for future high-accuracy greenhouse gas monitoring efforts. *Atmos Meas Tech* 7:647–687. <https://doi.org/10.5194/amt-7-647-2014>

Angert A, Biraud S, Bonfils C, Henning CC, Buermann W, Pinzon J, Tucker CJ, Fung I (2005) Drier summers cancel out the CO₂ uptake enhancement induced by warmer springs. *Proc Natl Acad Sci* 102:10823–10827. <https://doi.org/10.1073/pnas.0501647102>

Bacardit M, Camarero L (2009) Fluxes of Al, Fe, Ti, Mn, Pb, Cd, Zn, Ni, Cu, and As in monthly bulk deposition over the Pyrenees (SW Europe): the influence of meteorology on the atmospheric component of trace element cycles and its implications for high mountain lakes. *J Geophys Res Biogeosci* 114:1–17. <https://doi.org/10.1029/2008JG000732>

Bakwin PS, Davis KJ, Yi C, Wofsy SC, Munger JW, Haszpra L, Barcza Z (2004) Regional carbon dioxide fluxes from mixing ratio data. *Tellus B* 56:301–311. <https://doi.org/10.1111/j.1600-0889.2004.00111.x>

Bamberger I, Oney B, Brunner D, Henne S, Leuenberger M, Buchmann N, Eugster W (2017) Observations of atmospheric methane and carbon dioxide mixing ratios: tall-tower or mountain-top stations? *Bound-Layer Meteorol* 164:135–159. <https://doi.org/10.1007/s10546-017-0236-3>

Bergamaschi P, Krol M, Meirink JF, Dentener F, Segers a, van Aardenne J, Monni S, Vermeulen AT, Schmidt M, Ramonet M, Yver C, Meinhardt F, Nisbet EG, Fisher RE, O'Doherty S, Dlugokencky EJ (2010) Inverse modeling of European CH₄ emissions 2001–2006. *J Geophys Res* 115:D22309. <https://doi.org/10.1029/2010JD014180>

Berry J, Bjorkman O (1980) Photosynthetic response and adaptation to temperature in higher plants. *Annu Rev Plant Physiol* 31:491–543. <https://doi.org/10.1146/annurev.pp.31.060180.002423>

Bouwman AF (1989) The role of soils and land use in the greenhouse effect. *Netherlands. J Agric Sci* 37:13–19

Bowman AW, Azzalini A (1997) Applied smoothing techniques for data analysis: the kernel approach with S-plus illustrations. Oxford University Press, Oxford

Camarero L, Catalan J (1993) Chemistry of bulk precipitation in the central and eastern Pyrenees, northeast Spain. *Atmos Environ Part A Gen Top* 27:83–94. [https://doi.org/10.1016/0960-1686\(93\)90073-8](https://doi.org/10.1016/0960-1686(93)90073-8)

Camarero L, Morgui J-A (2009) Recerca ecològica a llarg termini en el Parc Nacional: el node LTER-Aiguestortes. In: *Actas de las VIII Jornadas sobre Investigación del Parque Nacional de Aiguestortes i Estany de Sant Maurici*. Espot, (Lleida). Departament de Medi Ambient i Habitatge, Generalitat de Catalunya. Barcelona. España

Camarero L, Bacardit M, de Diego A, Arana G (2017) Decadal trends in atmospheric deposition in a high elevation station: effects of climate and pollution on the long-range flux of metals and trace elements over SW Europe. *Atmos Environ* 167:542–552. <https://doi.org/10.1016/j.atmosenv.2017.08.049>

Camarero L, Catalan J, Morgui JA, Gacia E (2018) Daily meteorology (wind, air temperature, humidity, precipitation, radiation) from 2005–2017 for Aiguestortes LTER site. <https://b2share.eudat.eu/records/3e0968efd0994389b2ce59f0bca3adc5>. Accessed 20 Jun 2018

Carreras J, Diego F (2007) Cartografia dels hàbitats a Catalunya 1: 50.000. In: *General. Catalunya*

Carrillo E, Ninot JM (1992) Flora i vegetació de les valls d'Espot i Boi. *Arx. la secció Ciències l'Institut d'Estudis Catalans* 99:474 + 350

Carlsaw DC, Ropkins K (2012) Openair — an R package for air quality data analysis. *Environ Model Softw* 27–28:52–61. <https://doi.org/10.1016/j.envsoft.2011.09.008>

Catalan J (1989) The winter cover of a high-mountain Mediterranean lake (Estany Redó, Pyrenees). *Water Resour Res* 25:519–527. <https://doi.org/10.1029/WR025i003p00519>

Chappuis E, Gacia E, Ballesteros E (2011) Changes in aquatic macrophyte flora over the last century in Catalan water bodies (NE Spain). *Aquat Bot* 95:268–277. <https://doi.org/10.1016/j.aquabot.2011.08.006>

Chappuis E, Gacia E, Ballesteros E (2014) Environmental factors explaining the distribution and diversity of vascular aquatic macrophytes in a highly heterogeneous Mediterranean region. *Aquat Bot* 113:72–82. <https://doi.org/10.1016/j.aquabot.2013.11.007>

Chevallier F, Ciais P, Conway TJ, Aalto T, Anderson BE, Bousquet P, Brunke EG, Ciattaglia L, Esaki Y, Fröhlich M, Gomez A, Gomez-Pelaez AJ, Haszpra L, Krummel PB, Langenfelds RL, Leuenberger M, MacHida T, Maignan F, Matsueda H, Morgui JA, Mukai H, Nakazawa T, Peylin P, Ramonet M, Rivier L, Sawa Y, Schmidt M, Steele LP, Vay SA, Vermeulen AT, Wofsy S, Worthy D (2010) CO₂ surface fluxes at grid point scale estimated from a global 21 year reanalysis of atmospheric measurements. *J Geophys Res Atmos* 115:1–17. <https://doi.org/10.1029/2010JD013887>

Ciais P, Peylin P, Bousquet P (2000) Regional biospheric carbon fluxes as inferred from atmospheric CO₂ measurements. *Ecol Appl* 10:1574. <https://doi.org/10.2307/2641225>

Ciais P, Rayner P, Chevallier F, Bousquet P, Logan M, Peylin P, Ramonet M (2010) Atmospheric inversions for estimating CO₂ fluxes: methods and perspectives. *Clim Chang* 103:69–92. <https://doi.org/10.1007/s10584-010-9909-3>

Ciattaglia L (1983) Interpretation of atmospheric CO₂ measurements at Mt. Cimone (Italy) related to wind data. *J Geophys Res* 88:1331. <https://doi.org/10.1029/JC088iC02p01331>

Ciattaglia L, Cundari V, Colombo T (1987) Further measurements of atmospheric carbon dioxide at Mt. Cimone, Italy: 1979–1985. *Tellus B* 39 B:13–20. <https://doi.org/10.1111/j.1600-0889.1987.tb00266.x>

Colombo T, Santaguida R, Capasso A, Calzolari F, Evangelisti F, Bonasoni P (2000) Biospheric influence on carbon dioxide

- measurements in Italy. *Atmos Environ* 34:4963–4969. [https://doi.org/10.1016/S1352-2310\(00\)00366-6](https://doi.org/10.1016/S1352-2310(00)00366-6)
- Conway TJ, Tans PP, Waterman LS, Thoning KW, Kitzis DR, Masarie KA, Zhang N (1994) Evidence for interannual variability of the carbon cycle from the National Oceanic and Atmospheric Administration Climate Monitoring and Diagnostics Laboratory Global Air Sampling Network. *J Geophys Res - Atmos* 99:22831–22855. <https://doi.org/10.1029/94JD01951>
- Cooperative Global Atmospheric Data Integration Project. 2013 updated annually. M compilation of synchronized and gap-filled atmospheric carbon dioxide records for the period 1979–2012 (2013) obspack_co2_1_GLOBALVIEW-CO2_2013_v1.0.4 2013-12-23. Compiled by NOAA Global Monitoring Division: Boulder, Colorado, U.S.A. <https://doi.org/10.3334/OBSPACK/1002>. Accessed 1 Oct 2018
- Cooperative Global Atmospheric Data Integration Project; (2017): Multi-laboratory compilation of atmospheric carbon dioxide data for the period 1957–2016 (2017) obspack_co2_1_GLOBALVIEWplus_v3.2_2017_11_02; NOAA Earth System Research Laboratory, Global Monitoring Division. <https://doi.org/10.15138/G3704H>. Accessed 1 Oct 2018
- Cristofanelli P, Brattich E, Decesari S, Landi TC, Maione M, Putero D, Tositti L, Bonasoni P (2018) The “O. Vittori” observatory at Mt. Cimone: a “Lighthouse” for the Mediterranean troposphere. *High-Mountain Atmos Res SpringerBriefs Meteorol Springer, Cham*. https://doi.org/10.1007/978-3-319-61127-3_1
- Crosson ER (2008) A cavity ring-down analyzer for measuring atmospheric levels of methane, carbon dioxide, and water vapor. *Appl Phys B Lasers Opt* 92:403–408. <https://doi.org/10.1007/s00340-008-3135-y>
- Cundari V, Colombo T, Papini G, Benedicti G, Ciattaglia L (1990) Recent improvements on atmospheric CO₂ measurements at Mt. Cimone observatory, Italy. *Nuovo Cim C* 13:871–882. <https://doi.org/10.1007/BF02512003>
- Cundari V, Colombo T, Ciattaglia L (1995) Thirteen years of atmospheric carbon dioxide measurements at Mt. Cimone station, Italy. *Nuovo Cim C* 18:33–47. <https://doi.org/10.1007/BF02561457>
- Curcoll R, Recolons M, Font A, Agraz L, Parga E, Bacardit E, Camarero L, Pueyo S, Rodó X, Morgui JA (2010) First 2 years of atmospheric CO₂ measurements in the Estany Llong plain (2100 masl, Parc Nacional d'Aigüestortes i Estany de Sant Maurici, Pyrenees, Catalonia, Spain). In: *EGU General Assembly Conference Abstracts*. p 3608
- Dlugokencky EJ, Lang PM, Mund JW, Crotwell AM, Crotwell MJ, Thoning KW (2017a) Atmospheric carbon dioxide dry air mole fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network. 1968–2016, Version: 2017-07-28. ftp://aftp.cmdl.noaa.gov/data/trace_gases/co2/flask/surface/. Accessed 1 Oct 2018
- Dlugokencky EJ, Thoning KW, Lang PM, Tans PP (2017b) NOAA greenhouse gas reference from atmospheric carbon dioxide dry air mole fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network. ftp://aftp.cmdl.noaa.gov/data/trace_gases/co2/flask/surface/. Accessed 1 Oct 2018
- Geels C, Gloor M, Ciais P, Bousquet P, Peylin P, Vermeulen A, T, Dargaville R, Aalto T, Brandt J, Christensen JH, Frohn LM, Haszpra L, Karstens U, Rödenbeck C, Ramonet M, Carboni G, Santaguida R (2007) Comparing atmospheric transport models for future regional inversions over Europe - part 1: mapping the atmospheric CO₂ signals. *Atmos Chem Phys* 7:3461–3479. <https://doi.org/10.5194/acp-7-3461-2007>
- Gomez-Pelaez AJ, Ramos R (2011) Improvements in the carbon dioxide and methane continuous measurement programs at Izaña Global GAW Station (Spain) during 2007–2009. In: *Rep. 15th WMO/IAEA Meet. Expert. Carbon Dioxide, Other Greenh. Gases, Relat. Tracer Meas. Tech.* 7–10 Sept. 2009, GAW Rep. number 194, WMO TD 1553. https://library.wmo.int/pmb_ged/wmo-td_1553.pdf. Accessed 1 Oct 2018
- Goulden ML (1996) Exchange of carbon dioxide by a deciduous forest: response to interannual climate variability. *Science* (80-) 271:1576–1578. <https://doi.org/10.1126/science.271.5255.1576>
- Hase F, Frey M, Blumenstock T, Groß J, Kiel M, Kohlhepp R, Mengistu Tsidu G, Schäfer K, Sha MK, Orphal J (2015) Application of portable FTIR spectrometers for detecting greenhouse gas emissions of the major city Berlin. *Atmos Meas Tech* 8:3059–3068. <https://doi.org/10.5194/amt-8-3059-2015>
- Hervàs A, Camarero L, Reche I, Casamayor EO (2009) Viability and potential for immigration of airborne bacteria from Africa that reach high mountain lakes in Europe. *Environ Microbiol* 11:1612–1623. <https://doi.org/10.1111/j.1462-2920.2009.01926.x>
- Houghton RA (2003) Revised estimates of the annual net flux of carbon to the atmosphere from changes in land use and land management 1850–2000. *Tellus Ser B Chem Phys Meteorol* 55:378–390. <https://doi.org/10.1034/j.1600-0889.2003.01450.x>
- Hu C, Griffith TJ, Lee X, Millet DB, Chen Z, Baker JM, Xiao K (2018) Top-down constraints on anthropogenic CO₂ emissions within an agricultural-urban landscape. *J Geophys Res Atmos* 123:4674–4694. <https://doi.org/10.1029/2017JD027881>
- Le Quéré C, Andrew RM, Canadell JG, Sitch S, Korsbakken JJ, Peters GP, Manning AC, Boden TA, Tans PP, Houghton RA, Keeling RF, Alin S, Andrews OD, Anthoni P, Barbero L, Bopp L, Chevallier F, Chini LP, Ciais P, Currie K, Delire C, Doney SC, Friedlingstein P, Gkritzalis T, Harris I, Hauck J, Haverd V, Hoppema M, Klein Goldewijk K, Jain AK, Kato E, Körtzinger A, Landschützer P, Lefèvre N, Lenton A, Lienert S, Lombardozi D, Melton JR, Metzl N, Millero F, Monteiro PMSS, Munro DR, Nabel JEMSMS, Nakaoka SI, O'Brien K, Olsen A, Omar AM, Ono T, Pierrot D, Poulter B, Rödenbeck C, Salisbury J, Schuster U, Schwinger J, Séférian R, Skjelvan I, Stocker BD, Sutton AJ, Takahashi T, Tian H, Tilbrook B, van der Laan-Luijkx IT, van der Werf GR, Viovy N, Walker AP, Wiltshire AJ, Zaehle S, Ivar Korsbakken J, Peters GP, Manning AC, Boden TA, Tans PP, Houghton RA, Keeling RF, Alin S, Andrews OD, Anthoni P, Barbero L, Bopp L, Chevallier F, Chini LP, Ciais P, Currie K, Delire C, Doney SC, Friedlingstein P, Gkritzalis T, Harris I, Hauck J, Haverd V, Hoppema M, Klein Goldewijk K, Jain AK, Kato E, Körtzinger A, Landschützer P, Lefèvre N, Lenton A, Lienert S, Lombardozi D, Melton JR, Metzl N, Millero F, Monteiro PMSS, Munro DR, Nabel JEMSMS, Nakaoka SI, O'Brien K, Olsen A, Omar AM, Ono T, Pierrot D, Poulter B, Rödenbeck C, Salisbury J, Schuster U, Schwinger J, Séférian R, Skjelvan I, Stocker BD, Sutton AJ, Takahashi T, Tian H, Tilbrook B, van der Laan-Luijkx IT, van der Werf GR, Viovy N, Walker AP, Wiltshire AJ, Zaehle S (2016) Global carbon budget 2016. *Earth Syst Sci Data* 8:605–649. <https://doi.org/10.5194/essd-8-605-2016>
- Levin I (1987) Atmospheric CO₂ in continental European alternative approach to clean air CO₂ data. *Tellus* 39B:21–28. <https://doi.org/10.1111/j.1600-0889.1987.tb00267.x>
- Lloyd J, Taylor J (1994) On the temperature dependence of soil respiration. *Funct Ecol* 8:315–323. <https://doi.org/10.2307/2389824>
- Marquis M, Tans P (2008) Climate change: carbon crucible. *Science* (80-) 320:460–461. <https://doi.org/10.1126/science.1156451>
- Masarie KA, Tans PP (1995) Extension and integration of atmospheric carbon dioxide data into a globally consistent measurement record. *J Geophys Res* 100:11593. <https://doi.org/10.1029/95JD00859>
- Mast MA, Wickl KP, Striegl RT, Clow WD (1998) Winter fluxes of CO₂ and CH₄ from subalpine soils in Rocky Mountain National Park, Colorado. *Glob Biogeochem Cycles* 12:607–620. <https://doi.org/10.1029/98GB02313>
- Meijer HAJ, Smid HM, Perez E, Keizer MG (1996) Isotopic characterisation of anthropogenic CO₂ emissions using isotopic and

- 900 radiocarbon analysis. *Phys Chem Earth* 21:483–487. [https://doi.org/](https://doi.org/10.1016/S0079-1946(97)81146-9)
901 [10.1016/S0079-1946\(97\)81146-9](https://doi.org/10.1016/S0079-1946(97)81146-9)
- 902 Merbold L, Steinlin C, Hagedorn F (2013) Winter greenhouse gas fluxes
903 (CO₂, CH₄ and N₂O) from a subalpine grassland. *Biogeosciences*
904 10:3185–3203. <https://doi.org/10.5194/bg-10-3185-2013>
- 905 Ninyerola M, Pons X, Roure JM, Martin VJ, Raso J, Clavero P (2003)
906 *Atles climàtic digital de Catalunya*. In: *Serv. Meteorològic*
907 *Catalunya i Dep. Medi Ambient General*. Catalunya Barcelona.
908 http://www.opengis.uab.cat/acdc/en_index.htm. Accessed 1
909 Oct 2018
- 910 Peylin P, Law RM, Gurney KR, Chevallier F, Jacobson AR, Maki T,
911 Niwa Y, Patra PK, Peters W, Rayner PJ, Rödenbeck C, Van Der
912 Laan-Luijkx IT, Zhang X (2013) Global atmospheric carbon budget:
913 results from an ensemble of atmospheric CO₂ inversions.
914 *Biogeosciences* 10:6699–6720. [https://doi.org/10.5194/bg-10-](https://doi.org/10.5194/bg-10-6699-2013)
915 [6699-2013](https://doi.org/10.5194/bg-10-6699-2013)
- 916 Piao S, Sitch S, Ciais P, Friedlingstein P, Peylin P, Wang X, Ahlström A,
917 Anav A, Canadell JG, Cong N, Huntingford C, Jung M, Levis S,
918 Levy PE, Li J, Lin X, Lomas MR, Lu M, Luo Y, Ma Y, Myneni RB,
919 Poulter B, Sun Z, Wang T, Viovy N, Zaehle S, Zeng N (2013)
920 Evaluation of terrestrial carbon cycle models for their response to
921 climate variability and to CO₂ trends. *Glob Chang Biol* 19:2117–
922 2132. <https://doi.org/10.1111/gcb.12187>
- 923 Pickers PA, Manning AC (2015) Investigating bias in the application of
924 curve fitting programs to atmospheric time series. *Atmos Meas Tech*
925 8:1469–1489. <https://doi.org/10.5194/amt-8-1469-2015>
- 926 Pillai D, Gerbig C, Ahmadov R, Rödenbeck C, Kretschmer R, Koch T,
927 Thompson R, Neininger B, Lavrié JV (2011) High-resolution sim-
928 ulations of atmospheric CO₂ over complex terrain-representing the
929 Ochsenskopf mountain tall tower. *Atmos Chem Phys* 11:7445–7464.
930 <https://doi.org/10.5194/acp-11-7445-2011>
- 931 Pino D, Kaikkonen J-P, de Arellano JV-G (2013) Quantifying the uncer-
932 tainties of advection and boundary layer dynamics on the diurnal
933 carbon dioxide budget. *J Geophys Res Atmos* 118:9376–9392.
934 <https://doi.org/10.1002/jgrd.50677>
- 935 Post WM, Peng T-HH, Emanuel WR, King AW, Dale VH, DeAngelis DL
936 (1990) The global carbon cycle. *Am Sci* 78:892–895
- 937 Potter CSC, Randerson J (1993) Terrestrial ecosystem production: a pro-
938 cess model based on global satellite and surface data. *Glob*
939 *Biogeochem Cycles* 7:811–841. [https://doi.org/10.1029/](https://doi.org/10.1029/93GB02725)
940 [93GB02725](https://doi.org/10.1029/93GB02725)
- 941 Pumpanen J, Ilvesniemi H, Kulmala L, Siivola E, Laakso H, Kolari P,
942 Helenelund C, Laakso M, Uusimaa M, Hari P (2008) Respiration in
943 boreal forest soil as determined from carbon dioxide concentration
944 profile. *Soil Sci Soc Am J* 72:1187. [https://doi.org/10.2136/](https://doi.org/10.2136/sssaj2007.0199)
945 [sssaj2007.0199](https://doi.org/10.2136/sssaj2007.0199)
- 946 Randerson JT, Field CB, Fung IY, Tans PP (1999) Increases in early
947 season ecosystem uptake explain recent changes in the seasonal
948 cycle of atmospheric CO₂ at high northern latitudes. *Geophys Res*
949 *Lett* 26:2765–2768. <https://doi.org/10.1029/1999GL900500>
- 950 Reichstein M, Falge E, Baldocchi D, Papale D, Aubinet M, Berbigier P,
951 Bernhofer C, Buchmann N, Gilmanov T, Granier A, Grunwald T,
952 Havrankova K, Ilvesniemi H, Janous D, Knohl A, Laurila T, Lohila
953 A, Loustau D, Matteucci G, Meyers T, Miglietta F, Ourcival J-M,
954 Pumpanen J, Rambal S, Rotenberg E, Sanz M, Tenhunen J, Seufert
955 G, Vaccari F, Vesala T, Yakir D, Valentini R (2005) On the separa-
956 tion of net ecosystem exchange into assimilation and ecosystem
957 respiration: review and improved algorithm. *Glob Chang Biol* 11:
958 1424–1439. <https://doi.org/10.1111/j.1365-2486.2005.001002.x>
- 959 Rella CW, Chen H, Andrews AE, Filges A, Gerbig C, Hatakka J, Karion
960 A, Miles NL, Richardson SJ, Steinbacher M, Sweeney C, Wastine
961 B, Zellweger C (2013) High accuracy measurements of dry mole
962 fractions of carbon dioxide and methane in humid air. *Atmos Meas*
963 *Tech* 6:837–860. <https://doi.org/10.5194/amt-6-837-2013>
- 964 Richardson SJ, Miles NL, Davis KJ, Crosson ER, Rella CW, Andrews
965 AE (2012) Field testing of cavity ring-down spectroscopy analyzers
966 measuring carbon dioxide and water vapor. *J Atmos Ocean Technol* 966
967 29:397–406. <https://doi.org/10.1175/JTECH-D-11-00063.1>
- 968 Rigby M, Toumi R, Fisher R, Lowry D, Nisbet EG (2008) First contin-
969 uous measurements of CO₂ mixing ratio in central London using a
970 compact diffusion probe. *Atmos Environ* 42:8943–8953. [https://doi.](https://doi.org/10.1016/j.atmosenv.2008.06.040)
971 [org/10.1016/j.atmosenv.2008.06.040](https://doi.org/10.1016/j.atmosenv.2008.06.040)
- 972 Rödenbeck C, Houweling S, Gloor M, Heimann M (2003) CO₂ flux
973 history 1982–2001 inferred from atmospheric data using a global
974 inversion of atmospheric transport. *Atmos Chem Phys* 3:1919–
975 1964. <https://doi.org/10.5194/acp-3-1919-2003>
- 976 Sánchez ML, Pérez IA, García MA (2010) Study of CO₂ variability at
977 different temporal scales recorded in a rural Spanish site. *Agric For*
978 *Meteorol* 150:1168–1173. [https://doi.org/10.1016/j.agrformet.2010.](https://doi.org/10.1016/j.agrformet.2010.04.018)
979 [04.018](https://doi.org/10.1016/j.agrformet.2010.04.018)
- 980 Schibig MF, Steinbacher M, Buchmann B, Van Der Laan-Luijkx IT, Van
981 Der Laan S, Ranjan S, Leuenberger MC (2015) Comparison of
982 continuous in situ CO₂ observations at Jungfraujoch using two dif-
983 ferent measurement techniques. *Atmos Meas Tech* 8:57–68. [https://](https://doi.org/10.5194/amt-8-57-2015)
984 doi.org/10.5194/amt-8-57-2015
- 985 Schmidt M (2003) The Schauinsland CO₂ record: 30 years of continental
986 observations and their implications for the variability of the
987 European CO₂ budget. *J Geophys Res* 108:4619. [https://doi.org/](https://doi.org/10.1029/2002JD003085)
988 [10.1029/2002JD003085](https://doi.org/10.1029/2002JD003085)
- 989 Smith NG, Dukes JS (2013) Plant respiration and photosynthesis in
990 global-scale models: incorporating acclimation to temperature and
991 CO₂. *Glob Chang Biol* 19:45–63. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2012.02797.x)
992 [2486.2012.02797.x](https://doi.org/10.1111/j.1365-2486.2012.02797.x)
- 993 Sturm P, Leuenberger M, Sirignano C, Neubert REM, Meijer HAJ,
994 Langenfelds R, Brand WA, Tohjima Y (2004) Permeation of atmo-
995 spheric gases through polymer O-rings used in flasks for air sam-
996 pling. *J Geophys Res Atmos* 109:n/a-n/a. [https://doi.org/10.1029/](https://doi.org/10.1029/2003JD004073)
997 [2003JD004073](https://doi.org/10.1029/2003JD004073)
- 998 Tans PP, Fung IY, Takahashi T (1990a) Observational constraints on the
999 global atmospheric CO₂ budget. *Science* (80-) 247:1431–1438.
1000 <https://doi.org/10.1126/science.247.4949.1431>
- 1001 Tans PP, Thoning KW, Elliott WP, Conway TJ (1990b) Error estimates of
1002 background atmospheric CO₂ patterns from weekly flask samples. *J*
1003 *Geophys Res* 95:14063. <https://doi.org/10.1029/JD095iD09p14063>
- 1004 Thoning KW, Tans PP, Komhyr WD (1989) Atmospheric carbon dioxide
1005 at Mauna Loa observatory: 2. Analysis of the NOAA GMCC data,
1006 1974–1985. *J Geophys Res Atmos* 94:8549–8565. [https://doi.org/](https://doi.org/10.1029/JD094iD06p08549)
1007 [10.1029/JD094iD06p08549](https://doi.org/10.1029/JD094iD06p08549)
- 1008 Turnbull JC, Sweeney C, Karion A, Newberger T, Lehman SJ, Tans PP,
1009 Davis KJ, Lauvaux T, Miles NL, Richardson SJ, Cambaliza MO,
1010 Shepson PB, Gurney K, Patarasuk R, Razlivanov I (2015) Toward
1011 quantification and source sector identification of fossil fuel CO₂
1012 emissions from an urban area: results from the INFLUX experiment.
1013 *J Geophys Res Atmos* 120:292–312. [https://doi.org/10.1002/](https://doi.org/10.1002/2014JD022555)
1014 [2014JD022555](https://doi.org/10.1002/2014JD022555)
- 1015 Vaisala (2017) Vaisala CARBOCAP Carbon Dioxide Probe GMP343. In:
1016 Vaisala Oyj, Finl. [https://www.vaisala.com/sites/default/files/](https://www.vaisala.com/sites/default/files/documents/GMP343-Datasheet-B210688EN-F.pdf)
1017 [documents/GMP343-Datasheet-B210688EN-F.pdf](https://www.vaisala.com/sites/default/files/documents/GMP343-Datasheet-B210688EN-F.pdf). Accessed 1
1018 Oct 2018
- 1019 Verger A, Filella I, Baret F, Peñuelas J (2016) Vegetation baseline phe-
1020 nology from kilometeric global LAI satellite products. *Remote Sens*
1021 *Environ* 178:1–14. <https://doi.org/10.1016/j.rse.2016.02.057>
- 1022 Vogel FR, Hammer S, Steinhof A, Kromer B, Levin I (2010) Implication
1023 of weekly and diurnal ¹⁴C calibration on hourly estimates of CO-
1024 based fossil fuel CO₂ at a moderately polluted site in southwestern
1025 Germany. *Tellus Ser B Chem Phys Meteorol* 62:512–520. <https://doi.org/10.1111/j.1600-0889.2010.00477.x>
- 1026 Young SG, Bowman AW (1995) Non-parametric analysis of covariance.
1027 *Biometrics* 51:920–931. <https://doi.org/10.2307/2532993>
- 1028 Zhao CL, Tans PP (2006) Estimating uncertainty of the WMO mole
1029 fraction scale for carbon dioxide in air. *J Geophys Res* 111:
1030 D08S09. <https://doi.org/10.1029/2005JD006003>
- 1031

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES.

- Q1. Please check if the corresponding author's affiliation and telecommunication data are correctly captured/indicated. Also, please check if the author group and affiliations are presented correctly.
- Q2. The citation “Dlugokencky et al. 2017” has been changed to “Dlugokencky et al., 2017a” to match the author name/date in the reference list. Please check if the change is fine in this occurrence and modify the subsequent occurrences, if necessary.
- Q3. Please check if figure captions and their corresponding images are captured and presented correctly.

UNCORRECTED PROOF