**Vent-proximal sub-seafloor replacement clastic-carbonate hosted SEDEX-type
mineralization in the Mehdiabad world-class Zn-Pb-Ba-(Cu-Ag) deposit, Southern Yazd
Basin, Iran**

Sajjad Maghfouri^{a*} (Corresponding Author)

^a*Department of Geology, Faculty of Basic Sciences, Tarbiat Modares University, Tehran, Iran*

E-Mail: s.maghfouri@modares.ac.ir

Mohammad Reza Hosseinzadeh^b

^b*Department of Earth Sciences, Faculty of Natural Sciences, University of Tabriz, Tabriz, Iran*

E-Mail: mr-hosseinzadeh@tabrizu.ac.ir

Flavien Choulet^c

^c*Chrono-Environnement, Université de Franche-Comté/CNRS, 25030 Besançon cedex, France*

E-Mail: flavien.choulet@univ-fcomte.fr

Pura Alfonso^d

^d*Departament d'Enginyeria Minera i Recursos Minerals, Universitat Politècnica de*

Catalunya, Catalunya, Spain

E-Mail: pura@emrn.upc.edu

Amir M. Azim Zadeh ^e

^e *Department of Geology, Faculty of Science, University of Zanjan, Zanjan, Iran*

E-Mail: amir.azimzadeh@znu.ac.ir

Abdorrahman Rajabi ^f

^f *School of Geology, College of Science, University of Tehran, Tehran, Iran*

E-Mail: abdorrahman.rajabi@gmail.com

Abstract

The Mehdiabad Zn-Pb-Ba (Cu-Ag) deposit in the southern Yazd Basin, central Iran, is the largest sediment-hosted Zn-Pb deposit in Iran. This deposit is hosted by organic carbon matter-rich shale, fine-grained black siltstone, and dolomite interlayered with sandstone of the Taft Formation. Sedimentological and geochemical studies of the Taft Formation have shown that these organic carbon matter-rich shales formed during a period of basin deepening and under anoxic conditions. Based on the orebody structure, mineralogy, and ore fabrics, we recognize five different ore facies types in the Mehdiabad deposit: (1) a stockwork/feeder zone, consisting of a discordant sulfide mineralization, forming a stockwork of sulfide-bearing dolomite veins, cutting the sedimentary rocks of the footwall; (2) massive-replacement ore, including pervasive replacement carbonate by pyrite, chalcopyrite, galena, and sphalerite with minor barite; (3) a bedded ore, with laminated to disseminated pyrite, sphalerite, and galena; (4) barite ore, which contains accessory sulfide minerals and rare calcite at the top of the deposit, and (5) a distal facies, with minor disseminated and laminated pyrite, banded cherts, and disseminated barite. Two stages of base metal sulfide replacement have separate origins. Fine-grained sulfide bands (stage I), intricately interlayered with organic carbon-rich beds and thin turbidite beds, exhibit lamina and bedding textures, supporting a syn-sedimentary (onto the sea floor) origin. Coarse-grained base metal sulfides (stage II), occurring within breccias and veins to veinlets are considered to have formed by replacement during post sedimentation sub-seafloor fluid flow. $\delta^{34}\text{S}$ values of pyrite, sphalerite chalcopyrite and galena range from -22 to +4.6.8‰. The highest $\delta^{34}\text{S}$ values correspond to the feeder zone (+4.6 and -10.5‰), whereas massive-replacement (+3.7 to - 13.7‰) and bedded (-22 to -17‰) ore facies display depleted compositions. The overall range of $\delta^{34}\text{S}$ values is remarkably higher than typical magmatic values, suggesting that

sulfides formed from the reduction of seawater sulfate by bacteriogenic sulfate reduction in a closed or semi-closed system in the bedded ore, whereas thermochemical sulfate reduction likely played an important role in the feeder zone. The formation of the Mehdiabad deposit follows the evolution of the sedimentary basin. Abrupt lateral changes in facies and thickness, along with the existence of synsedimentary breccias and debris flows within the ore sequence, suggest the proximity to synsedimentary faults and tectonic activity contemporaneous with the sedimentation in the Lower Cretaceous, favorable for the formation of the ore deposit. The deposit formed from a combination of hydrothermal and syn-sedimentary processes. Sulfur isotopes, together with sedimentological, textural, mineralogical, and geochemical evidences, suggest that this deposit should be classified as a vent-proximal sub-seafloor replacement SEDEX ore deposit.

Keywords: Lower Cretaceous; Mehdiabad Zn-Pb-Ba (Cu - Ag) deposit; SEDEX-type, Southern Yazd Basin; Iran

1- Introduction

Sediment-hosted Zn-Pb ore deposits are major sources of base metals such as Zn-Pb-Ag ± Cu-Ba and are the primary sources of Zn and Pb (Large et al., 2005). These deposits are diversely hosted by carbonate or siliciclastic rocks, without significant genetic contributions of magmatic activity (Leach et al., 2005). The deposits display a broad range of relationships to enclosing host rocks, including stratiform, strata-bound, and discordant ores (Leach et al., 2005). About 600 sediment-hosted Zn-Pb deposits and occurrences are presently known in Iran (Fig. 1). Only a few deposits of these have been explored and/or exploited although <70 deposits are currently being mined (Rajabi et al., 2012; 2013). Despite numerous scientific debates, these deposits are considered to form a metallogenic range from exhalative sedimentary (SEDEX) to

Mississippi Valley-types (MVTs) [deposits](#) (Rajabi et al., 2102; 2013). The most important [Iranian](#) sediment-hosted Zn-Pb ore deposits include the world-class Mehdiabad Zn-Pb-Ba (Cu-Ag) deposit (Maghfouri et al., 2015), the deposits world-class Irankuh Zn-Pb ($\pm$ Ag) district (Rastad, 1981; Ghazban et al., 1994; Boveiri and Rastad, 2016; Boveiri et al., 2017), the world-class Angouran Zn-Pb deposit (Gilg et al., 2006; Boni et al., 2007; Daliran et al., 2013), the Ahangaran Pb-Ag ($\pm$ Zn) deposit (Momenzadeh, 1976; Momenzadeh et al., 1979), the Koushk Zn-Pb vent-proximal SEDEX-type deposit (Yaghubpur and Mehrabi, 1997; Rajabi et al., 2012), the Emarat Zn-Pb MVT-type deposit (Ehya et al., 2010), the Tiran Zn-Pb ($\pm$ Ag $\pm$ Cu $\pm$ Ba) Sedex-type deposit (Yarmohammadi et al., 2016), the Nakhlak Pb-Zn MVT-type deposit (Rasa, 1987), and the Kuh-e-Sormeh Zn-Pb MVT-type deposit (Liaghat et al., 2000). These Zn-Pb deposits belong to the Yazd-Anarak metallogenic belt (YAMB), the Malayer-Esfahan metallogenic belt (MEMB), and the Central Alborz metallogenic belt (CAMB) (Rajabi et al., 2012; 2103) (Fig. 1).

The Yazd-Anarak Metallogenic Belt (YAMB) is located in the western section of the Central East Iranian Microcontinent (CEIM) (Fig. 1). The YAMB hosts abundant mineral deposits, particularly those of the Cu (Fe) porphyry-skarn (Darreh Zarashk and Hasanabad deposits (Oligocene–Miocene)), W-Cu-Ni-Au-As vein-types (Chah Palang (post-Jurassic), Talmesi and Meskani deposits (post-Eocene)), Zn-Pb MVT-types (Nakhlak, Darreh-Zanjir, Chahake Ardakani and Howze Safid deposits (post-late Cretaceous)), and Zn-Pb SEDEX types (Mehdiabad, Farahabad, Mansourabad (lower Cretaceous) and Haft-Har (lower Cambrian) deposits), being one of the main metallogenic provinces in Iran (Fig. 2) (Maghfouri et al., 2015; 2016a; 2016b; Maghfouri, 2017).

95 The southern Yazd Basin in the YAMB, which includes three lower Cretaceous [sediment-hosted](#)
 96 Zn-Pb $\pm$ Ba deposits, including [the](#) Mehdiabad deposit and the Farahabad and Mansourabad
 97 deposits (Figs. 2 and 3). Sedimentation during this [geological time span](#) was controlled by basin
 98 subsidence and by the reactivation [of half-graben](#) faults. In the southern Yazd Basin,
 99 sedimentation [has occurred](#) asymmetrically throughout the basin. The lower part of the lower
 100 Cretaceous sequence is characterized by major lateral facies changes, observable between the
 101 Mehdiabad and the Farahabad areas, and by the [wedge-shaped](#) geometry of the sedimentary
 102 units; these stratigraphic features are indicative of rift related half-grabens (e.g., Betts and Lister,
 103 2001; Betts et al., 1998). The Mehdiabad deposit, located 115 km southeast of Yazd city, can be
 104 subdivided in four ore zones (Fig. 4): 1) the Black-Hill ore (BH), the Central Valley Orebody
 105 (CVOB), the East Ridge ore (ER) and the Calamine Mine (CM). Throughout this paper, the term
 106 “Mehdiabad deposits” refers collectively to the BH, CVOB, ER and CM ore bodies. The
 107 Mehdiabad deposit is the largest accumulation of carbonate-hosted zinc in the world (Leach et
 108 al., 2005), with a total resource of 394 Mt, grading 4.2% Zn, 1.6% Pb and 36 g/t Ag. In addition,
 109 the barite content of this deposit is unusually large (>40 Mt) and covers an area of at least 1.5
 110 km². Early studies at Mehdiabad (Ghasemi, 2006) [described](#) the initial geologic framework of
 111 the deposit. This work showed that [the](#) Mehdiabad deposit bears geologic similarities with other
 112 sediment-hosted deposits of the southern Yazd Basin. Ghasemi (2006) also [proposed a](#)
 113 “Mehdiabad-type”, [of deposits](#), to [distinguish](#) the YAMB deposits from other classical sediment-
 114 hosted Zn-Pb deposits (SEDEX, MVT, Irish and [massive sulfides](#)). Mohseni (2011) further
 115 studied the barite mineralization, located in the western part of the deposit, and classified it as an
 116 Irish-type deposit. [Recently petrological](#) and geochemical studies of the deposit were carried out
 117 in the frame of a Ph.D. research project between 2012 and 2017 and Maghfouri et al. (2015) and

Maghfouri (2017) presented geologic descriptions of the deposit and a preliminary genetic model. Based on the occurrence of exhalative hydrothermal distinctive characteristics such as laminated sulfide textures, lateral metal zonation, diagenetic replacement textures and the presence of a normal syn-sedimentary fault (Black-Hill Fault), a SEDEX model was proposed for the Mehdiabad deposit. Maghfouri and Hosseinzadeh (2018) studied the Mansourabad deposit, which is located in the southern Yazd Basin (Fig.3), and classified it as a vent-proximal sub-seafloor replacement SEDEX deposit model. They further proposed that the formation of SEDEX deposits in this basin is related to the evolution of an extensional Cretaceous environment that affected the Central Iranian microcontinent. This paper aims at providing advanced evidence from regional sedimentology, geochemistry, and sulfide-sediment relationships to support the synsedimentary origin for the Mehdiabad deposit.

2- Geological context and main features of the Mehdiabad deposit

2-1- Geology of the Yazd Block

The Mehdiabad deposit belongs to the Central-East Iranian Microcontinent (CEIM; Takin, 1972), which forms the central part of the present-day Iran Plate (Fig. 1). The CEIM consists, from east to west, of three structural units, i.e., the Lut, the Tabas, and the Yazd Blocks (Fig. 1) (Berberian and King, 1981; Berberian et al., 1982, and Davoudzadeh, 1997; Alavi, 1994). The stratigraphy of the Yazd Block is globally similar to the other parts of Central Iran; it contains sequences of Neoproterozoic to Middle Triassic sedimentary rocks, covered by extensive coal-bearing Upper Triassic to Mid-Jurassic sandstones, siltstones and shales (Shemshak Formation) (Fig. 2). The thick Jurassic sequence is well exposed in the Tabas Block and here it has been studied in details (see Nabavi, 1972a,b; Aistov et al., 1984; Wilmsen et al., 2003, 2009a;

Wilmsen et al., 2005; Wilmsen et al., 2014). During the Cretaceous, the CEIM was separated from the southern Eurasian margin by narrow oceanic basins (Sistan Ocean in the east, Sabzevar Ocean in the north, Nain–Baft Ocean in the west; Lindenberg et al., 1983; Tirrul et al., 1983; Dercourt et al., 1986; Philip and Floquet, 2000; Barrier and Vrielynck, 2008a,b). These oceanic basins opened during the **Early** Cretaceous (‘Neocomian’) (e.g., find of Valanginian tintinnids and ammonites from the Sabzevar Zone; Seyed-Emami et al., 1972) and their development may be connected to an inferred post-Triassic tectonic rotation of the CEIM around a vertical axis of about 135° with respect to Eurasia (e.g., Davoudzadeh et al., 1981; Soffel et al., 1996). Rotation most probably took place **predominantly** in post-Jurassic times (Esmaeily et al., 2007; Bagheri and Stampfli, 2008; Wilmsen et al., 2009a), although the timing and the amount of rotation are still debated (e.g., Muttoni et al., 2009a,b). Cifelli et al. (2013) demonstrated that the movements at the block-bounding fault between the Tabas and Lut Blocks of the CEIM changed from extensional during the Jurassic to a right-lateral transpression between the Early Cretaceous and the Paleocene. Mattei et al. (2012) have also documented evidence of significant Early Cretaceous and Neogene counterclockwise rotation of the CEIM. The various small oceans around the CEIM closed at the end of **the** Cretaceous and during the Paleocene, due to the closure of the Neotethys followed by the collision of the Arabian Plate (e.g., Tirrul et al., 1983; Dercourt et al., 1986; Stampfli and Borel, 2002; Agard et al., 2005; Rosetti et al., 2010).

2-2- Local geological setting of the Mehdiabad deposit

In the southern Yazd Basin, a thick **so-called** Lower Cretaceous Sedimentary Sequence (LCSS) is exposed, resting unconformably on the Jurassic Shir-Kuh granite and **on** metamorphic rocks of the Shemshak Group (Fig. 3). In the following **chapter**, the Lower Cretaceous

Formations of the Yazd area, identified by Nabavi (1972a,b; see also Majidifard, 1996) will be briefly described from base to top (Figs. 4 and 5).

2-2-1- The Sangestan Formation

The Early Cretaceous sequence in the southern Yazd Basin and at the Mehdiabad deposit started with the terrigenous sediments of the Sangestan Formation (Fig. 4), which unconformably overlies the metasediments of the Shemshak Group and the Shir Kuh granite (Figs. 4 and 5). The lower part of the Formation is exclusively siliciclastic (reddish conglomerate, sandstone and siltstone) reflecting semi-arid continental depositional conditions and upwardly-decreasing relief energy (Wilmsen et al., 2014) (Figs. 4 and 5A). Towards the top, a marine influence is indicated by evaporites and carbonates, in part containing restricted marine biota (calcareous algae, agglutinating foraminifera) (Wilmsen et al., 2014). The carbonates from the Upper Sangestan Formation are probably of Hauterivian age (Bucur et al., 2003). The Sangestan Formation is mostly equivalent to the Barremian, with a diachronic top due to the Hauterivian transgression (Wilmsen et al., 2014).

2-2-2- The Taft Formation

The Sangestan Formation is conformably overlain by organic carbon matter-rich shale, silty shale, sandy limestone, dolomite and siltstone of the Barremian to probably Aptian Taft Formation (Figs. 4 and 5) (Nabavi, 1972a), consisting of thick-bedded to massive rudists, algae, and orbitolinid-bearing shallow-water facies (e.g., Bucur et al., 2012; Schlagintweit et al., 2013a). The thickness of the Taft Formation reaches up to 400 m (section at Forouzandeh; Fig. 5C) and this series is part of the Lower Cretaceous carbonate platform system of the Yazd Block (Wilmsen et al., 2013a). On the basis of benthic foraminifera, Schlagintweit et al. (2013a,b)

proposed an Early Aptian (Bedoulian) age of the lower part of the Taft Formation. However, the base of the Taft Formation may well range locally into the Barremian (see Bucur et al., 2012). The Taft Formation includes two units (Fig. 5A). The lower unit (70–85 m thick) comprises organic carbon matter-rich shale, silty shale, silty limestone, dolomite and siltstone (Figs. 5A and 5D), whereas the upper unit (315–330 m thick) includes limestone and thick-bedded dolomite (Fig. 5A).

The lower calcareous siliciclastic rocks of the Taft Formation conformably overlie those of the Sangestan Formation (Figs. 5A and 6A) in the Mehdiabad area. The lower unit displays abrupt changes in thickness, attaining <70 m in the ER but tapers rapidly westward (80 m) as it approaches the edge of the Back-Hill Fault. The transition from the lower unit to the upper unit rocks of the Taft Formation implies major changes in the sedimentary regime, from dominantly siliciclastic (lower unit) to calcareous (upper unit) (Fig. 5A). This change indicates the sag phase of the rift basin evolution (Betts et al., 1998; Large et al., 2002). The main ore horizon of Zn-Pb-Ba (Cu ± Ag) mineralization (ore horizon I) in the Mehdiabad deposit occurs within the lower unit and at the base of the upper unit of the Taft Formation (Figs. 4 and 5A).

2-2-3- The Abkuh Formation

The upper part of the Taft Formation is covered by 20 to 30 m thick cherty limestone, which forms the base of the Abkuh Formation (Figs. 4 and 5A). It is affected by irregular dolomitization, which cuts across the stratification. Above the cherty limestone is a unit of dolomitic shale with calcarenite and chert-bearing limestone, generally identical to those seen in the drill holes. The vertical transition between the cherty limestone and the dolomitic shale is gradual and is marked by a decrease of the chert content. Lenses and olistoliths of parareef

limestone (1-10 m size) are common in the upper part of the unit. A variably thick (0 to 60 m) level of para-reef limestone is also developed towards the top of the sequence. The lenticular shape of the unit is probably due to syn-sedimentary deformation, further inverted during tangential tectonic events. The top of this unit consists of 60-80 m thick conformable limy shale and laminar limestone (Fig. 4). Ore horizon II (non-sulfide ore) is located in the upper part of this unit (Figs. 4 and 5A). Chert-bearing bedded limestone, massive limestone and limy shale form the upper part of the Abkuh Formation (Fig. 4).

2-3- Main features of the Mehdiabad deposit

The Zn-Pb-Ba (Cu - Ag) mineralization of the Mehdiabad deposit occurs along two horizons that extend over a length of at least 3.4 km (Fig. 6). The trend of the mineralization is SW–NE, dipping 20°-30° to the northwest, consistent with the bedding of the host rock. The sulfide and non-sulfide ore of the ore horizon I (OH I) is hosted by the organic carbon-rich shale, silty limestone, dolomite and siltstone of the Taft Formation, whereas non-sulfide ores of the ore horizon II (OH II) are hosted in the limy shale and thinly bedded limestone of the Abkuh Formation (Fig. 6A, B) (Reichert et al., 2003). The main part of the ore body or OH I, the so-called Black hill ore (BH), Central valley orebody (CVOB) and East Ridge (ER), is located in a depression surrounded by hills and mountains (Fig. 6B). The second ore horizon (OH II) is only observed at the Calamine Mine (CM) (Fig. 6A). The BH, ER and CM expose completely oxidized ore (Figs. 6, 7 and 8A), whereas the CVOB mainly consist of sulfide mineralization. The restrictive occurrence of orebodies in the Taft Formation and in the Abkuh Formation documents the large vertical (stratigraphic) extent of the mineralization (Fig. 6). Detailed geological mapping of host rocks and synsedimentary structures in the Mehdiabad deposit shows

228 that Zn-Pb-Ba (Cu $\pm$ Ag) mineralization occur to the east of the Black Hill syn-sedimentary fault
229 in the LCSS (Figs. 4, 6A).

230 2-3-1- Ore Horizon I (OH I)

231 The BH vein and brecciated mineralization occurs adjacent to the Black-Hill syn-
232 sedimentary fault (Fig. 8B). The BH ore consists of irregular veins of barite, quartz, chalcopyrite,
233 pyrite, sphalerite, galena, Zn-Pb carbonate, and dolomite, but also limonite and hematite, so that
234 it is often called the 'Black-Hill gossan'. The veins cut siltstone, dolomite, and shales of the
235 lower unit of the Taft Formation. Veins range from <1 mm up to 80 cm in width and constitute
236 up to ~80 % of the host rock. The abundance of veins increases stratigraphically upward, toward
237 the barite ore zone. Stockwork veins similar to those at Mehdiabad have been interpreted as
238 conduits through which mineralizing fluids vented onto the seafloor (Goodfellow and Lydon,
239 2007); therefore, this stockwork system likely represents the feeder zone of the deposit.

240 Toward the east, the BH mineralization changes into the CVOB (Fig. 6A). This lateral
241 transition is gradual and is marked by a progressive increase of breccias and of the sulfide
242 content. The organic matter-rich shale, dolomite and silty limestone of the Taft Formation (Fig.
243 5D), hosting the sulfide mineralization in the CVOB, are characterized by intensive and
244 extensive brecciation. Coarse angular sulfide-matrix breccias occur throughout the CVOB but
245 are spectacularly developed near the base where fragments of shale, dolomite, siltstone, and a
246 variety of other material including barite, and sulfide breccia occur in a fine-grained sulfide
247 matrix (Figs. 8, 9 and 12). Evidence that these breccias are related to the mineralizing process
248 and not unrelated faulting includes: interlaying of the breccias with nonbreccia-textured sulfides
249 (Fig. 12), breccias which are cut by and show alteration effects from later sulfide veins (Figs.

16D and 19A), and the fact that breccias are internal to the deposit and not spatially related to faults in the CVOB (Fig. 12). The well-developed sulfide-matrix breccia appears matrix-supported; clasts compose 20 to 75 percent of the volume of the rock. Clasts are angular to subrounded and range in size from <1 to >100 mm; the estimation of the maximum size of clasts is limited by the diameter of drill core (Figs. 18, 19 and 20). The most abundant ore minerals within the CVOB are sphalerite, galena, barite, pyrite, and rare chalcopyrite. Disseminated sulfides commonly occur as low-grade mineralization within the dolomite and the lower part of the Taft Formation. Towards the west, sulfide mineralization of the CVOB grades into synsedimentary stratiform horizons of ER (Fig. 6A).

The ER orebody mainly consists of stratabound sulfide and non-sulfide mineralization developed within the lower unit (calcareous shale, siltstone, silty limestone) of the Taft Formation (Figs. 6B, 7, 8B). It extends about 2 km along a SW-NE trend and dips at 10-30 degrees to the NW (Figs. 4 and 6B). The ER orebody, known from surface exposures and drilling, is a nearly flat lying, elongated stack of mineralized lenses. This zone consists of iron oxides (hematite, limonite), manganese oxi-hydroxides and Zn-Pb carbonates. The occurrence of galena, sphalerite, and pyrite in the drill cores sampled at 100m below the surface suggests the presence of a sulfide zone at depth. The textures of sulfides in the shale and siltstone are laminated and bedded. The ER occurs between the sandstone and oolitic limestone of the Sangestan and the silty limestone of the Taft Formation (Figs. 4 and 6B).

2-3-2- Ore Horizon II (OH II)

The CM represents the highest parts of the exposed mineralization of Mehdiabad (Fig. 6A). In contrast to the mixed main sulfide/non-sulfide orebody of the CVOB, the CM contains no

zinc- and lead sulfides. The CM is exposed in numerous adits, which follow the folded and duplicated ore horizon at different mining levels. The non-sulfide zinc-lead ore occurs on mainly three different levels within the Abkuh Formation, which is the result of fold repetition. Overall, the mineralization is predominantly hosted within the limy shale with intercalations of laminar limestone of the Abkuh Formation and particularly at the contact of the limy shale to the chert-bearing bedded limestone (Fig. 7). The host rocks to the CM are intensively faulted, brecciated, and locally mylonitized. Zinc-lead mineralization mainly consists of hemimorphite, smithsonite, and hydrozincite, locally associated with iron oxi/hydroxides (Reichert, 2007; Maghfouri, 2017). Only few sulfide minerals have been observed by microscope (Maghfouri, 2017). By analogy to OH I, the pre-existing sulfide mineral association of the OH II zone was probably galena-sphalerite-barite-pyrite. In the following we will only focus on the Ore Horizon I and readers may refer to Maghfouri et al. (2015) for complementary informations on OH II and CM.

3- Methodology

A field campaign and sampling were done during the autumn of 2015. To assess the geochemical characteristics of 250 systematic samples (least fractured and altered) of the ore-bearing zone and of host rocks were collected from the deposit for polished thin section examination. In order to correctly characterize their chemical compositions, 13 least-altered and fractured samples were chosen for major, trace, and rare-earth-elements (REE) analyses. Also 132 ore-samples were analysed by Mehdiabad Mining Company in the ACME Analytical Laboratories. Rock chips (~400 g) from each sample were carefully selected, washed with distilled water at room temperature, and then dried. Finally samples for whole-rock analyses were crushed and powdered in a tungsten carbide swing mill. Bulk-rock chemical analyses of major, trace, and REE elements were determined for all samples by Inductively Coupled Plasma-Emission

Spectrometry (ICP-ES) and Mass Spectrometry (ICP-MS) at ACME Analytical Laboratories, Vancouver, Canada. For the major elements, the detection limit is between 0.002 and 0.01 wt.%, 0.02–2 ppm for the trace elements, and 0.1–0.01 ppm for the REE elements. Major and trace element data are presented in Table 1 and REE elements data in Table 1.

Sulfide and sulfate minerals were separated from monomineralic samples using a microdrill and a binocular microscope. The microdrill was cleaned between each sample extraction to avoid contamination. Six $\delta^{34}\text{S}$ determinations by laser ablation-mass spectrometer were conducted at the Centres Científics i Tecnològics de la Universitat de Barcelona, Spain and in the Institute of Geochemistry, Chinese Academy of Sciences, Guizhou, China. The analytical procedure is similar to the one described in Huston et al. (1995). A 200- μm -thick polished slice of each sample was used for analyses. CO_2 separation was conducted on gas derived from samples.

4- Results

4-1- Sedimentary and geochemical characteristics of the host rocks

Most of the sedimentary rocks (Taft Formation) in the immediate vicinity of the Mehdiabad deposit are fine to very fine grained, gray-black, indistinctly to well-laminated organic carbon-rich shale, siltstone, dolomite, and limestone beds. These latter ones host the mineralization in the two horizons of the Mehdiabad deposit (Figs. 7 and 9). The main facies consists of organic carbon-rich shale and siltstone marked by interbeds of grey dolomitic siltstone, dolomite and thin-bedded black (carbonaceous) limestone (Fig. 9). The thickness of individual beds of organic matter-rich shale, siltstone and dolomite range from several centimeters to about 60 m, with an average of 40-50 m (Figs. 9A-9D). The shale and siltstone rocks may be calcareous, non-calcareous and/or carbonaceous. They are indistinctly to well laminated (<1 mm to 20 cm) (Fig.

9). Thin, sandy lenses or beds containing pale grey calcareous grains are locally observed (Fig. 9E). Organic matter-rich shale and siltstone are composed of quartz, carbonate, chert fragments, and carbonaceous material, occurring as sub-micron sized inclusions, organic matter nodules, pyrite and Fe-oxide minerals (Fig. 9F). The presence of abundant carbonates indicates that the sediments were not deposited at depths below the carbonate compensation depth (Peter and Scott, 1999). Sedimentary structures include normal layering, scours, pebble dents, and concretions composed of pyrite, calcite, ferroan dolomite and rare blebs of sphalerite. Soft-sediment deformation, represented by slump structures, indicates that the basin was tectonically active during sedimentation. Organic matter-rich shale and siltstone contains, very fine to coarse-grained, disseminated pyrite and/or fine grained sphalerite, galena and barite, of either syngenetic or diagenetic origin (Fig. 9).

The geochemical composition of the host rock shales from the Taft Formation is reported in Table 1. Unaltered shale samples from the Taft Formation have Mn and Fe contents ranging from 231 ppm to 6391 ppm and from 1.46% to 4.12%. respectively (Figs. 10 and 11). The V/Cr ratios for shale samples from the Mehdiabad deposit range from 0.60 to 4.43, whereas the V/(V+Ni) ratios range from 0.67 to 0.91 (Fig. 11). Samples are also marked by high U contents and the Th/U ratio ranges from 0.14 to 1.53, with an average value of 0.50.

4.2- Discrimination of ore facies and their distribution in the deposit

In the Central Valley, the sulfide orebodies of the Mehdiabad deposit do not crop out at the surface, and drill core logs and lithogeochemical data have been used to reconstruct a cross-section of the deposit (Fig. 12). The major sulfide mineralization is located east of the Black-Hill synsedimentary fault (Fig. 4). The sulfide zone at Mehdiabad (OH I) ranges greatly in thickness,

from few meters to more than 300 m. It displays a lens shape with a relatively flat top and a convex base (Fig. 12). The greatest thickness of high-grade mineralization (>7.52% Zn) is just to the east of the Black-Hill Fault, in the CVOB (Fig. 12). This zone also has the highest Zn grades (10–17%), highest Ag concentrations (21–51.7 g/t) and highest Pb grade (1.9-6%) (Maghfouri, 2017). Sulfide mineralization occurs as a single wedge within the organic matter-rich shale, siltstone, silty limestone and dolomite (Fig. 12). The western part (BH and CVOB) of the deposit has stratabound geometry, but toward the northeast (ER), as thickness decreases, the mineralization becomes stratiform and concordant with the host sedimentary rocks (Fig. 12).

The Mehdiabad deposit consists of five ore facies *types*, which are, from proximal (BH and CVOB) to distal (ER) relative to the synsedimentary fault (Fig. 12): (1) a stockwork zone, formed by sulfide veins accompanied by carbonate or quartz; (2) *massive*-replacement ore, formed by a massive sulfide zone and sulfide breccias; (3) *bedded* ore, consisting of a banded orebody made of pyrite, sphalerite, and disseminated galena; (4) a barite ore facies, which predominantly occur near the top of the sulfide mineralization; and (5) a distal facies, formed by chert beds with barite and disseminated pyrite.

- The stockwork or stringer zone

This zone consists of an irregular vein network cutting across the altered and brecciated siltstones, dolomites, and shales (Fig. 13). The veins are composed of fine grained barite, quartz, pyrite, chalcopyrite, sphalerite, galena, and dolomite (Fig. 13). This type of mineralization occurs adjacent to the Black-Hill Fault (Fig. 12). Dolomite and quartz occur in the core of the veins *whereas* sulfides are mainly found in the outer part. Veins range from <1 mm up to 1 m wide and may constitute up to ~20 % of the host rock (Fig. 13). The *number* of veins increases upward,

progressively passing into massive-replacement ore. Debris flow and abrupt facies changes occur adjacent to the stockwork zone (Fig. 12). Sulfide-rich veins generally present an alteration halo (silicification and dolomitization) and a partial replacement of pseudocubic pyrite by chalcopyrite and sphalerite is suggested by the irregular and crenulated grain boundaries in the pyrite-sphalerite aggregates (Fig.13). Copper contents in the stockwork zone may reach up to 2 wt% (Maghfouri, 2017). Sulfide veins also crosscut the barite ore facies and massive sulfides (Fig.13) suggesting multiple stages of stringer ore mineralization. Five main generations of veins and veinlets were identified within the stringer zone, based on the textural relationships and mineral paragenesis of the ore veins (V1 to V5) (Fig.13). These veinlets from early to late are (V1): chalcopyrite, quartz and barite-rich vein-veinlet's; (V2): sphalerite, galena, pyrite and barite-rich vein-veinlet's; (V3): sphalerite and galena-rich vein-veinlet's; (V4): barite-rich vein-veinlet's; and (V5): pyrite-bearing calcite vein-veinlet's. Banding, crustification, and comb textures indicate deposition in open space (Fig. 13). Wall rock fragments are commonly included in the veins and resulted in formation of cockade ore textures. Sulfide-cemented breccias are developed in areas of intense veining, particularly near the base of the deposit.

- The massive-replacement ore facies

The massive-replacement ore consists of a high-grade orebody that forms the thickest part of the deposit (Figs. 12 and 14A), principally in the CVOB, to a lesser degree, on the eastern margin of the BH. This facies comprises massive zones, replacement zones, brecciated sulfides, and a very irregular network of sulfide veins and veinlets (Figs. 14B to 14D). Mineralization in this facies is hosted by a series of stacked, planar-wedge shaped breccia bodies (Figs. 12 and 14). Some of the breccia units are dolomitized and only the lowermost breccia contains stratiform massive Zn-Pb-Fe sulfides. Dissolution breccias are perhaps the most distinctive type of

hydrothermal replacement at Mehdiabad and are characterized by coarse-grained sulfides that both replaced and infill veins and cavities within dolomitized limestone of the Taft Formation (Fig. 12). Well-developed sulfide-bearing breccias appear to be matrix-supported and clasts form 20 to 75 percent of the rock volume. Clasts are angular to subrounded and range in size from <1 to >100 mm, but estimation of the maximum size of clasts is limited by the diameter of drill core. The clasts are angular and are partially replaced and crosscut by thin veinlets of coarse-grained sulfide, which also form the matrix of the breccia. Most of the clasts are composed of hydrothermal dolomite (Fig. 14E), although clasts of shale, not dolomitized limestone of the Taft limestone, early sulfide aggregates, and massive barite were also observed. The bulk of the well-developed sulfide-bearing breccia occurs as a series of roughly stratiform lenses at or near the base of the dolomitized Taft Formation, in the hanging wall of the Black-Hill Fault.

The massive-replacement ore facies shows mineralogical zonation from bottom to top with an upward decrease in chalcopyrite and galena and increase in sphalerite, pyrite and barite (Fig. 12). Based on mineralogical and textural features, the massive-replacement ore facies is divided into five sub-facies (Figs. 12 and 14): (1) a chalcopyrite-sphalerite-galena-barite sub-facies (CSGB) (Figs. 12 and 14G), which occurs on the margin of the massive-replacement ore, especially in the BH area; (2) a sphalerite-galena-barite sub-facies (SGB) (Figs. 12, 14C, 14E and 14F); (3) a pyrite-sphalerite-barite-galena sub-facies (PSBG), mostly observed in the central part of the massive-replacement ore zone (Figs. 12, 14B, 14D and 14H); (4) a massive oxide Zn-Pb-Ba ore sub-facies (Figs. 12 and 14A); and (5) a barite ore sub-facies, located at the top of the system (Figs. 12 and 14A).

Stratiform beds and disseminations of barite also appear higher in the Mehdiabad stratigraphic section as well as locally in the massive-replacement ore facies within dolomitic

shales of the Taft Formation. The barite ore is composed of stratiform lenses, which are typically massive, but the bedding is locally defined by the presence of base metal sulfide or oxide layers (Fig. 14A). Fine- to coarse-grained euhedral to anhedral barite grains are intergrown with disseminated sphalerite, galena, and pyrite. Sulfide minerals also form thin to thick layers enclosed by barite. The massive sulfide-bearing barite lenses and beds locally grade in to quartz and apparently formed on the sea floor. Such textural evidence indicates that the barite predated and was partially replaced by the sulfides, but also co-precipitated with sulfides (Maghfouri, 2017). Barite ores of Mehdiabad can be divided into four textural types: massive, laminated, banded, and nodular barite ores and most of these are light gray in color. The massive ores are composed of high-grade (>85 %) BaSO₄ with minor sulfides and quartz. The barite crystals range from 0.05 to 0.18 mm. Generally, this massive variety is located at the top of the ore beds (Fig. 14A). Laminated barite commonly overlies the bedded sulfide ore facies and is distinguished by the number of sulfide and oxide layers interlaminated with barite. Banded barite ores, occurring at the base of the ore beds are of low grade, with relatively high sulfide contents.

- The bedded ore facies

The facies (bedded barite) is located above and to the east (bedded sulfide) of the massive-replacement ore facies in the CVOB (Fig. 12) and in the ER (Fig. 15). It is characterized by laminated sphalerite, galena, and iron sulfide (pyrite and marcasite) (Fig. 15). In some core samples (e.g., DDH 806, 810), iron sulfides predominate at the top of the zone, whereas the amount of base metal sulfides increases with depth. The lateral transition between the bedded ore and the massive-replacement ore facies is generally gradual and is recognized by an increase in laminae. The contact with the underlying silty limestone and dolomite is usually sharp (Figs. 16B and 16C). This facies displays stratiform banded sulfide-rich layers (Figs. 16D to G), concordant

with the host rock bedding (Fig. 16A). Pyrite, galena, and sphalerite are the dominant sulfide minerals (Fig. 16). Fine bedding defined by the distribution of sulfides associated with silt and organic matter may be observed from a macroscopic to microscopic scale (Fig. 16). The bedded ore facies can be divided into low-grade (1–2 % Zn + Pb) and high-grade (>4 % Zn + Pb) units (Fig. 16). At the microscopic scale, pyrite forms framboidal aggregates. Sphalerite laminae are less common than pyrite laminae (Fig. 16D); they are only ubiquitous in the high-grade part of the bedded ore facies.

- The distal facies

The banded silicic dolomite is only exposed in the eastern part of the deposit (ER; Figs. 4 and 10A) and is associated with strong Fe and Mn anomalies (Figs 16 H, I, J). This facies has no economic value and represents the most distal part of the hydrothermal deposit. The boundary between the bedded ore and the distal facies is gradual and is indicated by the absence of sphalerite and the increase of disseminated pyrite with respect to laminated pyrite (Figs 16 I, J, K). Disseminated barite occurs in association with Fe- and Mn-rich dolomite in the distal facies.

4-3- Relationships between mineralization, host rocks and faults

Similar to the sedimentary host rocks, the sulfides display characteristic layering on a macroscopic and microscopic scale (Figs. 16, 17, 18 and 19). Despite complex banding and a high variability of textural and compositional variations within the laminae, we may distinguish three end-member types of microbands in the ore (Figs. 16 and 17). The organic matter-rich shale layers (Type 1) are commonly less than 100 µm thick (Figs. 17A and 17B); they are principally composed of illite, carbonaceous matter, pyrite, and quartz; and are interpreted to be the product of normal pelagic sedimentation from the anoxic water column. The shale and silt

layers (Type 2) are commonly 100 to 500 μm thick, but laminae up to several centimeters thick are also present (Fig. 17E). The zinc-lead-iron-rich sulfide layers (Type 3) are commonly 50 to 300 μm thick (Figs. 17C and 17D) and consist of uniform mixtures of sphalerite-galena-pyrite with minor barite and quartz and rare chalcopyrite. Sphalerite and pyrite are the main sulfide minerals and the sphalerite/galena ratio varies from 2:1 to 9:1. Although it is common for the three end-member microband types to be regularly interlayered in a given sample (Figs. 16 and 17), mixture of these and end members may also occur (i.e. type 1 + type 3 in figs. 13B and 14A; type 2 + type 3 in figs. 16D, 16E, 17C and 17I D). From ore microscopy, different types of pyrites were observed within the shales collected from the drill holes. We may distinguish colloform pyrite (Fig. 17F) from framboid pyrite (Fig. 17G). Beds of galena and sphalerite are also present (Figs. 16G and 17H).

Rounded and subrounded sulfide clasts are common within the thick ($> 1\text{cm}$) turbidite beds (Figs 18A and 18B); these are interpreted as the result of a reworking of sulfide layers originally deposited on the sea floor. Compactional load-cast features are also common in the turbidites rocks, locally bending or folding the sulfide-rich layers (Figs. 18D, 18E and 18F). Detrital clasts of carbonate and lithic fragments occur at the base of the sedimentary rocks and they display evidence of compressive deformation, beneath the base metal layer. Flame structures are also visible in the sphalerite-galena-pyrite beds, extending upward into the mass flow units. Sulfide microbreccia layers are common in the different ores (Fig.18G). These layers are composed of rounded to angular clasts of sulfide minerals enclosed in a matrix of sediments with disseminated sulfides (Fig. 18G). They are typically non-graded, measuring 0.7 to 2 mm in thickness, and indicate open-cast reworking of a proximal source of preexisting sulfides.

Mineralization occurs to the east of the Black-Hill (BH) Fault (Fig. 8E), which was intersected by drilling. In drill core, evidence of hydrothermal alteration, marked by dolomitization, silicification, and sulfide mineralization, was recognized as extremely variable (Fig.13A). This corresponds to the stockwork facies, described in the previous section. Whereas the thickness of the sedimentary units of the Sangestan Formation beneath the Taft Formation is relatively constant, the thickness of the Taft Formation increases in the hanging-wall of the BH fault. As the compressive deformation related to Tertiary inverted tectonics was quite gentle, such present-day variations probably represent the original deposition, most likely controlled by the syn-sedimentary activity of the BH normal fault. It is also worth to notice that the thickest and highest grades of Zn-Pb mineralization, as well as pervasive hydrothermal alteration occur within the Taft Formation in the hanging- wall rocks of this major fault as well (Figs.12 and 19). Exploration drilling along the orebody also indicates that major facies changes occur in the Taft Formation, with shallow-water shelf facies to the east of the BH fault and basinal rocks in the CVOB area.

Additional evidence of the syn-sedimentary activity of the BH fault is the presence of mass flows (Fig.20). The latter correspond to sedimentary breccias that are common throughout the Taft Formation (Fig.20). To the west and to the south, the mass flows become thicker as the size of fragments increases; this indicates that the clasts are sourced from mass shedding along the tectonically active BH fault (Fig.20).

4-4- Sulfur isotopes: results and discussion

Sulfur isotopes at the Mehdiabad deposit were measured from 52 samples of pyrite, chalcopyrite, barite, sphalerite, and galena collected from all ore facies except the distal facies.

The analytical procedure is given in the supplementary material and the results of sulfur isotope analyses are listed in the table 2 and shown on Figure 17. The sulfides have a wide range of $\delta^{34}\text{S}$ values from -22‰ to 22.4‰ , with a majority at about -14 to 4 per mil (Fig. 21). Extremely low $\delta^{34}\text{S}$ values in the range of -20.1‰ to -22‰ are only observed in bedded ore facies pyrite (Fig. 21A), whereas the massive ore and stockwork sulfides have higher $\delta^{34}\text{S}$ values ranging between -13.7‰ and $+4.6\text{‰}$ (avg; -4.2‰).

The $\delta^{34}\text{S}$ values of sphalerite ($n = 11$) fall within the range of -18‰ to $+4.6\text{‰}$. There is a close correlation between texture and $\delta^{34}\text{S}$ values. Fine-grained, framboidal and colloform sphalerite has primarily negative values (-17.53‰ $n = 3$), whereas coarser-grained sulfides found at the base of the deposit in the massive-replacement ore (-6.27‰ for $n = 4$) and in the stockwork zone (-1.35‰ for $n = 4$) have significantly higher values (Figs. 21B). A similar pattern is observed for galena, whose $\delta^{34}\text{S}$ values range from -18.1‰ to $+4.4\text{‰}$ ($n = 16$). Galena from the massive-replacement ore facies (-9.34‰ for $n = 5$) and from the feeder zone (-2.72‰) are isotopically different from the galena locally intergrown with framboidal pyrite of the bedded ore (-17‰ to -18.1‰).

$\delta^{34}\text{S}$ values in chalcopyrite ($n = 9$) range from -9.6‰ to $+4.6\text{‰}$ (Fig. 21B). Vein-veinlet chalcopyrite hosted by hydrothermal dolomite in the deep feeder zone has the $\delta^{34}\text{S}$ between 4.6‰ and -7.3‰ per mil, whereas massive-replacement ore has a mean $\delta^{34}\text{S}$ of -3.6‰ . Barite intergrown with sulfides in the feeder zone ($n = 3$) has $\delta^{34}\text{S}$ values ranging from $+22.4\text{‰}$ to $+22$ per mil. A similar range of values was obtained in the bedded ore ($+17.8\text{‰}$ to $+22.3$ for $n = 4$) and in the massive-replacement ore facies, ($+19.65\text{‰}$ to 20.1‰ for $n = 3$).

5- Discussion

5-1- Conditions of deposition of Mehdiabad host rocks

Interbedded black siltstone documents the ambient depositional conditions and hosts the ores and mineralization at Mehdiabad. The dolomite beds represent a small volume in the Mehdiabad Basin. This facies is characterized by an accumulation of fine-grained hemi-pelagic suspensions in a quiet, reduced, sub-wave base subaqueous environment. Thicker, discrete fine-grained shale intervals are rich in pyrite and organic matter. This sedimentary facies, which is the most common at the base of the Taft Formation and in the middle part of Abkuh Formation, is considered to be relatively poor in clastic sediment input, and is interpreted to indicate the maximum of water depth and/or shoreline retreat in the basin.

Several geochemical indices based on redox-sensitive major and trace elements have been used to determine the paleo-oxygenation conditions of the water column (Hoffman et al., 1998; Jones and Manning, 1994; Quinby-Hunt and Wilde, 1994; Whitehead, 1973). Among the major elements, Fe and Mn can be used to indicate bottom water conditions as their solubility is affected by ambient pE and pH (Whitehead, 1973). Quinby-Hunt and Wilde (1994) used solubilities of Mn and Fe to determine the paleo oxygenation levels and the prevailing oxidizing agent in the water column. In oxic bottom water conditions, oxygen is the primary oxidizing agent; Quinby-Hunt and Wilde (1994) argued that low Mn and Fe solubilities would favor their fixation in sediment, yielding high Mn and Fe contents in shales. In anoxic bottom water conditions with nitrate as the primary oxidizing agent, Mn solubility increases whereas pE decreases, resulting in low Mn contents in the sediments. In contrast, Fe remains bound in oxide minerals within the sediments, although high Fe solubility at very low pE values may lead to a decrease of the Fe content in sediments. Although low Mn and high Fe concentrations may characterize nitrate-reducing conditions, low Mn and high Fe may also occur in anoxic

conditions in which sulfate is the primary oxidizing agent, as Mn remains dissolved in the water column and reduced sulfur fixes Fe in the sediments. The low Mn contents in the unaltered shale of the Taft Formation are consistent with a deposition in anoxic bottom waters (Fig. 10A), whereas the Fe contents may support oxic or anoxic sulfate-reducing conditions (Fig. 10). However, Mn was shown to have been mobile during hydrothermal alteration, compared to Fe, thus caution must be taken to use this proxy in such polyphase mineralizing system.

In addition, the low oxygen levels are strongly limited the benthic activity, and ichnofabric criteria cannot be used to discriminate between oxic (>2.0 ml O_2 /l H_2O), suboxic (0.5 – 2.0 ml O_2 /l H_2O), dysoxic (0.2 – 0.5 ml O_2 /l H_2O) and anoxic (<0.2 ml O_2 /l H_2O) conditions. Hoffman et al. (1998) proposed the V/Cr and V/(V + Ni) ratios, which are sensitive to anoxic and dysoxic conditions. Whereas V, by bounding to organic matter, is considered as a useful indicator of its preservation (Brumsack, 1986), Cr, which substitutes for Al in clays or occurs as chromite, is associated with the detrital fraction of the shale (Jones and Manning, 1994), Ni is preferentially incorporated into the sulfide minerals. The V/Cr and V/(V+Ni) ratios of the Mehdiabad shales indicate anoxic to dysoxic conditions (Fig. 11). The high U content of the Mehdiabad shales is typical of sediments deposited under anoxic conditions (Nath et al. 1997), and results from the reduction of U^{+6} into U^{+4} , which is insoluble and deposited in the sediments. The low value for the Th/U ratio (av. 0.85) shows that no fractionation occurred between U and Th, unlike what is classically observed in shallow oxic environment (Whittaker and Kyser 1993; Wingal and Twitchett, 1996).

Most SEDEX deposits are hosted by basinal marine, reduced facies, fine-grained sedimentary rocks that consist mostly of carbonaceous siltstone and shale (Large et al., 1996; Kelley et al., 2004a,b; Leach et al., 2005; Goodfellow and Lydon, 2007; Leach et al., 2010;

Wilkinson, 2014; Gadd et al., 2016a,b). These sedimentary rocks generally represent pelagic and hemipelagic sediments. In some deposits (e.g., Sullivan), pelagic and/or hemipelagic sedimentary rocks are interbedded with turbiditic siltstones and sandstones, and locally include coarse-grained clastic sediments, **derived** from the erosion of uplifted blocks during extensional tectonism (Goodfellow and **Lydon, 2007**; Magnall et al., 2015; Gadd et al., 2016b; Magnall et al., 2016; Gadd et al., 2016a). In addition, the geochemical redox proxies used to characterize the shales hosting the mineralization support an anoxic **environment** and our results are very similar to what has been reported for many other SEDEX deposits (Goodfellow and Lydon, 2007; Sáez et al., 2011; Yarmohammadi et al. 2016; Boveiri et al., 2017). In the case of the Irish-type or MVT deposits, the dominant **lithotypes** are limestone and dolomite (Lee and Wilkinson, 2002; Wilkinson, 2003; Reed and Wallace, 2004; Wilkinson, 2010; Elliott, 2015); MVT deposits generally form **in** reduced environments (Pfaff et al., 2010).

5-2- Textural and metal zonation

One of the most characteristic features of SEDEX deposits is the vertical and lateral zonation of textures, minerals, and metals (Goodfellow and Lydon, 2007). Deep drill holes have shown that the Taft Formation is mineralized in at least three areas (BH, CVOB and ER) in the hanging wall of the major Black-Hill synsedimentary fault (Figs. 6A, 12). The largest **of** such sulfide zones located **so** far are within the CVOB between the BH and ER orebodies (Maghfouri, 2017) (Fig.12). In this deposit, there is a gradual change, away from the stockwork zone (or BH synsedimentary fault), from veins and massive replacement textures in the massive-replacement ore (CVOB) to sedimentary textures (laminated, disseminated, and framboidal) in the bedded ore (ER) and **in the** distal facies (Maghfouri, 2017) (Fig. 12). This **systematic** trend is accompanied

by a decrease in the number and thickness of individual sulfide beds (Fig.12). The ore thickness and grade in this deposit decrease also systematically away from the Black-Hill Fault (Fig.12).

The deposit is also marked by lateral chemical zonation. In terms of metal ratios, there is a decrease in the Pb/Zn ratio from the massive-replacement ore, with a maximum value of combined base metals (Zn+Pb=12.5 %) in the thickest part of the deposit toward the bedded ore (Maghfouri, 2017) (Fig. 12) (Table 3). There is still a gradual increase in zinc grade to the CVOB, away from to Black-Hill Fault (Fig.12). These lateral variations in zinc grade demonstrate systematic changes from as much as 40 vol percent sphalerite to about 5 vol percent sphalerite without any stratigraphic correlation. This is accompanied by a decrease in Pb grades towards the northeast of the deposit (Maghfouri, 2017). Iron is mostly contained in pyrite, and the Fe/Pb and Fe/Zn ratios, together with the Mn content increase away from the massive-replacement ore as base metals decrease (Fig. 12).

Copper is strongly zoned, and Cu-rich ores have been found close to the Black-Hill Fault (Maghfouri, 2017) (Table 3). Silver displays a zonation pattern similar to zinc throughout much of the deposit, although the highest values occur in the central part (CVOB) of the sulfide zone in the area of high lead values (Fig. 12). Barium contents are uniformly high at Mehdiabad. The highest values occur adjacent to the Black-Hill Fault (Fig.12). There is a broad overlapping lateral zonation marked from west to the northeast by the following trend $Cu > Pb > Zn > Fe$. Maghfouri (2017) interpreted this trend as a manifestation of the prevailing direction of lateral fluid flow and solubility-controlled precipitation through the mineralizing system, possibly also due to a thermal gradient or a chemical gradient independent of temperature (e.g., pH, oxidation state, or chemical kinetic effect). The relatively high Cu and low Fe in the western part of the

deposit is inferred to represent the location of fluid entry or venting site into the Mehdiabad orebody.

Moreover, the Mehdiabad deposit also shows a strong vertical zonation, especially within the massive-replacement ore (Fig. 7). There is an upward increase of Zn, Pb, and Ba/(Zn + Pb) and a decrease of Cu/(Zn + Pb), Ag/(Pb + Ag), and Cu (Maghfouri, 2017) (Fig. 12) (Table 3). The vertical geochemical variations may reflect changes in the overall abundances of ore minerals, or alternatively, they could reflect the chemical composition of the sulfides. As an example, there is a greater abundance of iron sulfide in massive-replacement ore compared with vein sulfide ore, and sphalerite in vein ores is Fe-rich compared with that in other ore types.

5-3- Mineralizing process of the Mehdiabad deposit

5-3-1- Syn-sedimentary/diagenetic versus replacement

On the basis of mineralogical and textural studies, Maghfouri (2017) considered a three-stage (Fig. 22) mineralization model at Mehdiabad. Early sulfides and Ba sulfate (Py1, Sph1, Gn1, Cpy1 and Ba1) formed by exhalation and precipitated from an overlying water column during an early diagenetic stage (Stage 1), whereas late sulfides and Ba sulfate (Py2, Sph2, Gn2, Cpy2 and Ba2) resulted from a subsurface hydrothermal replacement event (Stage 2) (Fig. 19). The third stage is seen as a covering layer of barite (Ba3) that also contains pyrite (Py3), sphalerite (Sph3) and galena (Gn3) (Fig. 12).

The early syn-diagenetic stage is characterized by the presence of framboidal pyrite (Py1) (Fig.17G), colloform pyrite (Fig.17F) and laminated and disseminated sphalerite (Sph1), galena (Gn 1) and barite (Ba 1) (Figs. 9C, 17C,D, 16G, D and 17H), commonly found in the bedded ore

and rarely in the distal facies (Fig. 22). In addition to their shape, the small size of such pyrite grains rather supports a sedimentary-diagenetic origin. The sphalerite-galena-pyrite-rich bands are intricately interlayered with shale and siltstone beds and carbonate turbidites (Fig. 16D); the boundary between mineralized and non-mineralized siltstone beds is sharp and cannot be explained by a replacement process (Figs. 16 and 17C). In addition, there is a lack of significant sphalerite and galena in many of the siltstone, whereas this lithotype is favorable for a replacement model (Ghasemi, 2006). The relations between sulfide minerals and some sedimentary structures (Fig. 18) clearly indicate that sulfide layers of Stage 1 were already present on the seafloor when the turbidite units formed.

The second ore stage occurs as massive ore, veins, and replacements, located mainly in the massive-replacement ore and stockwork zones (Fig. 19, 14 and 22). This stage is also characterized by complex mineral assemblages (Fig. 22). Detailed mineralogical and textural observation revealed that replacement is the dominant process, with early sulfide minerals replaced by late and coarse-grained sulfide minerals (Figs. 14 and 19). Textural styles include bedding-parallel to cross-cutting extensional brecciation and veining, delicate to massive-replacement of host rock, and dissolution of limestones to form open spaces that were subsequently completely filled with sulfides (Figs. 18, 19 and 20). Based on geological, textural and mineralogical studies, Maghfouri, (2017) suggested that breccia's and open space filling mineralization (second stage) at Mehdiabad formed sub-seafloor epigenetically from mineralizing solutions that flowed laterally through the still permeable sediments along deep channel-ways from the Black-Hill Fault.

Second stage of mineralization (massive-replacement) is overlain by bedded barite (Ba₃) (third stage) (Fig. 22). The vertical transition between the massive-replacement ore facies and the

stratiform barite beds is gradual and is recognized by increasing of bedded barite mineralization. This stage of mineralization is characterized by massive to laminated barite and minor pyrite (Py3), sphalerite (Sph3), and galena (Gn3) (Fig. 22).

5-3-2- The role of the Black-Hill Fault

Detailed studies of the distribution of sedimentary facies suggest that fault-related, hanging-wall subsidence began along the BH fault system in the [Early](#) Cretaceous with contemporaneous sedimentation leading to the deposition of the Taft Formation. In addition to these sedimentary processes, normal faulting also played a fundamental role for the localization of mineralization and hydrothermal alteration at Mehdiabad (Reichert et al., 2003; Reichert, 2007; Maghfouri et al., 2015; Pourfaraj et al., 2016; Maghfouri, 2017). The maximum thickness of sulfides is found close to the Black-Hill Fault, suggesting that this syn-sedimentary structure acted as the most important [conduit](#) for hydrothermal fluid flow (Fig. 12). This area, which is characterized by abundant tectonic and sedimentary [breccias](#), displays evidence for the development of a vein network. Stockwork veins, similar to those at Mehdiabad have been interpreted as conduits through which mineralizing fluids vented to the seafloor (Goodfellow, 2004); therefore, the stockwork system likely represents the feeder zone. Based on fluid inclusion studies, Maghfouri and Hosseinzadeh (2018) suggested that the [neighboring](#) Mansourabad deposit formed from basinal, hydrothermal fluids rising through synsedimentary faults to the seafloor, which might be similar in the Mehdiabad deposit. Maghfouri (2017) noticed the presence of abundant clasts of sulfides, which are identical in texture and composition to the ore lens material, immediately underlying the mass flows (Fig. 20). This provides convincing evidence that base metal sulfide mineralization at Mehdiabad occur in the uppermost level of sea-floor sediments, where the laminated ores were susceptible to be affected by [syn-depositional faulting](#) and erosion caused

by mass flows. The relationship between ore lenses, organic matter-rich shale, and mass-flow breccias strongly suggests that the BH fault system controlled both ore fluid migration and sediment breccia deposition, with mineralization limited to the deepest part of the sub-basin adjacent to the active fault system. A variety of sulfide mineral (galena, sphalerite, and pyrite) clasts occur in the mass-flow breccias (Figs. 20 a,b,c,d). In SEDEX deposits, such syn-sedimentary faults are recognized as the pathways for metal-bearing, hydrothermal fluids to enter onto the seafloor (Logan, 1979; Large et al., 1996; Large et al., 2005). Tectono-sedimentary controls have been well documented in other SEDEX deposits (i.e. the Ikalukrok Unit in the Anarraaq deposit; Kelley et al., 2004a,b). Away from the fault, the outer northeastern margin of the deposit consists of banded silicic dolomite with pyrite, chert, and manganese and is interpreted to represent more distal hydrothermal exhalites (Fig. 15).

5-3-3- Processes involved in the formation of sulfide minerals

The $\delta^{34}\text{S}$ values found at Mehdiabad are much lower than expected if sulfur would have derived from a magmatic source. The source of the sulfur is probably seawater sulfate. The barite $\delta^{34}\text{S}$ values are comparable to the results reported by Maghfouri (2017) and are identical to the range of values (+15 to +21‰) for Lower Cretaceous seawater sulfate (Claypool et al., 1980; Bottrell and Newton, 2006). This suggests that barite precipitation at Mehdiabad is related to unmodified Cretaceous marine sulfates, as inferred for barite in the Carboniferous Red Dog deposit (Johnson et al., 2004; Kelley et al., 2004a) (Fig. 21C). The transformation of sulfate to sulfide can be achieved either by bacteriogenic sulfate reduction (BSR) or by thermochemical sulfate reduction (TSR), which implies isotopic fractionations greater than 20‰ (Ohmoto et al. 1990). As temperature increases, fractionation would decrease. Nevertheless, TSR processes usually reach metastability and therefore little or no isotopic fractionation occurs between sulfur

species (Cross and Bottrell, 2000). The Mehdiabad deposit is mineralogically and chemically zoned away from the feeder zone, reflecting steep temperature gradients. Silicic alteration found around the feeder zone suggests high-temperature conditions for this part of the deposit. However, the zonation, controlled mostly by zone refining and sub-sea floor replacement in the massive ore (Goodfellow and Lydon 2007; Wilkinson et al., 2005; Kelley et al., 2004a; Johnson et al., 2004), along with the $\delta^{34}\text{S}$ difference found between the bedded ore and the massive-replacement ore, suggest that, at the margins of the system (bedded ore and distal facies), temperatures would have been low enough to sustain BSR process. Maghfouri and Hosseinzadeh (2018) reported a similar zone-refining paragenetic succession in the vent complex (massive ore) of the Mansourabad deposit. If the temperature of ore-forming fluids is less than 150° C, microbial processes can play a significant role in sulfate reduction (Southam and Saunders, 2005). The presence of framboidal and colloform pyrite in the bedded ore facies within the lower parts of the massive-replacement ore supports this interpretation. Although the development of pyrite framboids can occasionally occur by means of abiotic processes (Berner, 1969; Prol-Ledesma et al., 2010), laminated and framboidal pyrites are usually associated with biogenic deposition (Alfonso et al., 2005).

Sulfides from the bedded ore and parts of massive-replacement ore and feeder zone have $\delta^{34}\text{S}$ values between -5.2 and -22‰. Those low values alone could account for a BSR process if it occurred in a closed or semi-closed system for sulfates, but open for sulfide following a Rayleigh distillation process (e.g. Canet et al. 2005). Microbial-mediated sulfate reduction is known to have played a prominent in situ role in the formation of SEDEX deposits (Southam and Saunders, 2005; Gadd et al., 2016a) (Fig. 21C). Compared to other sulfides, pyrite has generally lower $\delta^{34}\text{S}$ values, a feature typically reported in SEDEX and Irish type deposits (e.g., Anderson

et al., 1998; Large et al., 1996; Fallick et al., 2001; Ireland et al., 2004a,b; Kelley et al., 2004a; Leach et al., 2005; Gadd et al., 2016a). Colloform pyrite (stage 1) from the massive-replacement ore and feeder zone was replaced by galena and sphalerite (stage 2) with $\delta^{34}\text{S}$ values from +3.1 to +4.4 and from +2.3 to +4.6‰, respectively. These high isotopic values of late sulfides cannot be produced by BSR in the proposed closed system for sulfate and suggest at least a contribution of TSR to the sulfide sulfur that formed these minerals.

5-4- Type of Mehdiabad deposit

Different genetic types have been proposed for the Mehdiabad Zn-Pb-Ba-(Cu-Ag) deposit, including a MVT-type (Leach et al., 2010) and a sub-seafloor replacement SEDEX-type mineralization (Maghfouri et al., 2015; Maghfouri, 2017). Except for Leach et al. (2010) review paper, there is no mention of the Mehdiabad MVT-type deposit in any other article. However, our results are in not in agreement with the scenario proposed by Leach et al. (2010) for this deposit. Based on the location of mineralization in the Taft Formation, given that this formation is generally known as a carbonatic formation, Leach et al. (2010) considered Mehdiabad as a MVT-type deposit. The total thickness of the Taft Formation in the deposit area is about 400 meters (Figs. 5A and 23). However, Zn-Pb-Ba-(Cu-Ag) mineralization does not occur in the whole Taft Formation, but is located in 120 meters of the lower part of this formation (Figs. 5A, 6C, 7, 23 and 24). From these 120 meters, 80 meters consist of organic matter-rich shale, fine-grained black carbonate-rich siltstone, dolomite and the rest is carbonate (Figs. 5A, 23 and 24). Therefore, the main host rocks of mineralization in the Mehdiabad deposit are the clastic-carbonatic lithotypes: a feature similar to other SEDEX-type deposits (Figs. 23 and 24).

Most MVT Pb–Zn deposits are located in the regional thrust belts and along secondary reverse faults, and most researchers have considered that the Pb–Zn ore deposition was controlled by thrusting (Leach et al., 2005; 2010; Wilkinson, 2014). In fact, the ages of the carbonate hosted Pb–Zn mineralization stages in the MVT-type deposits are closely related to the timing of thrust faulting in the region of mineralization in the world. The MVT carbonate hosted Pb–Zn mineralization's are epigenetic (formed after the sedimentation of the host rock) (Leach et al., 2005; 2010; Wilkinson, 2014). Unlike most MVT-type deposits, Mehdiabad typically occurs within fine-grained marine sedimentary rocks (lower Unit of Taft Formation) (Figs. 23B and 24) overlying coarse-grained ones (Sangestan Formation) deposited in a continental rift basin, and are commonly spatially associated with the BH synsedimentary normal fault (Figs. 6A, 8B, 12 and 23A). The sharp contact between the Sangestan Formation and the Taft Formation near Mehdiabad (Fig. 4c) suggests the presence of a synsedimentary fault (BH), which may have acted as a pathway for the mineralizing fluids (Figs. 6A and 12). The hydrothermal alteration and the presence of mass flows observed in the proximal feeder zone at Mehdiabad (Figs. 12, 13, 20 and 23A) is reported in several SEDEX deposits (Tom, Jason, Vangorda, Rammelsberg, Century, Red Dog) and is generally interpreted to precede or overlap the mineralization event (Broadbent et al., 1998; Leach et al., 2005; Slack et al., 2004).

In the Mehdiabad deposit the ore minerals are fine-grained, and there are layered, banded and replacement textures (Figs. 16, 17 and 24). The chemistry of the ore minerals (e.g. higher Ag and Cu contents) reflects the high temperatures of the ore-forming fluid and is not typical of many MVT deposits. The feeder zone of the Mehdiabad deposit contains a copper mineralization (Figs. 13 and 24 (drill hole 7905)) that show a zonal pattern away from the main fault zone, with copper concentrating closest to the BH synsedimentary normal fault in the most altered rocks

(Figs. 12 and 24 (drill hole 7905)). Again this is unlike the numerous deposits typical of MVT districts.

Wilkinson (2014) defines sub-seafloor replacement deposits as deposits where the ores are hosted by carbonate and clastic rocks; they also have synsedimentary normal faults controlling the deposition of massive sulfides, formally known as sub-seafloor replacement SEDEX. These deposits (e.g., Anarraq) have many morphological and textural similarities to Irish-type ore bodies. In the sub-seafloor replacement SEDEX deposits, synsedimentary faults controlled the formation of seafloor depressions that may have trapped brines, acted as conduits for ore fluids, and generated debris flow breccias that can become mineralized (Figs. 6A, 12, 18, 19, 20, 23 and 24). Based on Wilkinson (2014), it is clear that the sub-seafloor replacement SEDEX-type deposits ore bodies are more akin to typical examples of the SEDEX family than to typical examples of the MVT clan. The principal similarity that the sub-seafloor replacement SEDEX-type deposits ores have with MVT deposits is that a part of the mineralization's can be carbonate-hosted but, as noted earlier, this is a poor classification criterion for sediment-hosted deposits given that classic shale-hosted SEDEX camps can include carbonate-hosted deposits, and many SEDEX deposits are hosted by carbonate-rich mudstones or siltstones (Wilkinson, 2014). Similar to SEDEX-type deposits, the Mehdiabad mineralization is hosted by organic matter-rich shale, fine-grained black carbonate-rich siltstone, and dolomite interlayered with a sandstone sequence (Figs. 9, 16, 17, 23A and 24). SEDEX deposits are generally tabular in shape and predominantly contain various amounts of Zn, Pb, Cu, Au, Ag, Ba, Fe and Mn oxides.

The geometry of the Mehdiabad deposit, shown in Figs. 6A and 12, and its mineralogical, structural and textural features suggest that the deposit is a SEDEX-style. However, typically, in these sub-seafloor replacement SEDEX-type deposits, there can be some evidence of exhalation

of the fluids onto, or below the seafloor, to form laminated and stratabound ores (see Wilkinson, 2014). There is evidence of laminated ores, derived from exhalative processes at the Mehdiabad deposit (Figs. 16, 17 and 24). Mehdiabad has characteristics similar to those of the Anarraaq sub-seafloor replacement SEDEX deposit, located in the Red Dog basin, Canada. These are: the exhalative textures deposited on the seafloor, the sub-seafloor replacement textures and the synsedimentary fault that controls the mineralization parameters (Kelley et al., 2004).

We have refined the sub-seafloor replacement SEDEX genetic type for Mehdiabad (Maghfouri et al., 2015; Maghfouri, 2017), at a level of detail useful to explain the prevailing textural and chemical characteristics of the deposit (Figs. 17–18). In fact, the characteristics defining Mehdiabad as a sub-seafloor replacement SEDEX-type deposit are: 1) the marine extensional basin setting; 2) the clastic-carbonatic marine sedimentary host rock sequence (Figs. 5A, 8A, 9, 23 and 24); 3) the bedding-parallel Zn-Pb-Ba-(Cu-Ag) sulfide lenses (Figs. 8, 12, 15, 16 and 17); 4) the synsedimentary faults that controlled the formation of seafloor depressions that may have trapped the metal-rich brines, acted as conduits for ore fluids, and generated debris flow breccias that can become mineralized (Figs. 6A, 12 and 20); 5) the well-developed lateral and vertical geochemical and mineralogical zoned patterns, extending for kilometers to tens of meters from the BH synsedimentary fault (Fig. 12); 6) similarly to other SEDEX-type deposits, the barite content in the Mehdiabad deposit is variable and occurs above the sulfide mineralization (Figs. 6A, 12 and 14A); 7) the sulfides display a variety of textures; the most characteristic are fine-grained layered and banded lithotypes, but coarser-grained brecciated, veined and replacement textures are also commonly observed (Figs. 13, 14, 16, 17, 18, 19 and 24). Based on the above characteristics, we consider Mehdiabad as a vent-proximal sub-seafloor

replacement clastic-carbonate hosted SEDEX-type deposit, and not a typical Mississippi Valley-type deposit (MVT) (Leach et al., 2005).

5-5- Model for the formation of the hypogene mineralization in the Mehdiabad deposit

The Mehdiabad deposit has a prominent feeder zone and a well-developed massive-replacement ore, forming what has been interpreted as a vent complex, similar to many well-known shale-hosted massive-replacement sulfide deposits. Thus, the Mehdiabad deposit can be classified as a vent-proximal sub-seafloor replacement clastic-carbonate hosted SEDEX-type deposit (cf. Lydon, 1996; Sangster 2002; Large et al. 2005; Goodfellow and Lydon 2007).

The formation of the Mehdiabad deposit was strongly affected by the structural setting of the basin (Fig. 25). The stratiform and stratabound orebodies of this deposit occur in the thickest section of organic matter-rich shale and dolomite units, immediately adjacent to the synsedimentary Black-Hill normal fault (Maghfouri. 2017) (Fig. 25). This fault is considered to be a favorable conduit for hydrothermal fluid upflow and played an important role for the formation of the deposit, similar to other deposits described by Large et al. (2002), Leach et al. (2005), and Rajabi et al. (2014). In the Mehdiabad deposit, presumed feeder zones developed next to the syn-sedimentary normal fault (Fig. 25). The metal zonation patterns, the presence of synsedimentary breccias and debris flows, the increase of hydrothermal alteration and of the temperature of the ore fluids towards the faults, suggest that this fault was active during ore deposition and provided major conduits for the metalliferous hydrothermal fluids.

In contrast to the laminated shale-sulfide textures found in other SEDEX-type deposits (Sangster 2002; Large et al. 2005; Leach et al. 2005; Goodfellow and Lydon 2007; Wilkinson, 2014), textural evidence suggests that the Mehdiabad deposit formed both by deposition on the

832 sea floor and by subsequent further replacement of preexisting sediments and ore minerals.
 833 During the deposition of organic matter-rich shale (the Taft Formation), the water column in the
 834 southern Yazd Basin was likely stratified with deep anoxic waters (Fig. 25A). The regional
 835 anoxic conditions for the basin are indicated by physical evidence, such as well-preserved
 836 laminated sedimentary texture, the presence of organic matter, and the grain size of pyrite
 837 framboids. Early venting of the exhaling hydrothermal fluids in the basin led to the precipitation
 838 of an early assemblage of sulfide minerals in the bedded ore facies as a reaction of the mixing
 839 with seawater (Fig. 25B). Because the ore-forming fluids at Mehdiabad were probably slightly
 840 less dense than seawater (Maghfouri, 2017), they behaved as a buoyant plume (Sato, 1972), at
 841 least initially near and above the fluid vent. Subsequently mixing and cooling occurred, so that
 842 the fluid and the metals precipitating from the vent settled on the seafloor after exhalation (Fig.
 843 25B). The buoyant plume behavior of hydrothermal fluids in the Mehdiabad deposit is consistent
 844 with observations in modern back-arc sediment-covered rifts, such as the Okinawa trough
 845 (Halbach et al. 1991) and with similar ancient massive sulfide deposits, including the Heath
 846 Steele Belt in the Bathurst Mining Camp District (Goodfellow et al. 2003), the Red Dog deposit
 847 (Moore et al., 1986), and the Paleozoic Selwyn Basin deposits in Canada (Goodfellow, 2007).
 848 Fluid mixing with seawater produced cooling and alkalization, resulting in the destabilization
 849 of Pb and Zn chloride complexes and in the precipitation of sulfides. A significant proportion of
 850 reduced sulfur at the Mehdiabad deposit had a biogenic origin, which is supported by: 1) the
 851 broad range of $\delta^{34}\text{S}$ sulfide values, and 2) the textural evidence (framboidal and colloform
 852 textures) for microbial biota associated with sulfide deposition. This early stage of mineralization
 853 is economically insignificant at Mehdiabad, because the hydrothermal fluids, being less dense
 854 than seawater, dispersed metals away from the venting site within the sediments. The fine-

grained nature of the early stage minerals also suggests a rapid deposition (Herzig & Hannington 1995, Kelley et al. 2004b).

Continued up-flow of high-temperature, metal-bearing, dense hydrothermal fluids into and along the bedded ore facies (ore stage 1) and dolomite shows that early sulfide minerals, barite, and dolomitic rocks were later replaced by massive sulfides (ore stage 2) (Fig. 25C). Most of the Mehdiabad sulfide mineralization was deposited during this main ore stage due to subsurface replacement processes in the host rocks (Fig. 25C), due to favorable porosity and permeability conditions. Such a sulfide replacement can be ascribed to either sub-seafloor replacement of carbonate and sulfide lamina and bands, without ascent of fluids onto the seafloor (Fig. 25C), or by sinking of ore fluids into the seafloor sediments from a brine pool in a local depression (Sangster, 2002). In this model, metal-bearing fluids move up along the syn-sedimentary normal fault. Notwithstanding limited leakage of fluid onto the seafloor, these fluids mixed with saline seawater in the unconsolidated sediments and then spread laterally into the host rocks (Fig. 25C). Sulfide minerals formed by replacement of specific sedimentary laminae and bands, where metal chloride complexes in the fluid reacted with H₂S generated by the thermochemical reduction of sulfates at a temperature higher than about 178-238 °C (Maghfouri, 2017).

The ores from the second stage of mineralization (massive-replacement) are overlain by bedded barite (Fig. 12). The vertical transition between the massive-replacement ore facies and the stratiform barite beds is gradual and is characterized by increase of the bedded barite mineralization. These relations shown that brecciation and replacement of dolomite by sulfide minerals predates the closure of the basin. Stratiform barite mineralization forms the top of ore horizon I and conformably covers the sulfide mineralization (Fig. 25D). The barite ore body, which was deposited in oxic seawater, is well exposed in the Mehdiabad area. The transition

from the sulfide to sulfate ore implies major changes in the sedimentary regime, from dominantly anoxic seawater to oxic seawater (Figs. 25C and 25D). This stage of mineralization is characterized by massive to laminated barite and by minor framboidal pyrite, sphalerite, and galena. Barite mineralization in the Mehdiabad deposit formed predominantly on the seafloor (Fig. 18D). SO_4^{2-} required for the formation of barite in this deposit originated from Early Cretaceous seawater sulfate (Maghfouri, 2017) and H_2S requisite for sulfide mineralization was provided by bacterial reduction. Ultimately, the limestone rocks of the Taft Formation covered barite mineralization in the Mehdiabad deposit and the formation of Zn-Pb-Ba mineralization terminated in the ore horizon I (Fig. 25E).

Subsequently the area where the Mehdiabad ore deposit is located experienced inversion tectonics during Tertiary, leading to a modification of the initially Lower Cretaceous geometry. This late stage was accompanied by supergene oxidation and alteration (Reichert et al., 2003; Maghfouri et al., 2017) and a partial remobilization of the metals, especially in the BHO and ER areas. In the CM area, the origin of the upper Ore Horizon (OHII) is not clearly understood yet, as this orebody is completely oxidized.

6- Conclusions

The Mehdiabad deposit was formed in the early stages of a long-lived starved sedimentary basin. In the Mehdiabad sequence >1300 m of sediment accumulated in this basin during Lower Cretaceous. The deposit is hosted within the Taft Formation. The lower unit of the Taft Formation is a rhythmically banded, possibly distal turbiditic calcarenite and shale unit called the Sangestan Formation. The Sangestan Formation is the stratigraphic footwall to mineralization at Mehdiabad. The lower part of the Taft Formation, consisting of organic matter-rich shale,

siltstone and dolomite, contains the bulk of the sulfide mineralization. Sediments of the Taft Formation were deposited in a dysoxic to anoxic water column favorable to sulfate-reduction. Bacterial reduction of seawater sulfates provided reduced sulfur for the Mehdiabad sulfides, although partially oxygenated bottom water conditions may have restricted the sulfate-reducing bacteria to the underlying sediments.

Mehdiabad is a well-preserved vent-proximal SEDEX deposit consisting of (1) a stockwork of sulfide veins with a hydrothermal alteration halo, discordantly underlying the main orebody; (2) a massive-replacement sulfide orebody containing pyrite, sphalerite, barite and galena with quartz and dolomite; (3) bedded ore that includes laminated sulfides with sedimentary structures; (4) barite ore with minor pyrite, sphalerite and galena (5) distal exhalites made up of banded chert with disseminated barite and minor pyrite. The Mehdiabad deposit has a prominent feeder zone (stockwork) overlain by a well-developed vent complex (massive ore) that laterally gives way to a bedded, stratiform ore body, similar as in the Mansourabad, Sullivan and Tom deposits (Goodfellow and Lydon, 2007; Maghfouri and Hosseinzadeh, 2018). Sulfide minerals were deposited from buoyant hydrothermal fluids vented into an anoxic basin that was affected by synsedimentary faulting and developed in a submarine environment. A thick accumulation of sulfide-bearing barite locally caps the sulfide deposit. The upwardly limiting factor for sulfide precipitation at Mehdiabad, therefore, may not have been a failure at the vent but a change to oxidizing conditions, which favored deposition of barite rather than sulfides.

The Mehdiabad deposit formed proximal to the source vent. Most of mineralization is a massive-replacement or breccia textured to indistinctly bedded; facies changes from one ore facies to another occur over short distances. Host-rock replacement textures are common, mainly

in massive ore facies. Mehdiabad deposit formed by a combination of sulfides and barite deposition on the sea floor, and sediment replacement within the sub-seafloor.

7- Acknowledgements

This paper is part of the senior author's PhD research study at University of Tabriz, Tabriz, Iran. The authors thank the University of Tabriz Research Grant Council and the Mehdiabad Mining Complex for funding fieldwork studies and providing access to exploration data and core logs. The paper benefited from detailed comments by Prof. Ebrahim Rastad. The authors thank the board of Mehdiabad Mining Complex (M. Hosseini and A. Safari) and Hajghasemi for allowing access to the deposit and their records. Particular thanks go to H. Pourfaraj for helping the first author during data collection.

References

- Aghanabati, A. (1998) Major sedimentary and structural units of Iran (map): Geosciences, v. 7, p. 29–30.
- Aghanabati, A. (2004) Geology of Iran. Geological Survey and Mineral Exploration of Iran, Tehran, 586 pp.

- 942 Agard, A., Omrani, J., Jolivet, L., Mouthereau, F. (2005) Convergence history across Zagros
943 (Iran): constraints from collisional and earlier [deformation](#): *Int. J. Earth Sci.* v. 94, p. 401–419.
- 944 Alavi, M. 1991, Tectonic map of the Middle East: Tehran, Geological Survey of Iran, scale
945 1:5,000,000.
- 946 Alavi, M. (1994) Tectonics of the Zagros orogenic belt of Iran: New data and interpretations:
947 *Tectonophysics*, v. 229, p. 211– 238.
- 948 Alfonso, P., Prol-Ledesma, R.M., Canet, C., Melgarejo, J.C., and Fallick, A.E (2005) Isotopic
949 evidence for biogenic precipitation as a principal mineralization process in coastal
950 gasohydrothermal vents, Punta Mita, Mexico: *Chemical Geology*, v. 224, p. 113–121.
- 951 Aistov, L., Melnikov, B., Krivyakin, B., Morozov, L., Kiristaev, V. (1984) Geology of the Khur
952 area (Central Iran). Explanatory Text of the Khur Quadrangle Map 1:250.000. – V/O
953 Technoexport, Report 20, pp. 1–130.
- 954 Anderson, I.K., Ashton, J.H., Boyce, A.J., Fallick, A.E., Russell, M.J. (1998) Ore depositional
955 processes in the Navan Zn+Pb deposit, Ireland. *ECONOMIC GEOLOGY*, v.93, p. 535–563.
- 956 Anger, G., Nielsen, H., Puchelt, H., Ricke, W. (1966) Sulfur isotopes in the Rammelsberg ore
957 deposit (Germany). *ECONOMIC GEOLOGY*, v. 61, p. 511–536.
- 958 Bagheri, S., Stampfli, G.M. (2008) The Anarak, Jandaq and Posht-e-Badam metamorphic
959 complexes in central Iran: new geological data, relationships and tectonic implications:
960 *Tectonophysics*, v. 451, p. 123–155.
- 961 Barrier, E., Vrielynck, B. (2008a) Map 6: [Cenomanian](#) (99.6–93.5 Ma). In: Barrier, E.,
962 Vrielynck, B. (Eds.), *Palaeotectonic Maps of the Middle East – Tectono- Sedimentary-*

- 963 Palinspastic Maps from the Late Norian to Pliocene. Commission for the Geological Map of
964 the World (CGMW/CCGM), Paris.
- 965 Barrier, E., Vrielynck, B. (2008b) Map 5: [Middle Aptian \(121.0–115.0 Ma\)](#). In: Barrier, E.,
966 Vrielynck, B. (Eds.), Palaeotectonic maps of the Middle East – tectonosedimentary–
967 palinspastic maps from the Late Norian to Pliocene. Commission for the Geological Map of
968 the World (CGMW/CCGM), Paris.
- 969 Berberian, M., King, G.C.P. (1981) Towards a palaeogeography and tectonic evolution of Iran:
970 Can. J. Earth Sci, v. 18, p. 210–265.
- 971 Berberian, F., Muir, I.D., Pankhurst, R.J., Berberian, M. (1982) Late Cretaceous and early
972 Miocene Andean-type plutonic activity in northern Makran and Central Iran: J. Geol. Soc.
973 London, v. 139, p. 605–614.
- 974 Berner, R.A, (1969) The synthesis of framboidal pyrite: ECONOMIC GEOLOGY, p. 64, p. 383
975 – 384.
- 976 Betts, P.G., Lister, G.S., O'dea, M.G., (1998) Asymmetric extension of the Middle Proterozoic
977 lithosphere, Mount Isa terrane, Queensland, Australia: Tectonophysics, v. 296, 293–316.
- 978 Betts, P.G., Lister, G.S., (2001) Comparison of the ‘strike-slip’ versus the ‘episodic rift-sag’
979 models for the origin of the Isa Superbasin: Aust. J. Earth Sci., v. 48, 265–280.
- 980 Boni, M., Gilg, H.A., Balassone, G., Schneider, J., Allen, C.R., Moore, F. (2007) Hypogene Zn
981 carbonate ores in the Angouran deposit, NW Iran. Mineralium Deposita: v. 42, p. 799-820.
- 982 Bottrell, S.R., Newton, R.J. (2006) Reconstruction of changes in global sulfur cycling from
983 marine sulfate isotopes: Earth-Science Reviews, v. 75, p. 59– 83.

- 984 Boviri, M., Rastad, E. (2016) Nature and origin of dolomitization associated with sulphide
985 mineralization: new insights from the Tappehsorkh Zn-Pb (-Ag-Ba) deposit, Irankuh Mining
986 District, Iran. *Geological Journal*. [http://dx.doi.org/ 10.1002/gj.2875](http://dx.doi.org/10.1002/gj.2875).
- 987 Boveiri, M., Rastad, E., Peter, J. (2017) A sub-seafloor hydrothermal syn-sedimentary to early
988 diagenetic origin for the Gushfil Zn-Pb-(Ag-Ba) deposit, south Esfahan, Iran: *Neues Jahrbuch*
989 *für Mineralogie-Abhandlungen, Journal of Mineralogy and Geochemistry*, v. 194 (1), p. 61-
990 90.
- 991 Brumsack, H.-J. (1986) The Inorganic Geochemistry of Cretaceous Black Shales (DSDP 41) in
992 Comparison to Modern Upwelling Sediments from the Gulf of California. In: Summerhayes,
993 C.P., Shackelton, N.J. (Eds.), *North Atlantic Palaeoceanography*, Geological Society, London,
994 Special Publication, v. 21, p. 447–462.
- 995 Bucur, I.I., Senowbari-Daryan, B., Majidifard, M.R. (2003) Neocomian microfossil association
996 from the Taft area near Yazd (Central Iran): *Facies*, v. 49, p. 217–222.
- 997 Bucur, I.I., Rashidi, K., Senowbari-Daryan, B. (2012) Early Cretaceous calcareous algae from
998 central Iran (Taft Formation, south of Alibad, near Yazd): *Facies*, v. 58, p. 605–636.
- 999 Buschendorf, F., Nielsen, H., Puchelt, H., Ricke, W. (1963) Schwefelisotopen-untersuchungen
1000 am Pyrit-Sphalerit-Baryt-Lager Meggen/Lenne (Deutschland) und an verschiedenen Devon-
1001 Evaporiten: *Geochim Cosmochim Acta*, v. 27, p. 501–523.
- 1002 Canet, C., Alfonso, P., Melgarejo, J.C., and Fallick, A.E. (2005) Stable isotope geochemistry of
1003 the Carboniferous Zn-Pb-Cu sediment-hosted sulfide deposits, northeastern Spain:
1004 *International Geology Review*, v. 47, p. 1298–1315.

- 2521
2522
2523 1005 Cifelli, F., Mattei, M., Rashid, H., Ghalamghash, J. (2013) Right-lateral transpressional tectonics
2524
2525 along the boundary between Lut and Tabas Blocks (Central Iran): *Geophys. J. Int.*, v. 193, p.
2526 1006
2527 1153–1165.
2528 1007
2529
2530 1008 Claypool, G.E., Holser, W.T., Kaplan, I.R., Sakai, H., Zak, I. (1980) The age curves for sulfur
2531
2532 and oxygen isotopes in marine sulfate and their mutual interpretation: *Chemical Geology*, v.
2533 1009
2534 28, p. 199–260.
2535 1010
2536
2537 1011 Cross, M.M., and Bottrell, S.H. (2000) Reconciling experimentally observed sulphur isotope
2538
2539 fractionation during thermochemical sulphate reduction (TSR) with field data: A ‘steadystate’
2540 1012
2541 model of isotopic behaviour: *Goldschmidt, Journal of Conference Abstracts*, v. 5, p. 325.
2542 1013
2543
2544 1014 Daliran, F., Pride, K., Walther, J., Berner, Z.A., Bakker, R.J. (2013) The Angouran Zn (Pb)
2545
2546 deposit, NW Iran: evidence for a two stage, hypogene zinc sulfide zinc carbonates
2547 1015
2548 mineralization: *Ore Geology Reviews*, v. 53, p. 373-402.
2549 1016
2550
2551 1017 Davoudzadeh, M. (1997) Iran. In: Moores, E.M., Fairbridge, R.M. (Eds.), *Encyclopedia of*
2552
2553 *European and Asian Regional Geology*, *Encyclopedia of Earth Sciences Series*. Chapman and
2554 1018
2555 Hall, London, p. 384–405.
2556 1019
2557
2558 1020 Davoudzadeh, M., Soffel, H., Schmidt, K. (1981) On the rotation of the Central-East Iran
2559
2560 microplate: *Neues Jb. Geol. Paläontol. Monat* 5, p. 180–192.
2561 1021
2562
2563 1022 Dercourt, J., Zonenshain, L.P., Ricou, L.-E., Kazmin, V.G., Le Pichon, X., Knipper, A.L.,
2564
2565 Grandjacquet, C., Sbertshikov, I.M., Geyssant, J., Lepvrier, C., Pechersky, D.H., Boulin, J.,
2566 1023
2567 Sibuet, J.-C., Savostin, L.A., Sorokhtin, O., Westphal, M., Bazhenov, M.L., Lauer, J.P., Bijou-
2568 1024
2569
2570
2571
2572
2573
2574
2575
2576

- 2577
2578
2579 1025 Duval, B. (1986) Geological evolution of the Tethys belt from the Atlantic to the Pamir since
2580
2581 1026 the Lias: Tectonophysics, v. 123, p. 241–315.
2582
2583
2584 1027 Ehya, F., Lotfi, M., Rasa, I. (2010) Emarat carbonate-hosted Zn–Pb deposit, Markazi Province,
2585
2586 1028 Iran: A geological, mineralogical and isotopic (S, Pb) study: Journal of Asian Earth Sciences,
2587
2588 1029 v. 37, p. 186–194.
2589
2590
2591 1030 Elliott, H. (2015) Pb-Zn mineralisation within the Limerick Basin (SW Ireland): A role for
2592
2593 1031 volcanism? PhD thesis, National Oceanography Centre, Southampton School of Ocean and
2594
2595 1032 [Earth Science. Pp. 197.](#)
2596
2597
2598 1033 Esmaeily, D., Bouchez, J.L., Siqueira, R. (2007) Magnetic fabrics and microstructures of the
2599
2600 1034 Jurassic Shah-Kuh granite pluton (Lut Block, Eastern Iran) and geodynamic inference:
2601
2602 1035 Tectonophysics, v. 439, p. 149–170.
2603
2604
2605 1036 Fallick, A.E., Ashton, J.H., Boyce, A.J., Ellam, R.J., Russell, M.J. (2001) Bacteria were
2606
2607 1037 responsible for the magnitude of the world-class hydrothermal base metal-sulfide orebody at
2608
2609 1038 Navan, Ireland: ECONOMIC GEOLOGY, v. 96, p. 885–890.
2610
2611
2612 1039 Gadd, M.G., Matthews, L.D., Peter, J.M. (2016a) Non-hydrothermal origin of apatite in SEDEX
2613
2614 1040 mineralization and host rocks of the Howard’s Pass district, Yukon, Canada: American
2615
2616 1041 Mineralogist, v. 101, p. 1061–1071.
2617
2618
2619 1042 Gadd, M.G., Matthews, L.D., Peter, J.M., Paradis, S., Jonsson, R.I. (2016b) The world-class
2620
2621 1043 Howard’s Pass SEDEX Zn-Pb district, Selwyn Basin, Yukon. Part II: the roles of
2622
2623 1044 [thermochemical and bacterial sulfate reduction in metal fixation. Miner Deposita, Volume 52,](#)
2624
2625 [Issue 3, pp 405–419.](#)
2626
2627
2628
2629
2630
2631
2632

- 1046 Ghasemi, M. (2006) Formation Mechanism of the Mehdi Abad Zn-Pb Deposit and its
1047 Comparison with Other Near Lead and Zinc Deposits (Unpublished M.Sc. Thesis): Research
1048 Institute of Earth Sciences, Geological Survey of Iran, p. 238.
- 1049 Ghazban, F., McNutt, R.H., Schwarcz, H.P. (1994) Genesis of sediment-hosted Zn-Pb-Ba
1050 deposits in the Iran Kouh district, Esfahan area, west-Central Iran: ECONOMIC GEOLOGY,
1051 v. 89, p. 1262–1278.
- 1052 Gilg, H.A., Boni, M., Balassone, G., Allen, C.R., Banks, D., Moore, F. (2006) Marble hosted
1053 sulphide ores in the Angouran Zn- (Pb-Ag) deposit, NW Iran: interaction of sedimentary
1054 brines with a metamorphic core complex: Mineralium Deposita, v. 41, p. 1-16.
- 1055 Goodfellow, W.D., Jonasson, I.R. (1987) Environment of formation of the Howards Pass (XY)
1056 Zn-Pb deposits, Selwyn Basin, Yukon, in: Morin, J.A., (eds.), Mineral deposits of Northern
1057 Cordillera: Canadian Institute of Mining and Metallurgy Spec, v. 37, p. 19–50.
- 1058 [Goodfellow, W.D., Mccutcheon, S.R., Peter, J.M. \(2003\)](#) Massive sulfide deposits of the
1059 Bathurst Mining Camp, New Brunswick, and Northern Maine: introduction and summary of
1060 findings: [Econ Geology Monograph](#), 11:1–1.
- 1061 Goodfellow, W.D. (2004) Geology, genesis and exploration of SEDEX deposits, with emphasis
1062 on the Selwyn basin, Canada, in Deb, M., Goodfellow, W.D., (eds.), Sediment-hosted lead-
1063 zinc sulphide deposits: Attributes and models of some major deposits of India, Australia and
1064 Canada. Delhi, India, Narosa Publishing House, p. 24–99.
- 1065 Goodfellow, W.D., and Lydon, J.W. (2007) Sedimentary exhalative (Sedex) deposits, in
1066 Goodfellow, W.D., eds., Mineral deposits of Canada: A synthesis of major deposit types,

- 2689
2690
2691 1067 district metallogeny, the evolution of geological provinces, and exploration methods:
2692
2693 1068 Geological Association of Canada, Mineral Deposits Division, Special Publication 5, p. 163–
2694
2695 183.
2696 1069
2697
2698 1070 Halbach, P.E., Marten, A., Schwanold, G. (1991) Present-day Kuroko type ore formation: results
2699
2700 from the central Okinawa trough. In: Pagel M, Leroy JL (eds) Source, transport and
2701 1071
2702 deposition of metals: Balkema, Rotterdam, p. 623–626.
2703 1072
2704
2705 1073 Haghipour, A., Pelissier, G. (1970) Geological Quadrangle Map of Iran, H 8, Ardekan
2706
2707 1:250,000. Geological Survey of Iran, Tehran.
2708 1074
2709
2710 1075 Hajmolla Ali, A., Majidifard, M.R. (2000) Geological Map of Yazd, Scale 1:100,000. Geological
2711
2712 Survey of Iran.
2713 1076
2714
2715 1077 Herzig, P.M., Hannington, M.D. (1995) Polymetallic massive sulfides at the modern seafloor, a
2716
2717 review: Ore Geology Review, v. 10, p. 95–115.
2718 1078
2719
2720 1079 Hoffman, D.L., Algeo, T.J., Maynard, J.B., Joachimski, M.M., Hower, J.C., Jaminski, J. (1998)
2721
2722 Regional stratigraphic variation in bottom water anoxia in offshore core shales of Upper
2723 1080
2724 Pennsylvanian cyclothems from Eastern Midcontinent Shelf (Kansas), U.S.A. in: Schieber, J.,
2725 1081
2726 Zimmerle, W., Sethi, P., (eds.), Shales and mudstones I.E Schweizerbart'sche
2727 1082
2728 Verlagsbuchhandlung, Stuttgart, p. 243–269.
2729 1083
2730
2731 1084 Huston, D.L., Power, M., Gemmell, J.B., Large, R.R. (1995) Design, calibration and geological
2732
2733 application of the first operational Australian laser ablation sulphur isotope microprobe: Aust.
2734 1085
2735 J. Earth Sci, v. 42, p. 549–555.
2736 1086
2737
2738
2739
2740
2741
2742
2743
2744

- 1087 Ireland, T. (2001) Lateral Variability in the HYC Zn-Pb-Ag deposit, Northern Territory,
1088 Australia; implications for ore genesis. Unpubl Honours thesis, University of Tasmania,
1089 Australia.pp. 189.
- 1090 Ireland, T., Bull, S.W., Large, R.R. (2004a) Mass flow sedimentology within the HYC Zn-Pb-Ag
1091 deposit, Northern Territory, Australia: Evidence for syn-sedimentary ore genesis: Mineralium
1092 Deposita, v. 39, p. 143–158.
- 1093 Ireland, T., Large, R.R., McGoldrick, P., Blake, M. (2004b) Spatial distribution patterns of sulfur
1094 isotopes, nodular carbonate, and ore textures in the McArthur River (HYC) Zn-Pb-Ag deposit,
1095 Northern Territory, Australia: ECONOMIC GEOLOGY, v. 99, p. 1687-1709.
- 1096 Jones, B., Manning, D.A.C (1994) Comparison of geochemical indices used for the interpretation
1097 of paleoredox conditions in ancient mudstone: Chemical Geology, v. 111, p. 111–129.
- 1098 Johnson, C.A., Kelley, K.D., Leach, D.L (2004) Sulfur and oxygen isotopes in barite deposits of
1099 the western Brooks Range, Alaska, and implications for the origin of the Red Dog massive
1100 sulfide deposits: ECONOMIC GEOLOGY, v. 99, p. 1435–1448.
- 1101 Kelley, K.D., Dumoulin, J.A., Jennings, S (2004a) The Anarraq Zn-Pb- Ag and barite deposit,
1102 northern Alaska: Evidence for replacement of carbonate by barite and sulfides: ECONOMIC
1103 GEOLOGY, v. 99, p. 1577–1591.
- 1104 Kelley, K.D., Leach, D.L., Johnson, C.A., Clark, J.L., Fayek, M., Slack, J.F., Anderson, V.M,
1105 Ayuso, R.A, Ridley, W.I. (2004b) Textural, compositional, and sulfur isotope variations of
1106 sulfide minerals in the Red Dog Zn-Pb-Ag deposits, Brooks Range, Alaska: Implications for
1107 ore formation: ECONOMIC GEOLOGY, v. 99, p. 1509–1532.

- 1108 Large, R.R., Bull, S.W., Cooke, D.R., McGoldrick, P.J. (1996) Review of genetic models at
 1109 HYC: Constraints from new sedimentology, alteration halo studies and fluid chemical
 1110 modelling: Townsville, James Cook University of North Queensland: ECONOMIC
 1111 GEOLOGY Research Unit Contribution, v. 55, p. 72–74.
- 1112 Large, R.R., Bull, S.W., Yang, J., Cooke, D.R., Garven, G., McGoldrick, P.J., Selley, D. (2002)
 1113 Controls on the formation of giant stratiform sediment-hosted Zn-Pb-Ag deposits with
 1114 particular reference to the north Australian Proterozoic. University of Tasmania, Centre for
 1115 Special Ore Deposit and Exploration (CODES) Studies Publication 4, p. 107–149
- 1116 Large, R.R., Bull, S.W., McGoldrick, P.J., Walters, S., Derrick, G.M., Carr, G.R. (2005)
 1117 Stratiform and strata-bound Zn-Pb-Ag deposits in Proterozoic sedimentary basins, northern
 1118 Australia. ECONOMIC GEOLOGY 100th Anniversary, p. 931–963
- 1119 Leach, D.L., Sangster, D.F., Kelley, K.D., Large, R.R., Garven, G., Allen, C.R., Gutzmer, J.,
 1120 Walters, S. (2005) Sediment-hosted lead-zinc deposits: a global perspective. ECONOMIC
 1121 GEOLOGY 100th Anniversary, p. 561–607
- 1122 Leach, D.L., Bradley, D.C., Huston, D., Pisarevsky, S.A., Taylor, R.D., Gardoll, S.J (2010)
 1123 Sediment-hosted lead-zinc deposits in Earth history: ECONOMIC GEOLOGY, v. 105, p.
 1124 593–625.
- 1125 Lee, M.J., and Wilkinson, J.J. (2002) Cementation, hydrothermal alteration, and Zn-Pb
 1126 mineralization of carbonate breccias in the Irish Midlands: Textural evidence from the
 1127 Cooleen zone, near Silvermines, County Tipperary: ECONOMIC GEOLOGY, v. 97, p.
 1128 653–662.

- 1129 Liaghat, S., Moore, F., Jami, M. (2000) The Kouh Sourmeh mineralization, a carbonate-hosted
1130 Zn-Pb deposit in the Simply Folded Belt of the Zagros Mountains, SW Iran: Mineralium
1131 Deposita, v. 35, p. 72-78.
- 1132 Lindenberg, H.G., Görler, K., Ibbeken, H. (1983) Stratigraphy, structure and orogenic
1133 evolution of the Sabzevar Zone in the area of Oryan Khorasan, NE Iran. Geological Survey of
1134 Iran Report, v. 51, p. 120–143.
- 1135 [Logan, R.G. \(1979\) The Geology and Mineralogical Zoning of the H.Y.C. Ag-Pb-Zn Deposit,](#)
1136 [McArthur River, Northern Territory, Australia. Unpublished Thesis. Australian National](#)
1137 [University, pp. 341.](#)
- 1138 Lydon, J.W. (1996) Sedimentary exhalative sulphides (SEDEX), in Eckstrand, O.R., Sinclair,
1139 W.D., Thorpe, R.I., (eds.), Geology of Canadian Mineral Deposit Types, Geology of Canada,
1140 No. 8, Geological Survey of Canada (also Geological Society of America, The Geology of
1141 North America P-1), p. 130-152.
- 1142 Lydon, W.J. (2004) Geology of the Beit-Purcell Basin and the Sullivan deposit, in Deb, M.
1143 Goodfellow, W.D., (eds.), Sediment-hosted lead-zinc sulphide deposits: Attributes and models
1144 of some major deposits of India, Australia and Canada. Narosa Publishing House, Delhi,
1145 India, p. 100-148.
- 1146 Maghfouri, S., Hosseinzadeh, M.R., Rajabi, A., Azimzadeh, A.M., Choulet, F. (2015) Geology
1147 and Origin of Mineralization in the Mehdiabad Zn-Pb-Ba (Cu) Deposit, Yazd Block, Central
1148 Iran, 13th SGA biennial meeting, Nancy-France.pp. 3.

- 1149 Maghfouri, S., Hosseinzadeh, M.R., Rajabi, A., Azimzadeh, A.M. (2016a) Facies analysis and
1150 stratigraphy position of carbonate-clastic hosted Zn-Pb-Ba mineralization horizons in the
1151 Early Cretaceous sedimentary sequence, Southern Yazd Basin: Scientific Quarterly Journal,
1152 Geosciences, v. 26, p. 102-115.
- 1153 Maghfouri, S., Hosseinzadeh, M.R., Rajabi, A., Azimzadeh, A.M (2016b) Darreh-zanjir Deposit,
1154 a Typical Carbonate Hosted Zn-Pb Deposit (MVT) in Early Cretaceous Sedimentary
1155 Sequence, Southern Yazd Basin, Scientific Quarterly Journal, Geosciences, v. 26, No: 103.
- 1156 Maghfouri, S. (2017) Geology, Geochemistry, Ore Controlling Parameters and Genesis of Early
1157 Cretaceous Carbonate-clastic Hosted Zn-Pb Deposits in Southern Yazd Basin, with Emphasis
1158 on Mehdiabad Deposit (Unpublished Ph.D. Thesis). Tabriz University, Iran, p. 475.
- 1159 Maghfouri, S., and Hosseinzadeh, M.R. (2018) The early Cretaceous Mansourabad shale
1160 carbonate hosted Zn–Pb (-Ag) deposit, central Iran: An example of vent-proximal sub-
1161 seafloor replacement SEDEX mineralization. Ore Geology Reviews, v. 95, p. 20–39.
- 1162 Magnall, J.M., Stern, R.A., Gleeson, S.A., Paradis, S. (2015) Widespread euxinic conditions are
1163 not a prerequisite for sediment-hosted base metal (Pb-Zn-Ba) mineralization at MacMillan
1164 Pass, Yukon. In S. Paradis, Ed., Targeted Geoscience Initiative 4: Sediment-hosted Zn-Pb
1165 deposits: Processes and implications for exploration, p. 43–57. Geological Survey of Canada,
1166 Open File 7838.
- 1167 Magnall, M.J., Gleeson, S.A., Blamey, N.J.F., Paradis, S., Luo, Y. (2016) The thermal and
1168 chemical evolution of hydrothermal vent fluids in shale hosted massive sulphide (SHMS)
1169 systems from the MacMillan Pass district (Yukon, Canada): Geochimica et Cosmochimica
1170 Acta, v. 193, p. 251–273.

- 2969
2970
2971 1171 Majidifard, M.R. (1996) Stratigraphy, fossils and environment of Early Cretaceous rocks from
2972
2973 the northern hills of Shirkuh: Geological Survey of Iran, Earth Science 20, 2e31 (in Farsi).
2974 1172
2975
2976 1173 Majidifard, M.R., Vaziri, S.H. (2000) Geological Map of Bahadoran, Scale 1:100,000.
2977
2978 1174 Geological Survey of Iran.
2979
2980
2981 1175 Mattei, M., Cifelli, F., Muttoni, G., Zanchi, A., Berra, F., Mossavvari, F., Eshraghi, S.A. (2012)
2982
2983 Neogene block rotation in central Iran: evidence from paleomagnetic data: Geol. Soc, Am.
2984 1176
2985 Bull. v. 124, p. 943–956.
2986 1177
2987
2988 1178 Mohseni, M. (2011) Geology and mineralization of the Mehdi Abad Zn-Pb Deposit
2989
2990 (Unpublished M.Sc. Thesis). Dameghan University, Iran, 190 p.
2991 1179
2992
2993 1180 Momenzadeh, M. (1976) Stratabound Lead-Zinc Ores in the Early Cretaceous and Jurassic
2994
2995 Sediments in the Malayer-Esfahan District (West Central Iran), Lithology, Metal Content,
2996 1181
2997 Zonation and Genesis (Unpublished Ph.D. Thesis). University of Heidelberg, Heidelberg, 300
2998 1182
2999 p.
3000 1183
3001
3002 1184 Momenzadeh, M., Shafighi, S., Rastad, E., and Amstutz, G.S. (1979) The Ahangaran lead-silver
3003
3004 deposit, Se-Malayer, west Central Iran: Mineralium Deposita, v. 14, p. 323–341.
3005 1185
3006
3007 1186 Moore, D.E., Young, L.E., Modene, J.S., Plahuta, J.T. (1986) Geologic setting and genesis of the
3008
3009 Red Dog zinc-lead-silver deposit, Western Brooks Range, Alaska: ECONOMIC GEOLOGY,
3010 1187
3011 v. 81, p. 1696–1727.
3012 1188
3013
3014 1189 Muttoni, G., Mattei, M., Balini, M., Zanchi, A., Gaetani, M., Berra, F (2009a) The drift history
3015
3016 of Iran from the Ordovician to the Triassic. In: Brunet, M.-F., Wilmsen, M., Granath, J.W.
3017 1190
3018
3019
3020
3021
3022
3023
3024

- 1191 (Eds.), South Caspian to Central Iran Basins, v. 312. Geological Society London, Special
1192 Publications, p. 7–29.
- 1193 Muttoni, G., Gaetani, M., Kent, D.V., Sciunnach, D., Angiolini, L., Berra, F., Garzanti, E.,
1194 Mattei, M., Zanchi, A (2009b) Opening of the Neo-Tethys ocean and the Pangea B to Pangea
1195 A transformation during the Permian: GeoArabia, v. 14, p. 17–48.
- 1196 Nabavi, M (1972a) Lower Cretaceous deposits in the Taft-Yazd and Khur area. Geological
1197 Survey of Iran, Report 106, p. 1–127.
- 1198 Nabavi, M.H (1972b) Geological quadrangle map of Iran, H 9, Yazd 1:250,000. Geological
1199 Survey of Iran, Tehran.
- 1200 Nath, B. N., Bau, M., Ramlin Geswara Rao, B. & Rao, Ch.M (1997) Trace and rare earth
1201 elemental variation in Arabian Sea sediments through a transect across the oxygen minimum
1202 zone. Geochim: Cosmochim. Acta, v. 61, p. 2375 – 2388.
- 1203 Nosratian, J (1991) Geological Map of Mehdi Abad. Based on the Topographical Map Mehdi
1204 Abad-Yazd, 1:2000.
- 1205 Ohmoto, H., Kaiser, C.J., Geer, A (1990) Systematics of sulphur isotopes in recent marine
1206 sediments and ancient sediment-hosted base metal deposits, in: Herbert, H.K., Ho, S.E., (eds.),
1207 Stable isotopes and fluid processes in mineralization, v. 23, Perth, University of Western
1208 Australia Publishing, p. 70–120.
- 1209 Peter, J.M., Scott, S.D (1999) Windy Craggy, northwestern British Columbia: the world's largest
1210 Besshi deposit. Rev. ECONOMIC GEOLOGY. v. 8, p. 261–295.

- 1211 Pfaff, K., Hildebrandt, H., Leach, D., Jacob, D., Markl, G (2010) Formation of the Wiesloch
Mississippi Valley-type Zn-Pb-Ag deposit in the extensional setting of the Upper
Rhinegraben, SW Germany: *Miner Deposita*, v. 45, p. 647–666
- 1214 Philip, J., Floquet, M (2000) Late Cenomanian. In: Dercourt, J., Gaetani, M., Vrielynck, B.,
Barrier, E., Biju-Duval, B., Brunet, M.F., Cadet, J.P., Crasquin, S., Sandulescu, M. (Eds.),
Atlas Peri-Tethys Palaeogeographical Maps. CCGM/CGMW, Paris, p. 129–136.
- 1217 Pourfaraj, H., Maghfouri, S., Mohajjel, M., Rastad, E (2016) Kinematic change in syn-
sedimentary normal fault in Mehdi-Abad Mine of Yazd, 33rd National Geosciences
Symposium, Trhran.
- 1220 Prol-Ledesma, R.M., Canet, C., Villanueva-Estrada, R.E., and Ortega-Osorio, A (2010)
Morphology of pyrite in particulate matter from shallow submarine hydrothermal vents:
American Mineralogist, v. 95, p. 1500–1507.
- 1223 Quinby-Hunt, M.S., Wilde, P (1994) Thermodynamic zonation in the black shale facies based on
iron–manganese–vanadium content: *Chemical. Geology*, v. 113, p. 297–317.
- 1225 Rajabi, A., Rastad, E., Canet, C (2012) Metallogeny of Cretaceous carbonate hosted Zn–Pb
deposits of Iran: geotectonic setting and data integration for future mineral exploration,
International Geology Review, v. 54:14, p. 1649-1672.
- 1228 Rajabi A, Rastad E, Canet C (2013) Metallogeny of Permian–Triassic carbonate-hosted Zn–Pb
and F deposits of Iran: a review for future mineral exploration: *Aust J Earth Sci*, v. 60, p. 197–
216

- 1231 Rajabi, A., Rastad, E., Canet, C., Alfonso, P (2014) Basin evolution and Stratigraphic Correlation
1232 of Sedimentary-Exhalative Zn-Pb deposits, Zarigan-Chahmir Basin: Ore Geology Reviews, v.
1233 64, p. 328–353.
- 1234 Rasa, I (1987) Geologisch-petrographische untersuchungen in der Blei-lagerstatte Nakhlak,
1235 Zentraliran [Ph.D. thesis]: Heidelberg, Heidelberg University.
- 1236 Rastad, E (1981) Geological, Mineralogical, and Facies Investigations on the Early Cretaceous
1237 Stratabound Zn-Pb-(-Ba-Cu) Deposits of the Iran Kouh Mountain Range, Esfahan, West
1238 Central Iran (Unpublished Ph.D. Thesis). University of Heidelberg, Heidelberg, 334 p.
- 1239 Reed, C., Wallace, M (2004) Zn-Pb mineralisation in the Silvermines district, Ireland: a product
1240 of burial diagenesis: Mineralium Deposita, v. 39, p. 87–102.
- 1241 Reichert, J., Borg, G. & Rashidi, B. (2003) Mineralogy of calamine ore from the Mehdi Abad
1242 zinc-lead deposit, Central Iran. In: Eliopoulos, D.G. et al. (eds.) Mineral Exploration and
1243 Sustainable Development. vol. 1, 97-100, Millpress, Rotterdam.
- 1244 Reichert, J (2007) A Metallogenetic Model for Carbonatehosted Non-sulfide Zinc Deposits
1245 Based on Observations of Mehdi Abad and Iran Kouh, Central and Southwestern Iran
1246 (Unpublished Ph.D. Thesis). University of Martin Luther, Shillong, 129 p.
- 1247 Rossetti, F., Nasrabady, M., Vignaroli, G., Theye, T., Gerdes, A., Razavi, M., Moin Vaziri, H
1248 (2010) Early Cretaceous migmatitic mafic granulites from the Sabzevar range (NE Iran):
1249 implications for the closure of the Mesozoic peri-Tethyan oceans in central Iran: Terra Nova,
1250 v. 22, p. 26–34.

- 3193
3194
3195 1251 Sáez, R., Moreno, C., González, F., Almodóvar, G.R (2011) Black shales and massive sulfide
3196
3197 deposits: causal or casual relationships? Insights from Rammelsberg, Tharsis, and Draa Sfar:
3198 1252
3199 Mineralium Deposita, v. 46, no. 5–6, p. 585–614.
3200 1253
3201
3202 1254 Sangster, D.F (2002) The role of dense brines in the formation of vent-distal sedimentary
3203
3204 exhalative (SEDEX) lead-zinc deposits: field and laboratory evidence: Mineralium Deposita,
3205 1255
3206 v. 37, p. 149-157.
3207 1256
3208
3209 1257 Sato, T (1972) Behaviours of ore-forming solutions in seawater: Mining Geology, v. 22, p. 31–
3210
3211 42.
3212 1258
3213
3214 1259 Schlagintweit, F., Bucur, I.I., Rashidi, K., Hanifzadeh, R., Wilmsen, M (2013a) Torremiroella
3215
3216 hispanica Brun and Canérot, 1979 (benthic foraminifera) from the Lower Cretaceous of
3217 1260
3218 Central Iran and its palaeobiogeographic significance: Cretaceous. Research, v. 46, p. 272–279.
3219 1261
3220
3221 1262 Schlagintweit, F., Bucur, I.I., Rashidi, K., Saberzadeh, B (2013b) Praeorbitolina claveli n. sp.
3222
3223 (benthic foraminifera) from the Lower Aptian (Bedoulian) of Central Iran: Carnets de
3224 1263
3225 Géologie/Notebooks on Geology, v. 4, p. 255–272.
3226 1264
3227
3228 1265 Seyed-Emami, K., Bozorgnia, F., Eftekhari-Nezhad, J (1972) Der erste sichere Nachweis von
3229
3230 Valanginien im nordöstlichen Zentraliran (Sabzewar-Gebiet): Neues Jb. Geol. Paläontol.
3231 1266
3232 Monat, v.1, p. 52–67.
3233 1267
3234
3235 1268 Shanks, W.C., Woodruff, L.G., Jilson, G.A., Jennings, D.S., Modene, J.S., Ryan, B.D (1987)
3236
3237 Sulfur and lead isotope studies of stratiform Zn-Pb-Ag deposits, Anvil Range, Yukon: basinal
3238 1269
3239 brine exhalation and anoxic bottom-water mixing: ECONOMIC GEOLOGY, v. 82, p. 600–
3240 1270
3241 634.
3242 1271
3243
3244
3245
3246
3247
3248

3249
3250
3251 1272 Soffel, H., Davoudzadeh, M., Rolf, C., Schmidt, S (1996) New palaeomagnetic data from
3252
3253 Central Iran and a Triassic palaeo-reconstruction: *Geol. Rundsch*, v. 85, p. 293–302.
3254 1273
3255
3256 1274 Southam, G., Saunders, J.A (2005) The geomicrobiology of ore deposits: *ECONOMIC*
3257
3258 *GEOLOGY*, v. 100, p. 1067–1084.
3259 1275
3260
3261 1276 Stampfli, G.M., Borel, G.D (2002) A plate tectonic model for the Paleozoic and Mesozoic
3262
3263 constrained by dynamic plate boundaries and restored synthetic oceanic isochrones: *Earth*
3264 1277
3265 *Planet. Sci*, v. 196, p. 17–33.
3266 1278
3267
3268 1279 Takin, M (1972) Iranian geology and continental drift in the Middle East: *Nature*, v. 235, p. 147–
3269
3270 150.
3271 1280
3272
3273 1281 Taraz, H., Aghanabati, A (1983) Geological Quadrangle Map of Iran, G 9, Abadeh 1:250,000.
3274
3275 Geological Survey of Iran, Tehran.
3276 1282
3277
3278 1283 Tirrul, R., Bell, I.R., Griffis, R.J., Camp, V.E (1983) The Sistan suture zone of eastern Iran.
3279
3280 *Geol. Soc. Am. Bull*, v. 94, p. 134–150.
3281 1284
3282
3283 1285 Velasco, F., Herrero, J.M., Gil, P.P., Alvarez, L., Yusta, I (1994) Mississippi Valley-type,
3284
3285 SEDEX, and iron deposits in Lower Cretaceous rocks of the Basque-Cantabrian Basin,
3286 1286
3287 northern Spain. In: Fontbote, L., Boni, M., (eds.), *Sediment-hosted Zn-Pb Ores*. Society for
3288 1287
3289 *Geology Applied to Ore Deposits*, Springer-Verlag, Berlin-Heidelberg, v. 10, p. 246–270.
3290 1288
3291
3292 1289 Wilkinson JJ (2003) On diagenesis, dolomitisation and mineralisation in the Irish Zn-Pb orefield:
3293
3294 *Mineralium Deposita*, v. 38, p. 968–983.
3295 1290
3296
3297
3298
3299
3300
3301
3302
3303
3304

- 1291 Wilkinson JJ, Eyre, S.L., Boyce, A.J. (2005) Ore-forming processes in Irish-type carbonate-
- 1292 hosted Zn-Pb deposits: Evidence from mineralogy, chemistry and isotopic composition of
- 1293 sulfides at the Lisheen Mine: ECONOMIC GEOLOGY, v. 100, p. 63–86.
- 1294 Wilkinson, J.J. (2014) Sediment-Hosted Zinc–Lead Mineralization: Processes and Perspectives,
- 1295 Treatise on Geochemistry 2nd Edition.
- 1296 Wilmsen, M., Fürsich, F.T., Seyed-Emami, K. (2003) Revised lithostratigraphy of the Middle
- 1297 and Upper Jurassic Magu Group of the northern Tabas Block, Eastcentral Iran: Newsl.
- 1298 Stratigr, v. 39, p. 143–156.
- 1299 Wilmsen, M., Wiese, F., Seyed-Emami, K., Fürsich, F.T. (2005) First record and significance of
- 1300 Cretaceous (Turonian) ammonites from the Shotori Mountains, east-central Iran: Cretaceous
- 1301 Research, v. 26, p. 181–195.
- 1302 Wilmsen, M., Fürsich, F.T., Seyed-Emami, K., Majidifard, M.R., Taheri, J (2009) The
- 1303 Cimmerian orogeny in northern Iran: tectono-stratigraphic evidence from the foreland: Terra
- 1304 Nova, v. 21, p. 211–218.
- 1305 Wilmsen, M., Fürsich, F.T., Majidifard, J (2013) The Shah Kuh Formation, a latest Barremian –
- 1306 early Aptian carbonate platform of Central Iran (Khur area, Yazd Block). Cretaceous
- 1307 Research, v. 39, p. 183–194.
- 1308 Wilmsen, M., Fürsich, F. T., Majidifard, M. R (2014) An overview of the Cretaceous
- 1309 stratigraphy and facies development of the Yazd Block, western Central Iran, Journal of Asian
- 1310 Earth Sciences.pp. 18.

3361
3362
3363 1311 Whitehead, R.E. (1973) Environment of stratiform sulphide deposition; variation in Mn:Fe ratio
3364
3365 1312 in host rocks at Heath Steele Mine, New Brunswick, Canada: Mineral. Deposita, v. 8, p. 148–
3366
3367 160.
3368 1313
3369
3370 1314 Whittaker, S.G., Kyser, T.K. (1993) Variations in the neodymium and strontium isotopic
3371
3372 composition and REE content of molluscan shells from the Cretaceous Western Interior
3373 1315
3374 Seaway: Geochim. Cosmochim. Acta, v. 57, p. 4003 – 4014.
3375 1316
3376
3377 1317 Wingal, P.B., Twitchett, R.J. (1996) Oceanic anoxia and the end-Permian mass extinction:
3378
3379 Science, v. 272, p. 1155 –1158.
3380 1318
3381
3382 1319 Yaghubpur, A., Mehrabi, B. (1997) Kushk zinc-lead deposit: a typical black-shale-hosted deposit
3383
3384 in Yazd State, Iran: Journal of Sciences, Islamic Republic of Iran, v. 8, no. 2, p. 117–125.
3385 1320
3386
3387 1321 Yarmohammadi, A., Rastad, E., Rajabi, A. (2016) Geochemistry, fluid inclusion study and
3388
3389 genesis of the sediment-hosted Zn-Pb ($\pm$ Ag $\pm$ Cu) deposits of the Tiran basin, NW of
3390 1322
3391 Esfahan, Iran: N. Jb. Miner. Abh. (J. Min. Geochem.), v. 193/2, p. 183–203.
3392 1323
3393
3394
3395 1324
3396
3397
3398 1325
3399
3400
3401 1326
3402
3403
3404 1327
3405
3406 1328
3407
3408
3409 1329
3410
3411
3412
3413
3414
3415
3416

3417
3418
3419
3420
3421
3422
3423
3424
3425
3426
3427
3428
3429
3430
3431
3432
3433
3434
3435
3436
3437
3438
3439
3440
3441
3442
3443
3444
3445
3446
3447
3448
3449
3450
3451
3452
3453
3454
3455
3456
3457
3458
3459
3460
3461
3462
3463
3464
3465
3466
3467
3468
3469
3470
3471
3472

Table captions

Table 1. Major (%) and trace elements (ppm) contents of the Taft Formation shales hosting the mineralization of the Mehdiabad deposit. (BH: Black-Hill ore, CVOB: Central Valley Orebody, ER: East Ridge ore).

Table 2. Values of $\delta^{34}\text{S}_{\text{SDT}}$ in sulfide minerals and barite from the Mehdiabad deposit

Table 3. Assay analysis of ore minerals in the different drill holes of Mehdiabad deposits (Location of samples shown in the figure 12).

Figure captions

Fig.1. Distribution map of sediment hosted Zn-Pb deposits in the main tectonic units of Iran; Al: Alborz zone; CIGS: Central Iranian geological and structural gradual zone; E: East Iran ranges; K: Kopeh-Dagh; KR: Kermanshah Radiolarites subzone; KT: Khazar-Talesh-Ziveh structural zone; L: Lut Block; M: Makran zone; Oph: ophiolite belts; PB: Posht-e-Badam Block; SSZ: Sanandaj-Sirjan zone; T: Tabas Block; TM: tertiary magmatic rocks; UDMA: Urumieh-Dokhtar magmatic arc; Y: Yazd Block; Z: Zabol area; Za: Zagros ranges. Tectonic and structural map of Iran modified after Aghanabati (1998; 2004) and Alavi (1991).

Fig. 2. Simplified geological map of the YAMB in the CIEM, showing the location of the different types of ore deposits. Blue rectangle shows the location of the Southern Yazd Basin in the Fig. 3. (NDBF: Naein-Dehshir-Baft fault; ANF: Anar fault; PBF: Posht-e-Badam fault; CHF:

Chapedoni fault; BF: Biabanak fault; GKDF: Great Kavir Drounaeh fault) (modified after Haghypour and Pelissier, 1970; Nabavi, 1972b; Taraz and Aghanabati, 1983).

Fig. 3. Simplified geological map of the southern Yazd Basin, showing the location of Zn-Pb deposits within this basin (modified after Majidifard and Vaziri, 2000; Hajmolla Ali and Majidifard, 2000). Small rectangle shows the location of study area in the Fig. 4.

Fig. 4. Geological map of the Mehdiabad deposit showing the location of the barite deposit in the east of Black-Hill Fault and of the oxide Zn-Pb orebody in the northeast of the map. CVF: Central Valley fault; NTF: North Thrust fault; FF: Fourzandeh fault; BHO: Black-Hill ore; CVOB: Central Valley orebody; ER: East Ridge; CM: Calamine Mine (A-A', B-B' and C-C' sections are shown in Figs. 8 and 12) (modified after Nosrati, 1991).

Fig. 5. (A) Schematic columnar section of the Lower Cretaceous sedimentary sequence hosting the Mehdiabad deposit. The lower unit of the Taft Formation and the middle unit of the Abkuh Formation host the Zn-Pb-Ba mineralization. (B): General view of the Shir-Kuh granite, unconformably covered by the Sangestan Formation (view to the northeast). (C): View of the silicified silty limestone (distal facies) marking the limit between the Sangestan and Taft Formations (view to the northwest). (D): Hand specimen photograph of silty limestone with interbedded organic matter-rich shale host rock in the Mehdiabad deposit.

Fig. 6. (A): General view of the two ore horizons (OH I and OH II) with different ore facies, OH I is located within the Taft Formation, whereas OH II corresponding to the CM orebody that occurs within the Abkuh Formation. Mineralization in the Mehdiabad deposit occurs to the east of the Black-Hill Fault. Numbers show the location of cross-sections in the Fig. 7. (1 to 5 numbers of columnar sections and A-A', B-B', C-C' and D-D' sections are shown in Figs. 7, 8

and 12) (view to the north-northwest). (B): The location of the bedded ore facies (ER) and distal facies that are embedded with Sangestan and Taft Formations (view to the northwest).

Fig. 7. Lithostratigraphic correlation diagram of the Early Cretaceous sedimentary sequence in the Mehdiabad deposit from west (1) to east (5) showing the position of the ore horizons in the stratigraphic units. Numbers show the stratigraphical location of cross-sections in the Fig. 6A.

Fig. 8. Cross-sections and plan view of the ore sequence and extension of mineralization in the Mehdiabad deposit. (A): North-south cross section of the ore sequence with extension of sulfide and non-sulfide ore mineralization's in the CVOB. (B): Schematic cross section of the OH I in the Mehdiabad deposit, which include the BH orebody and ER mineralization. (C): Southeast-northwest section through the Mehdiabad deposit illustrating the geology, ore horizon, and structure of the area. The location of these sections is shown in Figs. 4 and 6A.

Fig. 9. Hand specimen (A, B, C, D and E) and microscopic (F) photographs of ore-bearing host rocks in the ore horizon I. (A): Drill core of organic matter-rich (Om) shale-siltstone (lower Unite) from the Taft Formation overlying the massive ore facies. (B): Hand specimen photograph of organic matter-rich (Om) shale with organic matter nodule (lower Unit of Taft Formation). (C): Alternation of silty limestone laminae (light-gray) and sphalerite (Sph)+galena (Gn)+pyrite (Py) sulfides (black gray) laminas. (D): Hand specimen photograph of shale. Gray laminas are shale and siltstone rich in pyrite (Py), whereas black laminae are rich in organic matter. (E): Typical laminated siltstone and silty sandstone hosting the ore horizon I. Note the precedence of normal micro-fractures, locally inversed in the lower part of specimen. (F): Silty shale laminae composed of feldspar, quartz, sericite, and clay minerals alternating with black organic matter (Om) laminae.

Fig. 10. Histograms showing the distribution of Fe and Mn contents of unaltered shale samples from the Taft Formation. These diagrams are used to discriminate the paleoenvironment and fields are taken from Quinby-Hunt and Wilde (1994). (A): the Mn content indicates the anoxic paleoconditions during the sedimentation of shale. (B): the Fe content is related to both anoxic sulfate reducing and nitrate-reducing conditions.

Fig. 11. Bivariate V/(V+Ni) vs. V/Cr plot indicating geochemical redox conditions at the time of the black shale deposition (modified after Sáez et al. 2011). Most samples plot in the anoxic conditions field. Field boundaries are from Jones and Manning (1994) and Hoffman et al. (1998).

Fig. 12. W-NE cross-section of the ore sequence and extension of ore facies in the Mehdiabad deposit (Maghfouri, 2017). The mineralization is located to the east of the Black-Hill syn-sedimentary fault. Location of this section is shown in Fig. 6A.

Fig. 13. Outcrop, hand specimen and microscopic photographs of main generations of veins and veinlets in the feeder zone of the Mehdiabad deposit.

Fig. 14. Outcrop (A), hand specimen (B, C, D, E, I and J) and microscopic (F, G and H) photographs of the massive-replacement ore facies in the Mehdiabad deposit. (A): General view of the relations between the massive-replacement ore facies and the barite ore. The massive oxide ore facies is capped by the barite ore mineralization. (B): Massive pyrite and sphalerite, showing pyrite (Py) replaced by sphalerite (Sph). (C): Barite (Ba) replaced by galena (Gn). (D): Massive sulfide ore facies consisting of sphalerite (Sph) and barite (Ba) minerals. (E): Dolomite host rock brecciated and replaced by galena (Gn) and sphalerite (Sph). (F): Early pyrite (Py1) replaced by secondary sphalerite (Sph2) and galena (Gn2). (G): Massive-replacement ore facies composed of galena (Gn2), sphalerite (Sph2), barite (Ba) and chalcopyrite (Cpy1). (H):

Replacement of early colloform pyrite (Py1) by late sphalerite (Sph2) and galena (Gn2). (I and J): Hand specimen photographs of massive-replacement sphalerite (Sph), galena (Gn) and pyrite (Py) sulfide mineralization in the lower unit of the Taft Formation.

Fig. 15. Geology map of the ER area showing the location of bedded ore and the distal facies at the boundary between the Taft and the Sangestan Formations.

Fig. 16. Outcrop (A and B), hand specimen (C, D, E and F) and microscopic (G) photographs of the bedded ore facies and distal facies in the Mehdiabad deposit in the lower Unit of the Taft Formation. (A and B): General view of the alternation of limy siltstone (light) and oxide Zn-Pb (black) layers in the ER orebody. (C): Photograph of typical pyritic (Py) laminae in the organic matter (Om) rich shale of the lower unit. The organic matter occurs as laminae and/or lenses within the sulfide laminae. (D): Inter-ore breccia resting on laminated sulfide-bearing shales. (Sphalerite (Sph), galena (Gn) and pyrite (Py)) laminae associated with black organic matter (Om) rich shale of lower Unit in the Taft Formation. (E): Alternation of shale (black-gray) and sulfides (light gray) laminae, including galena (Gn) and sphalerite (Sph) in the lower unit of the Taft Formation. (F): Laminated limy siltstone showing the three main types of laminae; organic matter-rich laminae (Type 1), siltstone laminae (Type 2), and pyritic rich laminae (Type 3). (G): Laminated sulfides including galena (Gn) and sphalerite (Sph1) from the bedded ore facies. (H): Photograph of alternation of chert laminae and Fe-oxide laminae in the distal facies of the Mehdiabad deposit. (I): Hand specimen photograph of chert laminae and Fe-oxide lamina in the distal facies. (J): Reflected light image of disseminated pyrite (Py) in the distal facies of the Mehdiabad deposit.

Fig. 17. Photographs (A, B, C, D and E) and photomicrographs (E, F, G and H) of sulfide-sediment textures in Mehdiabad deposit from ore horizon I in the lower unit of the Taft Formation. (A): Photograph of typical pyritic (Py) laminae associated with black organic matter (Om) rich shale of the lower unit. (B): Alternation of organic-matter (Om) laminae (black-gray) and siltstone host rock layers (light gray). (C): Alternation of shale laminae (black gray) and sulfides (sphalerite (Sph), galena (Gn) and pyrite (Py)) lamianas. (D): Sphalerite (Sph), galena (Gn) and pyrite (Py) sulfide layers assocaited with the lower unit black shale host rock. The sulfides occur as laminae and/or lenses within sulfide layer. (E): Siltstone laminae alternating with carbonate laminae. (F): Reflected light image of early colloform pyrite (Py1) crosscut by late sphalerite (Sph2) and galena (Gn2). (G): Early framboidal pyrite from the organic matter-rich shale. (H): Photomicrograph of laminated galena (Gn1) within the bedded ore facies in the lower unit of the Taft Formation.

Fig. 18. Photographs of sulfide-matrix breccia and sulfide-sediment textures in the Mehdiabad deposit from ore horizon I in the lower unit of the Taft Formation. (A and B): Sulfide-matrix breccia and Rounded-subrounded clasts of pyrite (Py) and galena (Gn) in a turbidite bed. In the Fig.A matrix is sphalerite (Sph), galena (Gn), pyrite (Py) and host rock and fragments of breccia is silty limestone and siltstone of the lower unit. In the Fig. B fragments and matrix both are pyrite (Py), galena (Gn) and sphalerite (Sph). (C): Photograph of sulfide-matrix breccia in the lower unit of the Taft Formation. (D): Load-cast texture, with a dolomite clast set on the thin-bedded shale. (E and F): Coarse-grained clasts, penetratating the underlying siltstone and shale bands. (G): Poorly sorted sulfide breccia layer with clasts of pyrite (Py), sphalerite (Sph) and galena (Gn). These layers are parallel to bedding.

Fig. 19. Textures of stage 2 (sub-seafloor replacement) sulfide mineralization in the ore horizon I. (A): Stage 1 (laminated sulfide and barite) crosscut and brecciated by stage 2 sulfide mineralization. Clasts of stage 1 relict within stage 2 sulfide matrix. (B and C): Stage 2 sulfides preferentially replacing matrix of dolomite breccia. (D): Replacement of dolomite (Dol) with stage 2 sphalerite (Sph) minerals is shown.

Fig. 20. Localization of the different mass flows textures, within the Mehdiabad deposit. Grain size of mass flows decreasing away from the BH syn-sedimentary fault. Sph: sphalerite; Gn: galena; Py: pyrite; Ba: barite; Om: organic matter. (See Fig. 12 for cross-section legend)

Fig. 21. (A): Histogram of the $\delta^{34}\text{S}$ values in the different ore facies of the Mehdiabad deposit . (B): Histogram of the $\delta^{34}\text{S}$ values for sulfide minerals and ore-associated barite from the Mehdiabad deposit. (C): Comparison between the $\delta^{34}\text{S}$ values of Mehdiabad sulfides and sulfates with other Zn-Pb deposits. References: Anvil District (Dy, Grum an Faro deposits), Canada (Shanks et al. 1987); Howards Pass, Canada (Goodfellow and Jonasson 1987); HYC, Australia (Ireland, 2001); Meggen, Germany (Buschendorf et al. 1963); Mt Isa, Australia (Large et al. 2005); Rammelsberg, Germany (Anger et al. 1966); Reocin and Troya, Spain (Velasco et al. 1994); Sullivan, Canada (Lydon, 2004); Tom and Jason, Canada (Goodfellow, 2004); Century, Australia (Wilkinson, 2014); George Fisher, Australia (Wilkinson, 2014); Navan and Silvermine, Ireland (Wilkinson, 2014); Red Dog, Canada (Wilkinson, 2014; Kelley et al., 2004a,b). Sulfur isotope composition of seawater sulfates is from Bottrell and Newton (2006).

Fig. 22. Paragenetic sequence of mineralization stages in the Mehdiabad deposit.

Fig. 23. (A): The six drill holes within the Sangestan, Taft and Abkuh Formations showing the sites of samples in the figure 24 and the boundaries of mineralization, BH syndimentary fault,

and different ore facies. (B): General view of the Sangestan Formation, conformably covered by Lower Unit of Taft Formation in the Farahabad area (view to the west). Lower Unit of Taft Formation is host rocks of main mineralization in the Mehdiabad deposit.

Fig. 24. Core photographs of Lower Unit of Taft Formation showing relation between ore minerals and rock forming minerals (location of samples shown in the figure 23A).

Fig. 25. Schematic diagram showing the inferred depositional environment of the Mehdiabad deposit. (A): Deposition of organic matter-rich shale (the Taft Formation) in the anoxic waters column. (B): Early venting of the exhaling hydrothermal fluid on the seafloor would have led to the precipitation of an early assemblage of the fine grain size minerals (stage 1) in the bedded ore facies. The temperature and salinity of hydrothermal fluids of this stage is 141 °C and 8.7% NaCl (Maghfouri, 2017). (C): Continued up-flow of high-temperature, metal-bearing, dense hydrothermal fluid into and along the bedded ore facies (ore stage I) and dolomite at a sub-sea floor level, show that the ore stage I minerals and dolomitic rocks were replaced by massive sulfide (ore stage II), normally towards their base. The temperature and salinity of hydrothermal fluids of this stage is 178-238 °C and 10.1% NaCl (Maghfouri, 2017). (D): This stage of mineralization consists of the precipitation of barite and minor sulfide minerals at and on the sea floor. Barite mineralization occurs from an oxic seawater column. (E): The upper units of the Taft Formation (limestone rocks) received the deposition of barite mineralization and finally the formation of Zn-Pb-Ba mineralization ended in the ore horizon I.

Highlights

► The Mehdiabad Zn-Pb-Ba (Cu - Ag) deposit in the southern Yazd basin, central Iran, is the largest sediment-hosted Zn-Pb deposit in Iran. ► Mineralization is mainly hosted in silty dolomite and shale. ► Sulfur isotopes, together with sedimentological, textural, mineralogical, and geochemical evidences, suggest that this deposit should be classified as a vent-proximal sub-seafloor replacement SEDEX ore deposit.

Massive ore facies

Massive, replacement, brecciated, colloform, vein-veinlet

Chalcopyrite, galena, sphalerite, pyrite, barite, dolomite, calcite, minor quartz

Bedded ore facies

Laminated, disseminated, framboidal

Sphalerite, galena, pyrite, Fe-oxide, minor barite, dolomite, calcite, Mn-minerals

**Vent-proximal sub-seafloor replacement clastic-carbonate hosted SEDEX-type
mineralization in the Mehdiabad world-class Zn-Pb-Ba-(Cu-Ag) deposit, Southern Yazd
Basin, Iran**

Sajjad Maghfouri^{a*} (Corresponding Author)

^a*Department of Geology, Faculty of Basic Sciences, Tarbiat Modares University, Tehran, Iran*
E-Mail: s.maghfouri@modares.ac.ir

Mohammad Reza Hosseinzadeh^b

^b*Department of Earth Sciences, Faculty of Natural Sciences, University of Tabriz, Tabriz, Iran*
E-Mail: mr-hosseinzadeh@tabrizu.ac.ir

Flavien Choulet^c

^c*Chrono-Environnement, Université de Franche-Comté/CNRS, 25030 Besançon cedex, France*
E-Mail: flavien.choulet@univ-fcomte.fr

Pura Alfonso^d

^d*Departament d'Enginyeria Minera i Recursos Minerals, Universitat Politècnica de
Catalunya, Catalunya, Spain*
E-Mail: pura@emrn.upc.edu

Amir M. Azim Zadeh ^e

^e *Department of Geology, Faculty of Science, University of Zanjan, Zanjan, Iran*
E-Mail: amir.azimzadeh@znu.ac.ir

Abdorrahman Rajabi ^f

^f *School of Geology, College of Science, University of Tehran, Tehran, Iran*
E-Mail: abdorrahman.rajabi@gmail.com

Abstract

The Mehdiabad Zn-Pb-Ba (Cu-Ag) deposit in the southern Yazd Basin, central Iran, is the largest sediment-hosted Zn-Pb deposit in Iran. This deposit is hosted by organic carbon matter-rich shale, fine-grained black siltstone, and dolomite interlayered with sandstone of the Taft Formation. Sedimentological and geochemical studies of the Taft Formation have shown that these organic carbon matter-rich shales formed during a period of basin deepening and under anoxic conditions. Based on the orebody structure, mineralogy, and ore fabrics, we recognize five different ore facies types in the Mehdiabad deposit: (1) a stockwork/feeder zone, consisting of a discordant sulfide mineralization, forming a stockwork of sulfide-bearing dolomite veins, cutting the sedimentary rocks of the footwall; (2) massive-replacement ore, including pervasive replacement carbonate by pyrite, chalcopyrite, galena, and sphalerite with minor barite; (3) a bedded ore, with laminated to disseminated pyrite, sphalerite, and galena; (4) barite ore, which contains accessory sulfide minerals and rare calcite at the top of the deposit, and (5) a distal facies, with minor disseminated and laminated pyrite, banded cherts, and disseminated barite. Two stages of base metal sulfide replacement have separate origins. Fine-grained sulfide bands (stage I), intricately interlayered with organic carbon-rich beds and thin turbidite beds, exhibit lamina and bedding textures, supporting a syn-sedimentary (onto the sea floor) origin. Coarse-grained base metal sulfides (stage II), occurring within breccias and veins to veinlets are considered to have formed by replacement during post sedimentation sub-seafloor fluid flow. $\delta^{34}\text{S}$ values of pyrite, sphalerite chalcopyrite and galena range from -22 to +4.6.8‰. The highest $\delta^{34}\text{S}$ values correspond to the feeder zone (+4.6 and -10.5‰), whereas massive-replacement (+3.7 to - 13.7‰) and bedded (-22 to -17‰) ore facies display depleted compositions. The overall range of $\delta^{34}\text{S}$ values is remarkably higher than typical magmatic values, suggesting that

sulfides formed from the reduction of seawater sulfate by bacteriogenic sulfate reduction in a closed or semi-closed system in the bedded ore, whereas thermochemical sulfate reduction likely played an important role in the feeder zone. The formation of the Mehdiabad deposit follows the evolution of the sedimentary basin. Abrupt lateral changes in facies and thickness, along with the existence of synsedimentary breccias and debris flows within the ore sequence, suggest the proximity to synsedimentary faults and tectonic activity contemporaneous with the sedimentation in the Lower Cretaceous, favorable for the formation of the ore deposit. The deposit formed from a combination of hydrothermal and syn-sedimentary processes. Sulfur isotopes, together with sedimentological, textural, mineralogical, and geochemical evidences, suggest that this deposit should be classified as a vent-proximal sub-seafloor replacement SEDEX ore deposit.

Keywords: Lower Cretaceous; Mehdiabad Zn-Pb-Ba (Cu - Ag) deposit; SEDEX-type, Southern Yazd Basin; Iran

1- Introduction

Sediment-hosted Zn-Pb ore deposits are major sources of base metals such as Zn-Pb-Ag ± Cu-Ba and are the primary sources of Zn and Pb (Large et al., 2005). These deposits are diversely hosted by carbonate or siliciclastic rocks, without significant genetic contributions of magmatic activity (Leach et al., 2005). The deposits display a broad range of relationships to enclosing host rocks, including stratiform, strata-bound, and discordant ores (Leach et al., 2005). About 600 sediment-hosted Zn-Pb deposits and occurrences are presently known in Iran (Fig. 1). Only a few deposits of these have been explored and/or exploited although <70 deposits are currently being mined (Rajabi et al., 2012; 2013). Despite numerous scientific debates, these deposits are considered to form a metallogenetic range from exhalative sedimentary (SEDEX) to

Mississippi Valley-types (MVTs) deposits (Rajabi et al., 2102; 2013). The most important Iranian sediment-hosted Zn-Pb ore deposits include the world-class Mehdiabad Zn-Pb-Ba (Cu-Ag) deposit (Maghfouri et al., 2015), the deposits world-class Irankuh Zn-Pb ($\pm$ Ag) district (Rastad, 1981; Ghazban et al., 1994; Boveiri and Rastad, 2016; Boveiri et al., 2017), the world-class Angouran Zn-Pb deposit (Gilg et al., 2006; Boni et al., 2007; Daliran et al., 2013), the Ahangaran Pb-Ag ($\pm$ Zn) deposit (Momenzadeh, 1976; Momenzadeh et al., 1979), the Koushk Zn-Pb vent-proximal SEDEX-type deposit (Yaghubpur and Mehrabi, 1997; Rajabi et al., 2012), the Emarat Zn-Pb MVT-type deposit (Ehya et al., 2010), the Tiran Zn-Pb ($\pm$ Ag $\pm$ Cu $\pm$ Ba) Sedex-type deposit (Yarmohammadi et al., 2016), the Nakhlak Pb-Zn MVT-type deposit (Rasa, 1987), and the Kuh-e-Sormeh Zn-Pb MVT-type deposit (Liaghat et al., 2000). These Zn-Pb deposits belong to the Yazd-Anarak metallogenic belt (YAMB), the Malayer-Esfahan metallogenic belt (MEMB), and the Central Alborz metallogenic belt (CAMB) (Rajabi et al., 2012; 2103) (Fig. 1).

The Yazd-Anarak Metallogenic Belt (YAMB) is located in the western section of the Central East Iranian Microcontinent (CEIM) (Fig. 1). The YAMB hosts abundant mineral deposits, particularly those of the Cu (Fe) porphyry-skarn (Darreh Zarashk and Hasanabad deposits (Oligocene–Miocene)), W-Cu-Ni-Au-As vein-types (Chah Palang (post-Jurassic), Talmesi and Meskani deposits (post-Eocene)), Zn-Pb MVT-types (Nakhlak, Darreh-Zanjir, Chahake Ardakani and Howze Safid deposits (post-late Cretaceous)), and Zn-Pb SEDEX types (Mehdiabad, Farahabad, Mansourabad (lower Cretaceous) and Haft-Har (lower Cambrian) deposits), being one of the main metallogenic provinces in Iran (Fig. 2) (Maghfouri et al., 2015; 2016a; 2016b; Maghfouri, 2017).

95 The southern Yazd Basin in the YAMB, which includes three lower Cretaceous sediment-hosted
96 Zn-Pb $\pm$ Ba deposits, including the Mehdiabad deposit and the Farahabad and Mansourabad
97 deposits (Figs. 2 and 3). Sedimentation during this geological time span was controlled by basin
98 subsidence and by the reactivation of half-graben faults. In the southern Yazd Basin,
99 sedimentation has occurred asymmetrically throughout the basin. The lower part of the lower
100 Cretaceous sequence is characterized by major lateral facies changes, observable between the
101 Mehdiabad and the Farahabad areas, and by the wedge-shaped geometry of the sedimentary
102 units; these stratigraphic features are indicative of rift related half-grabens (e.g., Betts and Lister,
103 2001; Betts et al., 1998). The Mehdiabad deposit, located 115 km southeast of Yazd city, can be
104 subdivided in four ore zones (Fig. 4): 1) the Black-Hill ore (BH), the Central Valley Orebody
105 (CVOB), the East Ridge ore (ER) and the Calamine Mine (CM). Throughout this paper, the term
106 “Mehdiabad deposits” refers collectively to the BH, CVOB, ER and CM ore bodies. The
107 Mehdiabad deposit is the largest accumulation of carbonate-hosted zinc in the world (Leach et
108 al., 2005), with a total resource of 394 Mt, grading 4.2% Zn, 1.6% Pb and 36 g/t Ag. In addition,
109 the barite content of this deposit is unusually large (>40 Mt) and covers an area of at least 1.5
110 km². Early studies at Mehdiabad (Ghasemi, 2006) described the initial geologic framework of
111 the deposit. This work showed that the Mehdiabad deposit bears geologic similarities with other
112 sediment-hosted deposits of the southern Yazd Basin. Ghasemi (2006) also proposed a
113 “Mehdiabad-type”, of deposits, to distinguish the YAMB deposits from other classical sediment-
114 hosted Zn-Pb deposits (SEDEX, MVT, Irish and massive sulfides). Mohseni (2011) further
115 studied the barite mineralization, located in the western part of the deposit, and classified it as an
116 Irish-type deposit. Recently petrological and geochemical studies of the deposit were carried out
117 in the frame of a Ph.D. research project between 2012 and 2017 and Maghfouri et al. (2015) and

Maghfouri (2017) presented geologic descriptions of the deposit and a preliminary genetic model. Based on the occurrence of exhalative hydrothermal distinctive characteristics such as laminated sulfide textures, lateral metal zonation, diagenetic replacement textures and the presence of a normal syn-sedimentary fault (Black-Hill Fault), a SEDEX model was proposed for the Mehdiabad deposit. Maghfouri and Hosseinzadeh (2018) studied the Mansourabad deposit, which is located in the southern Yazd Basin (Fig.3), and classified it as a vent-proximal sub-seafloor replacement SEDEX deposit model. They further proposed that the formation of SEDEX deposits in this basin is related to the evolution of an extensional Cretaceous environment that affected the Central Iranian microcontinent. This paper aims at providing advanced evidence from regional sedimentology, geochemistry, and sulfide-sediment relationships to support the synsedimentary origin for the Mehdiabad deposit.

2- Geological context and main features of the Mehdiabad deposit

2-1- Geology of the Yazd Block

The Mehdiabad deposit belongs to the Central-East Iranian Microcontinent (CEIM; Takin, 1972), which forms the central part of the present-day Iran Plate (Fig. 1). The CEIM consists, from east to west, of three structural units, i.e., the Lut, the Tabas, and the Yazd Blocks (Fig. 1) (Berberian and King, 1981; Berberian et al., 1982, and Davoudzadeh, 1997; Alavi, 1994). The stratigraphy of the Yazd Block is globally similar to the other parts of Central Iran; it contains sequences of Neoproterozoic to Middle Triassic sedimentary rocks, covered by extensive coal-bearing Upper Triassic to Mid-Jurassic sandstones, siltstones and shales (Shemshak Formation) (Fig. 2). The thick Jurassic sequence is well exposed in the Tabas Block and here it has been studied in details (see Nabavi, 1972a,b; Aistov et al., 1984; Wilmsen et al., 2003, 2009a;

Wilmsen et al., 2005; Wilmsen et al., 2014). During the Cretaceous, the CEIM was separated from the southern Eurasian margin by narrow oceanic basins (Sistan Ocean in the east, Sabzevar Ocean in the north, Nain–Baft Ocean in the west; Lindenberg et al., 1983; Tirrul et al., 1983; Dercourt et al., 1986; Philip and Floquet, 2000; Barrier and Vrielynck, 2008a,b). These oceanic basins opened during the Early Cretaceous (‘Neocomian’) (e.g., find of Valanginian tintinnids and ammonites from the Sabzevar Zone; Seyed-Emami et al., 1972) and their development may be connected to an inferred post-Triassic tectonic rotation of the CEIM around a vertical axis of about 135° with respect to Eurasia (e.g., Davoudzadeh et al., 1981; Soffel et al., 1996). Rotation most probably took place predominantly in post-Jurassic times (Esmaeily et al., 2007; Bagheri and Stampfli, 2008; Wilmsen et al., 2009a), although the timing and the amount of rotation are still debated (e.g., Muttoni et al., 2009a,b). Cifelli et al. (2013) demonstrated that the movements at the block-bounding fault between the Tabas and Lut Blocks of the CEIM changed from extensional during the Jurassic to a right-lateral transpression between the Early Cretaceous and the Paleocene. Mattei et al. (2012) have also documented evidence of significant Early Cretaceous and Neogene counterclockwise rotation of the CEIM. The various small oceans around the CEIM closed at the end of the Cretaceous and during the Paleocene, due to the closure of the Neotethys followed by the collision of the Arabian Plate (e.g., Tirrul et al., 1983; Dercourt et al., 1986; Stampfli and Borel, 2002; Agard et al., 2005; Rosetti et al., 2010).

2-2- Local geological setting of the Mehdiabad deposit

In the southern Yazd Basin, a thick so-called Lower Cretaceous Sedimentary Sequence (LCSS) is exposed, resting unconformably on the Jurassic Shir-Kuh granite and on metamorphic rocks of the Shemshak Group (Fig. 3). In the following chapter, the Lower Cretaceous

Formations of the Yazd area, identified by Nabavi (1972a,b; see also Majidifard, 1996) will be briefly described from base to top (Figs. 4 and 5).

2-2-1- The Sangestan Formation

The Early Cretaceous sequence in the southern Yazd Basin and at the Mehdiabad deposit started with the terrigenous sediments of the Sangestan Formation (Fig. 4), which unconformably overlies the metasediments of the Shemshak Group and the Shir Kuh granite (Figs. 4 and 5). The lower part of the Formation is exclusively siliciclastic (reddish conglomerate, sandstone and siltstone) reflecting semi-arid continental depositional conditions and upwardly-decreasing relief energy (Wilmsen et al., 2014) (Figs. 4 and 5A). Towards the top, a marine influence is indicated by evaporites and carbonates, in part containing restricted marine biota (calcareous algae, agglutinating foraminifera) (Wilmsen et al., 2014). The carbonates from the Upper Sangestan Formation are probably of Hauterivian age (Bucur et al., 2003). The Sangestan Formation is mostly equivalent to the Barremian, with a diachronic top due to the Hauterivian transgression (Wilmsen et al., 2014).

2-2-2- The Taft Formation

The Sangestan Formation is conformably overlain by organic carbon matter-rich shale, silty shale, sandy limestone, dolomite and siltstone of the Barremian to probably Aptian Taft Formation (Figs. 4 and 5) (Nabavi, 1972a), consisting of thick-bedded to massive rudists, algae, and orbitolinid-bearing shallow-water facies (e.g., Bucur et al., 2012; Schlagintweit et al., 2013a). The thickness of the Taft Formation reaches up to 400 m (section at Forouzandeh; Fig. 5C) and this series is part of the Lower Cretaceous carbonate platform system of the Yazd Block (Wilmsen et al., 2013a). On the basis of benthic foraminifera, Schlagintweit et al. (2013a,b)

proposed an Early Aptian (Bedoulian) age of the lower part of the Taft Formation. However, the base of the Taft Formation may well range locally into the Barremian (see Bucur et al., 2012). The Taft Formation includes two units (Fig. 5A). The lower unit (70–85 m thick) comprises organic carbon matter-rich shale, silty shale, silty limestone, dolomite and siltstone (Figs. 5A and 5D), whereas the upper unit (315–330 m thick) includes limestone and thick-bedded dolomite (Fig. 5A).

The lower calcareous siliciclastic rocks of the Taft Formation conformably overlie those of the Sangestan Formation (Figs. 5A and 6A) in the Mehdiabad area. The lower unit displays abrupt changes in thickness, attaining <70 m in the ER but tapers rapidly westward (80 m) as it approaches the edge of the Back-Hill Fault. The transition from the lower unit to the upper unit rocks of the Taft Formation implies major changes in the sedimentary regime, from dominantly siliciclastic (lower unit) to calcareous (upper unit) (Fig. 5A). This change indicates the sag phase of the rift basin evolution (Betts et al., 1998; Large et al., 2002). The main ore horizon of Zn-Pb-Ba (Cu ± Ag) mineralization (ore horizon I) in the Mehdiabad deposit occurs within the lower unit and at the base of the upper unit of the Taft Formation (Figs. 4 and 5A).

2-2-3- The Abkuh Formation

The upper part of the Taft Formation is covered by 20 to 30 m thick cherty limestone, which forms the base of the Abkuh Formation (Figs. 4 and 5A). It is affected by irregular dolomitization, which cuts across the stratification. Above the cherty limestone is a unit of dolomitic shale with calcarenite and chert-bearing limestone, generally identical to those seen in the drill holes. The vertical transition between the cherty limestone and the dolomitic shale is gradual and is marked by a decrease of the chert content. Lenses and olistoliths of parareef

limestone (1-10 m size) are common in the upper part of the unit. A variably thick (0 to 60 m) level of para-reef limestone is also developed towards the top of the sequence. The lenticular shape of the unit is probably due to syn-sedimentary deformation, further inverted during tangential tectonic events. The top of this unit consists of 60-80 m thick conformable limy shale and laminar limestone (Fig. 4). Ore horizon II (non-sulfide ore) is located in the upper part of this unit (Figs. 4 and 5A). Chert-bearing bedded limestone, massive limestone and limy shale form the upper part of the Abkuh Formation (Fig. 4).

2-3- Main features of the Mehdiabad deposit

The Zn-Pb-Ba (Cu - Ag) mineralization of the Mehdiabad deposit occurs along two horizons that extend over a length of at least 3.4 km (Fig. 6). The trend of the mineralization is SW–NE, dipping 20°-30° to the northwest, consistent with the bedding of the host rock. The sulfide and non-sulfide ore of the ore horizon I (OH I) is hosted by the organic carbon-rich shale, silty limestone, dolomite and siltstone of the Taft Formation, whereas non-sulfide ores of the ore horizon II (OH II) are hosted in the limy shale and thinly bedded limestone of the Abkuh Formation (Fig. 6A, B) (Reichert et al., 2003). The main part of the ore body or OH I, the so-called Black hill ore (BH), Central valley orebody (CVOB) and East Ridge (ER), is located in a depression surrounded by hills and mountains (Fig. 6B). The second ore horizon (OH II) is only observed at the Calamine Mine (CM) (Fig. 6A). The BH, ER and CM expose completely oxidized ore (Figs. 6, 7 and 8A), whereas the CVOB mainly consist of sulfide mineralization. The restrictive occurrence of orebodies in the Taft Formation and in the Abkuh Formation documents the large vertical (stratigraphic) extent of the mineralization (Fig. 6). Detailed geological mapping of host rocks and synsedimentary structures in the Mehdiabad deposit shows

that Zn-Pb-Ba ($\text{Cu} \pm \text{Ag}$) mineralization occur to the east of the Black Hill syn-sedimentary fault in the LCSS (Figs. 4, 6A).

2-3-1- Ore Horizon I (OH I)

The BH vein and brecciated mineralization occurs adjacent to the Black-Hill syn-sedimentary fault (Fig. 8B). The BH ore consists of irregular veins of barite, quartz, chalcopyrite, pyrite, sphalerite, galena, Zn-Pb carbonate, and dolomite, but also limonite and hematite, so that it is often called the 'Black-Hill gossan'. The veins cut siltstone, dolomite, and shales of the lower unit of the Taft Formation. Veins range from <1 mm up to 80 cm in width and constitute up to ~80 % of the host rock. The abundance of veins increases stratigraphically upward, toward the barite ore zone. Stockwork veins similar to those at Mehdiabad have been interpreted as conduits through which mineralizing fluids vented onto the seafloor (Goodfellow and Lydon, 2007); therefore, this stockwork system likely represents the feeder zone of the deposit.

Toward the east, the BH mineralization changes into the CVOB (Fig. 6A). This lateral transition is gradual and is marked by a progressive increase of breccias and of the sulfide content. The organic matter-rich shale, dolomite and silty limestone of the Taft Formation (Fig. 5D), hosting the sulfide mineralization in the CVOB, are characterized by intensive and extensive brecciation. Coarse angular sulfide-matrix breccias occur throughout the CVOB but are spectacularly developed near the base where fragments of shale, dolomite, siltstone, and a variety of other material including barite, and sulfide breccia occur in a fine-grained sulfide matrix (Figs. 8, 9 and 12). Evidence that these breccias are related to the mineralizing process and not unrelated faulting includes: interlaying of the breccias with nonbreccia-textured sulfides (Fig. 12), breccias which are cut by and show alteration effects from later sulfide veins (Figs.

16D and 19A), and the fact that breccias are internal to the deposit and not spatially related to faults in the CVOB (Fig. 12). The well-developed sulfide-matrix breccia appears matrix-supported; clasts compose 20 to 75 percent of the volume of the rock. Clasts are angular to subrounded and range in size from <1 to >100 mm; the estimation of the maximum size of clasts is limited by the diameter of drill core (Figs. 18, 19 and 20). The most abundant ore minerals within the CVOB are sphalerite, galena, barite, pyrite, and rare chalcopyrite. Disseminated sulfides commonly occur as low-grade mineralization within the dolomite and the lower part of the Taft Formation. Towards the west, sulfide mineralization of the CVOB grades into synsedimentary stratiform horizons of ER (Fig. 6A).

The ER orebody mainly consists of stratabound sulfide and non-sulfide mineralization developed within the lower unit (calcareous shale, siltstone, silty limestone) of the Taft Formation (Figs. 6B, 7, 8B). It extends about 2 km along a SW-NE trend and dips at 10-30 degrees to the NW (Figs. 4 and 6B). The ER orebody, known from surface exposures and drilling, is a nearly flat lying, elongated stack of mineralized lenses. This zone consists of iron oxides (hematite, limonite), manganese oxi-hydroxides and Zn-Pb carbonates. The occurrence of galena, sphalerite, and pyrite in the drill cores sampled at 100m below the surface suggests the presence of a sulfide zone at depth. The textures of sulfides in the shale and siltstone are laminated and bedded. The ER occurs between the sandstone and oolitic limestone of the Sangestan and the silty limestone of the Taft Formation (Figs. 4 and 6B).

2-3-2- Ore Horizon II (OH II)

The CM represents the highest parts of the exposed mineralization of Mehdiabad (Fig. 6A). In contrast to the mixed main sulfide/non-sulfide orebody of the CVOB, the CM contains no

zinc- and lead sulfides. The CM is exposed in numerous adits, which follow the folded and duplicated ore horizon at different mining levels. The non-sulfide zinc-lead ore occurs on mainly three different levels within the Abkuh Formation, which is the result of fold repetition. Overall, the mineralization is predominantly hosted within the limy shale with intercalations of laminar limestone of the Abkuh Formation and particularly at the contact of the limy shale to the chert-bearing bedded limestone (Fig. 7). The host rocks to the CM are intensively faulted, brecciated, and locally mylonitized. Zinc-lead mineralization mainly consists of hemimorphite, smithsonite, and hydrozincite, locally associated with iron oxide/hydroxides (Reichert, 2007; Maghfouri, 2017). Only few sulfide minerals have been observed by microscope (Maghfouri, 2017). By analogy to OH I, the pre-existing sulfide mineral association of the OH II zone was probably galena-sphalerite-barite-pyrite. In the following we will only focus on the Ore Horizon I and readers may refer to Maghfouri et al. (2015) for complementary informations on OH II and CM.

3- Methodology

A field campaign and sampling were done during the autumn of 2015. To assess the geochemical characteristics of 250 systematic samples (least fractured and altered) of the ore-bearing zone and of host rocks were collected from the deposit for polished thin section examination. In order to correctly characterize their chemical compositions, 13 least-altered and fractured samples were chosen for major, trace, and rare-earth-elements (REE) analyses. Also 132 ore-samples were analysed by Mehdiabad Mining Company in the ACME Analytical Laboratories. Rock chips (~400 g) from each sample were carefully selected, washed with distilled water at room temperature, and then dried. Finally samples for whole-rock analyses were crushed and powdered in a tungsten carbide swing mill. Bulk-rock chemical analyses of major, trace, and REE elements were determined for all samples by Inductively Coupled Plasma-Emission

Spectrometry (ICP-ES) and Mass Spectrometry (ICP-MS) at ACME Analytical Laboratories, Vancouver, Canada. For the major elements, the detection limit is between 0.002 and 0.01 wt.%, 0.02–2 ppm for the trace elements, and 0.1–0.01 ppm for the REE elements. Major and trace element data are presented in Table 1 and REE elements data in Table 1.

Sulfide and sulfate minerals were separated from monomineralic samples using a microdrill and a binocular microscope. The microdrill was cleaned between each sample extraction to avoid contamination. Six $\delta^{34}\text{S}$ determinations by laser ablation-mass spectrometer were conducted at the Centres Científics i Tecnològics de la Universitat de Barcelona, Spain and in the Institute of Geochemistry, Chinese Academy of Sciences, Guizhou, China. The analytical procedure is similar to the one described in Huston et al. (1995). A 200- μm -thick polished slice of each sample was used for analyses. CO_2 separation was conducted on gas derived from samples.

4- Results

4-1- Sedimentary and geochemical characteristics of the host rocks

Most of the sedimentary rocks (Taft Formation) in the immediate vicinity of the Mehdiabad deposit are fine to very fine grained, gray-black, indistinctly to well-laminated organic carbon-rich shale, siltstone, dolomite, and limestone beds. These latter ones host the mineralization in the two horizons of the Mehdiabad deposit (Figs. 7 and 9). The main facies consists of organic carbon-rich shale and siltstone marked by interbeds of grey dolomitic siltstone, dolomite and thin-bedded black (carbonaceous) limestone (Fig. 9). The thickness of individual beds of organic matter-rich shale, siltstone and dolomite range from several centimeters to about 60 m, with an average of 40-50 m (Figs. 9A-9D). The shale and siltstone rocks may be calcareous, non-calcareous and/or carbonaceous. They are indistinctly to well laminated (<1 mm to 20 cm) (Fig.

9). Thin, sandy lenses or beds containing pale grey calcareous grains are locally observed (Fig. 9E). Organic matter-rich shale and siltstone are composed of quartz, carbonate, chert fragments, and carbonaceous material, occurring as sub-micron sized inclusions, organic matter nodules, pyrite and Fe-oxide minerals (Fig. 9F). The presence of abundant carbonates indicates that the sediments were not deposited at depths below the carbonate compensation depth (Peter and Scott, 1999). Sedimentary structures include normal layering, scours, pebble dets, and concretions composed of pyrite, calcite, ferroan dolomite and rare blebs of sphalerite. Soft-sediment deformation, represented by slump structures, indicates that the basin was tectonically active during sedimentation. Organic matter-rich shale and siltstone contains, very fine to coarse-grained, disseminated pyrite and/or fine grained sphalerite, galena and barite, of either syngenetic or diagenetic origin (Fig. 9).

The geochemical composition of the host rock shales from the Taft Formation is reported in Table 1. Unaltered shale samples from the Taft Formation have Mn and Fe contents ranging from 231 ppm to 6391 ppm and from 1.46% to 4.12%. respectively (Figs. 10 and 11). The V/Cr ratios for shale samples from the Mehdiabad deposit range from 0.60 to 4.43, whereas the V/(V+Ni) ratios range from 0.67 to 0.91 (Fig. 11). Samples are also marked by high U contents and the Th/U ratio ranges from 0.14 to 1.53, with an average value of 0.50.

4.2- Discrimination of ore facies and their distribution in the deposit

In the Central Valley, the sulfide orebodies of the Mehdiabad deposit do not crop out at the surface, and drill core logs and lithogeochemical data have been used to reconstruct a cross-section of the deposit (Fig. 12). The major sulfide mineralization is located east of the Black-Hill synsedimentary fault (Fig. 4). The sulfide zone at Mehdiabad (OH I) ranges greatly in thickness,

from few meters to more than 300 m. It displays a lens shape with a relatively flat top and a convex base (Fig. 12). The greatest thickness of high-grade mineralization (>7.52% Zn) is just to the east of the Black-Hill Fault, in the CVOB (Fig. 12). This zone also has the highest Zn grades (10–17%), highest Ag concentrations (21–51.7 g/t) and highest Pb grade (1.9-6%) (Maghfouri, 2017). Sulfide mineralization occurs as a single wedge within the organic matter-rich shale, siltstone, silty limestone and dolomite (Fig. 12). The western part (BH and CVOB) of the deposit has stratabound geometry, but toward the northeast (ER), as thickness decreases, the mineralization becomes stratiform and concordant with the host sedimentary rocks (Fig. 12).

The Mehdiabad deposit consists of five ore facies types, which are, from proximal (BH and CVOB) to distal (ER) relative to the synsedimentary fault (Fig. 12): (1) a stockwork zone, formed by sulfide veins accompanied by carbonate or quartz; (2) massive-replacement ore, formed by a massive sulfide zone and sulfide breccias; (3) bedded ore, consisting of a banded orebody made of pyrite, sphalerite, and disseminated galena; (4) a barite ore facies, which predominantly occur near the top of the sulfide mineralization; and (5) a distal facies, formed by chert beds with barite and disseminated pyrite.

- The stockwork or stringer zone

This zone consists of an irregular vein network cutting across the altered and brecciated siltstones, dolomites, and shales (Fig. 13). The veins are composed of fine grained barite, quartz, pyrite, chalcopyrite, sphalerite, galena, and dolomite (Fig. 13). This type of mineralization occurs adjacent to the Black-Hill Fault (Fig. 12). Dolomite and quartz occur in the core of the veins whereas sulfides are mainly found in the outer part. Veins range from <1 mm up to 1 m wide and may constitute up to ~20 % of the host rock (Fig. 13). The number of veins increases upward,

progressively passing into massive-replacement ore. Debris flow and abrupt facies changes occur adjacent to the stockwork zone (Fig. 12). Sulfide-rich veins generally present an alteration halo (silicification and dolomitization) and a partial replacement of pseudocubic pyrite by chalcopyrite and sphalerite is suggested by the irregular and crenulated grain boundaries in the pyrite-sphalerite aggregates (Fig.13). Copper contents in the stockwork zone may reach up to 2 wt% (Maghfouri, 2017). Sulfide veins also crosscut the barite ore facies and massive sulfides (Fig.13) suggesting multiple stages of stringer ore mineralization. Five main generations of veins and veinlets were identified within the stringer zone, based on the textural relationships and mineral paragenesis of the ore veins (V1 to V5) (Fig.13). These veinlets from early to late are (V1): chalcopyrite, quartz and barite-rich vein-veinlet's; (V2): sphalerite, galena, pyrite and barite-rich vein-veinlet's; (V3): sphalerite and galena-rich vein-veinlet's; (V4): barite-rich vein-veinlet's; and (V5): pyrite-bearing calcite vein-veinlet's. Banding, crustification, and comb textures indicate deposition in open space (Fig. 13). Wall rock fragments are commonly included in the veins and resulted in formation of cockade ore textures. Sulfide-cemented breccias are developed in areas of intense veining, particularly near the base of the deposit.

- The massive-replacement ore facies

The massive-replacement ore consists of a high-grade orebody that forms the thickest part of the deposit (Figs. 12 and 14A), principally in the CVOB, to a lesser degree, on the eastern margin of the BH. This facies comprises massive zones, replacement zones, brecciated sulfides, and a very irregular network of sulfide veins and veinlets (Figs. 14B to 14D). Mineralization in this facies is hosted by a series of stacked, planar-wedge shaped breccia bodies (Figs. 12 and 14). Some of the breccia units are dolomitized and only the lowermost breccia contains stratiform massive Zn-Pb-Fe sulfides. Dissolution breccias are perhaps the most distinctive type of

hydrothermal replacement at Mehdiabad and are characterized by coarse-grained sulfides that both replaced and infill veins and cavities within dolomitized limestone of the Taft Formation (Fig. 12). Well-developed sulfide-bearing breccias appear to be matrix-supported and clasts form 20 to 75 percent of the rock volume. Clasts are angular to subrounded and range in size from <1 to >100 mm, but estimation of the maximum size of clasts is limited by the diameter of drill core. The clasts are angular and are partially replaced and crosscut by thin veinlets of coarse-grained sulfide, which also form the matrix of the breccia. Most of the clasts are composed of hydrothermal dolomite (Fig. 14E), although clasts of shale, not dolomitized limestone of the Taft limestone, early sulfide aggregates, and massive barite were also observed. The bulk of the well-developed sulfide-bearing breccia occurs as a series of roughly stratiform lenses at or near the base of the dolomitized Taft Formation, in the hanging wall of the Black-Hill Fault.

The massive-replacement ore facies shows mineralogical zonation from bottom to top with an upward decrease in chalcopyrite and galena and increase in sphalerite, pyrite and barite (Fig. 12). Based on mineralogical and textural features, the massive-replacement ore facies is divided into five sub-facies (Figs. 12 and 14): (1) a chalcopyrite-sphalerite-galena-barite sub-facies (CSGB) (Figs. 12 and 14G), which occurs on the margin of the massive-replacement ore, especially in the BH area; (2) a sphalerite-galena-barite sub-facies (SGB) (Figs. 12, 14C, 14E and 14F); (3) a pyrite-sphalerite-barite-galena sub-facies (PSBG), mostly observed in the central part of the massive-replacement ore zone (Figs. 12, 14B, 14D and 14H); (4) a massive oxide Zn-Pb-Ba ore sub-facies (Figs. 12 and 14A); and (5) a barite ore sub-facies, located at the top of the system (Figs. 12 and 14A).

Stratiform beds and disseminations of barite also appear higher in the Mehdiabad stratigraphic section as well as locally in the massive-replacement ore facies within dolomitic

shales of the Taft Formation. The barite ore is composed of stratiform lenses, which are typically massive, but the bedding is locally defined by the presence of base metal sulfide or oxide layers (Fig. 14A). Fine- to coarse-grained euhedral to anhedral barite grains are intergrown with disseminated sphalerite, galena, and pyrite. Sulfide minerals also form thin to thick layers enclosed by barite. The massive sulfide-bearing barite lenses and beds locally grade in to quartz and apparently formed on the sea floor. Such textural evidence indicates that the barite predated and was partially replaced by the sulfides, but also co-precipitated with sulfides (Maghfouri, 2017). Barite ores of Mehdiabad can be divided into four textural types: massive, laminated, banded, and nodular barite ores and most of these are light gray in color. The massive ores are composed of high-grade (>85 %) BaSO₄ with minor sulfides and quartz. The barite crystals range from 0.05 to 0.18 mm. Generally, this massive variety is located at the top of the ore beds (Fig. 14A). Laminated barite commonly overlies the bedded sulfide ore facies and is distinguished by the number of sulfide and oxide layers interlaminated with barite. Banded barite ores, occurring at the base of the ore beds are of low grade, with relatively high sulfide contents.

- The bedded ore facies

The facies (bedded barite) is located above and to the east (bedded sulfide) of the massive-replacement ore facies in the CVOB (Fig. 12) and in the ER (Fig. 15). It is characterized by laminated sphalerite, galena, and iron sulfide (pyrite and marcasite) (Fig. 15). In some core samples (e.g., DDH 806, 810), iron sulfides predominate at the top of the zone, whereas the amount of base metal sulfides increases with depth. The lateral transition between the bedded ore and the massive-replacement ore facies is generally gradual and is recognized by an increase in laminae. The contact with the underlying silty limestone and dolomite is usually sharp (Figs. 16B and 16C). This facies displays stratiform banded sulfide-rich layers (Figs. 16D to G), concordant

with the host rock bedding (Fig. 16A). Pyrite, galena, and sphalerite are the dominant sulfide minerals (Fig. 16). Fine bedding defined by the distribution of sulfides associated with silt and organic matter may be observed from a macroscopic to microscopic scale (Fig. 16). The bedded ore facies can be divided into low-grade (1–2 % Zn + Pb) and high-grade (>4 % Zn + Pb) units (Fig. 16). At the microscopic scale, pyrite forms framboidal aggregates. Sphalerite laminae are less common than pyrite laminae (Fig. 16D); they are only ubiquitous in the high-grade part of the bedded ore facies.

- The distal facies

The banded silicic dolomite is only exposed in the eastern part of the deposit (ER; Figs. 4 and 10A) and is associated with strong Fe and Mn anomalies (Figs 16 H, I, J). This facies has no economic value and represents the most distal part of the hydrothermal deposit. The boundary between the bedded ore and the distal facies is gradual and is indicated by the absence of sphalerite and the increase of disseminated pyrite with respect to laminated pyrite (Figs 16 I, J, K). Disseminated barite occurs in association with Fe- and Mn-rich dolomite in the distal facies.

4-3- Relationships between mineralization, host rocks and faults

Similar to the sedimentary host rocks, the sulfides display characteristic layering on a macroscopic and microscopic scale (Figs. 16, 17, 18 and 19). Despite complex banding and a high variability of textural and compositional variations within the laminae, we may distinguish three end-member types of microbands in the ore (Figs. 16 and 17). The organic matter-rich shale layers (Type 1) are commonly less than 100 µm thick (Figs. 17A and 17B); they are principally composed of illite, carbonaceous matter, pyrite, and quartz; and are interpreted to be the product of normal pelagic sedimentation from the anoxic water column. The shale and silt

layers (Type 2) are commonly 100 to 500 μm thick, but laminae up to several centimeters thick are also present (Fig. 17E). The zinc-lead-iron-rich sulfide layers (Type 3) are commonly 50 to 300 μm thick (Figs. 17C and 17D) and consist of uniform mixtures of sphalerite-galena-pyrite with minor barite and quartz and rare chalcopyrite. Sphalerite and pyrite are the main sulfide minerals and the sphalerite/galena ratio varies from 2:1 to 9:1. Although it is common for the three end-member microband types to be regularly interlayered in a given sample (Figs. 16 and 17), mixture of these and end members may also occur (i.e. type 1 + type 3 in figs. 13B and 14A; type 2 + type 3 in figs. 16D, 16E, 17C and 17D). From ore microscopy, different types of pyrites were observed within the shales collected from the drill holes. We may distinguish colloform pyrite (Fig. 17F) from framboid pyrite (Fig. 17G). Beds of galena and sphalerite are also present (Figs. 16G and 17H).

Rounded and subrounded sulfide clasts are common within the thick ($> 1\text{cm}$) turbidite beds (Figs 18A and 18B); these are interpreted as the result of a reworking of sulfide layers originally deposited on the sea floor. Compactional load-cast features are also common in the turbidites rocks, locally bending or folding the sulfide-rich layers (Figs. 18D, 18E and 18F). Detrital clasts of carbonate and lithic fragments occur at the base of the sedimentary rocks and they display evidence of compressive deformation, beneath the base metal layer. Flame structures are also visible in the sphalerite-galena-pyrite beds, extending upward into the mass flow units. Sulfide microbreccia layers are common in the different ores (Fig.18G). These layers are composed of rounded to angular clasts of sulfide minerals enclosed in a matrix of sediments with disseminated sulfides (Fig. 18G). They are typically non-graded, measuring 0.7 to 2 mm in thickness, and indicate open-cast reworking of a proximal source of preexisting sulfides.

Mineralization occurs to the east of the Black-Hill (BH) Fault (Fig. 8E), which was intersected by drilling. In drill core, evidence of hydrothermal alteration, marked by dolomitization, silicification, and sulfide mineralization, was recognized as extremely variable (Fig.13A). This corresponds to the stockwork facies, described in the previous section. Whereas the thickness of the sedimentary units of the Sangestan Formation beneath the Taft Formation is relatively constant, the thickness of the Taft Formation increases in the hanging-wall of the BH fault. As the compressive deformation related to Tertiary inverted tectonics was quite gentle, such present-day variations probably represent the original deposition, most likely controlled by the syn-sedimentary activity of the BH normal fault. It is also worth to notice that the thickest and highest grades of Zn-Pb mineralization, as well as pervasive hydrothermal alteration occur within the Taft Formation in the hanging- wall rocks of this major fault as well (Figs.12 and 19). Exploration drilling along the orebody also indicates that major facies changes occur in the Taft Formation, with shallow-water shelf facies to the east of the BH fault and basinal rocks in the CVOB area.

Additional evidence of the syn-sedimentary activity of the BH fault is the presence of mass flows (Fig.20). The latter correspond to sedimentary breccias that are common throughout the Taft Formation (Fig.20). To the west and to the south, the mass flows become thicker as the size of fragments increases; this indicates that the clasts are sourced from mass shedding along the tectonically active BH fault (Fig.20).

4-4- Sulfur isotopes: results and discussion

Sulfur isotopes at the Mehdiabad deposit were measured from 52 samples of pyrite, chalcopyrite, barite, sphalerite, and galena collected from all ore facies except the distal facies.

The analytical procedure is given in the supplementary material and the results of sulfur isotope analyses are listed in the table 2 and shown on Figure 17. The sulfides have a wide range of $\delta^{34}\text{S}$ values from -22‰ to 22.4‰ , with a majority at about -14 to 4 per mil (Fig. 21). Extremely low $\delta^{34}\text{S}$ values in the range of -20.1‰ to -22‰ are only observed in bedded ore facies pyrite (Fig. 21A), whereas the massive ore and stockwork sulfides have higher $\delta^{34}\text{S}$ values ranging between -13.7‰ and $+4.6\text{‰}$ (avg; -4.2‰).

The $\delta^{34}\text{S}$ values of sphalerite ($n = 11$) fall within the range of -18‰ to $+4.6\text{‰}$. There is a close correlation between texture and $\delta^{34}\text{S}$ values. Fine-grained, framboidal and colloform sphalerite has primarily negative values (-17.53‰ $n = 3$), whereas coarser-grained sulfides found at the base of the deposit in the massive-replacement ore (-6.27‰ for $n = 4$) and in the stockwork zone (-1.35‰ for $n = 4$) have significantly higher values (Figs. 21B). A similar pattern is observed for galena, whose $\delta^{34}\text{S}$ values range from -18.1‰ to $+4.4\text{‰}$ ($n = 16$). Galena from the massive-replacement ore facies (-9.34‰ for $n = 5$) and from the feeder zone (-2.72‰) are isotopically different from the galena locally intergrown with framboidal pyrite of the bedded ore (-17‰ to -18.1‰).

$\delta^{34}\text{S}$ values in chalcopyrite ($n = 9$) range from -9.6‰ to $+4.6\text{‰}$ (Fig. 21B). Vein-veinlet chalcopyrite hosted by hydrothermal dolomite in the deep feeder zone has the $\delta^{34}\text{S}$ between 4.6‰ and -7.3‰ per mil, whereas massive-replacement ore has a mean $\delta^{34}\text{S}$ of -3.6‰ . Barite intergrown with sulfides in the feeder zone ($n = 3$) has $\delta^{34}\text{S}$ values ranging from $+22.4\text{‰}$ to $+22$ per mil. A similar range of values was obtained in the bedded ore ($+17.8\text{‰}$ to $+22.3$ for $n = 4$) and in the massive-replacement ore facies, ($+19.65\text{‰}$ to 20.1‰ for $n = 3$).

5- Discussion

5-1- Conditions of deposition of Mehdiabad host rocks

Interbedded black siltstone documents the ambient depositional conditions and hosts the ores and mineralization at Mehdiabad. The dolomite beds represent a small volume in the Mehdiabad Basin. This facies is characterized by an accumulation of fine-grained hemi-pelagic suspensions in a quiet, reduced, sub-wave base subaqueous environment. Thicker, discrete fine-grained shale intervals are rich in pyrite and organic matter. This sedimentary facies, which is the most common at the base of the Taft Formation and in the middle part of Abkuh Formation, is considered to be relatively poor in clastic sediment input, and is interpreted to indicate the maximum of water depth and/or shoreline retreat in the basin.

Several geochemical indices based on redox-sensitive major and trace elements have been used to determine the paleo-oxygenation conditions of the water column (Hoffman et al., 1998; Jones and Manning, 1994; Quinby-Hunt and Wilde, 1994; Whitehead, 1973). Among the major elements, Fe and Mn can be used to indicate bottom water conditions as their solubility is affected by ambient pE and pH (Whitehead, 1973). Quinby-Hunt and Wilde (1994) used solubilities of Mn and Fe to determine the paleo oxygenation levels and the prevailing oxidizing agent in the water column. In oxic bottom water conditions, oxygen is the primary oxidizing agent; Quinby-Hunt and Wilde (1994) argued that low Mn and Fe solubilities would favor their fixation in sediment, yielding high Mn and Fe contents in shales. In anoxic bottom water conditions with nitrate as the primary oxidizing agent, Mn solubility increases whereas pE decreases, resulting in low Mn contents in the sediments. In contrast, Fe remains bound in oxide minerals within the sediments, although high Fe solubility at very low pE values may lead to a decrease of the Fe content in sediments. Although low Mn and high Fe concentrations may characterize nitrate-reducing conditions, low Mn and high Fe may also occur in anoxic

conditions in which sulfate is the primary oxidizing agent, as Mn remains dissolved in the water column and reduced sulfur fixes Fe in the sediments. The low Mn contents in the unaltered shale of the Taft Formation are consistent with a deposition in anoxic bottom waters (Fig. 10A), whereas the Fe contents may support oxic or anoxic sulfate-reducing conditions (Fig. 10). However, Mn was shown to have been mobile during hydrothermal alteration, compared to Fe, thus caution must be taken to use this proxy in such polyphase mineralizing system.

In addition, the low oxygen levels are strongly limited the benthic activity, and ichnofabric criteria cannot be used to discriminate between oxic (>2.0 ml O_2/l H_2O), suboxic (0.5 – 2.0 ml O_2/l H_2O), dysoxic (0.2 – 0.5 ml O_2/l H_2O) and anoxic (<0.2 ml O_2/l H_2O) conditions. Hoffman et al. (1998) proposed the V/Cr and V/(V + Ni) ratios, which are sensitive to anoxic and dysoxic conditions. Whereas V, by bounding to organic matter, is considered as a useful indicator of its preservation (Brumsack, 1986), Cr, which substitutes for Al in clays or occurs as chromite, is associated with the detrital fraction of the shale (Jones and Manning, 1994), Ni is preferentially incorporated into the sulfide minerals. The V/Cr and V/(V+Ni) ratios of the Mehdiabad shales indicate anoxic to dysoxic conditions (Fig. 11). The high U content of the Mehdiabad shales is typical of sediments deposited under anoxic conditions (Nath et al. 1997), and results from the reduction of U^{+6} into U^{+4} , which is insoluble and deposited in the sediments. The low value for the Th/U ratio (av. 0.85) shows that no fractionation occurred between U and Th, unlike what is classically observed in shallow oxic environment (Whittaker and Kyser 1993; Wingal and Twitchett, 1996).

Most SEDEX deposits are hosted by basinal marine, reduced facies, fine-grained sedimentary rocks that consist mostly of carbonaceous siltstone and shale (Large et al., 1996; Kelley et al., 2004a,b; Leach et al., 2005; Goodfellow and Lydon, 2007; Leach et al., 2010;

Wilkinson, 2014; Gadd et al., 2016a,b). These sedimentary rocks generally represent pelagic and hemipelagic sediments. In some deposits (e.g., Sullivan), pelagic and/or hemipelagic sedimentary rocks are interbedded with turbiditic siltstones and sandstones, and locally include coarse-grained clastic sediments, derived from the erosion of uplifted blocks during extensional tectonism (Goodfellow and Lydon, 2007; Magnall et al., 2015; Gadd et al., 2016b; Magnall et al., 2016; Gadd et al., 2016a). In addition, the geochemical redox proxies used to characterize the shales hosting the mineralization support an anoxic environment and our results are very similar to what has been reported for many other SEDEX deposits (Goodfellow and Lydon, 2007; Sáez et al., 2011; Yarmohammadi et al. 2016; Boveiri et al., 2017). In the case of the Irish-type or MVT deposits, the dominant lithotypes are limestone and dolomite (Lee and Wilkinson, 2002; Wilkinson, 2003; Reed and Wallace, 2004; Wilkinson, 2010; Elliott, 2015); MVT deposits generally form in reduced environments (Pfaff et al., 2010).

5-2- Textural and metal zonation

One of the most characteristic features of SEDEX deposits is the vertical and lateral zonation of textures, minerals, and metals (Goodfellow and Lydon, 2007). Deep drill holes have shown that the Taft Formation is mineralized in at least three areas (BH, CVOB and ER) in the hanging wall of the major Black-Hill synsedimentary fault (Figs. 6A, 12). The largest of such sulfide zones located so far are within the CVOB between the BH and ER orebodies (Maghfouri, 2017) (Fig.12). In this deposit, there is a gradual change, away from the stockwork zone (or BH synsedimentary fault), from veins and massive replacement textures in the massive-replacement ore (CVOB) to sedimentary textures (laminated, disseminated, and framboidal) in the bedded ore (ER) and in the distal facies (Maghfouri, 2017) (Fig. 12). This systematic trend is accompanied

by a decrease in the number and thickness of individual sulfide beds (Fig.12). The ore thickness and grade in this deposit decrease also systematically away from the Black-Hill Fault (Fig.12).

The deposit is also marked by lateral chemical zonation. In terms of metal ratios, there is a decrease in the Pb/Zn ratio from the massive-replacement ore, with a maximum value of combined base metals ($Zn+Pb=12.5\%$) in the thickest part of the deposit toward the bedded ore (Maghfouri, 2017) (Fig. 12) (Table 3). There is still a gradual increase in zinc grade to the CVOB, away from to Black-Hill Fault (Fig.12). These lateral variations in zinc grade demonstrate systematic changes from as much as 40 vol percent sphalerite to about 5 vol percent sphalerite without any stratigraphic correlation. This is accompanied by a decrease in Pb grades towards the northeast of the deposit (Maghfouri, 2017). Iron is mostly contained in pyrite, and the Fe/Pb and Fe/Zn ratios, together with the Mn content increase away from the massive-replacement ore as base metals decrease (Fig. 12).

Copper is strongly zoned, and Cu-rich ores have been found close to the Black-Hill Fault (Maghfouri, 2017) (Table 3). Silver displays a zonation pattern similar to zinc throughout much of the deposit, although the highest values occur in the central part (CVOB) of the sulfide zone in the area of high lead values (Fig. 12). Barium contents are uniformly high at Mehdiabad. The highest values occur adjacent to the Black-Hill Fault (Fig.12). There is a broad overlapping lateral zonation marked from west to the northeast by the following trend $Cu > Pb > Zn > Fe$. Maghfouri (2017) interpreted this trend as a manifestation of the prevailing direction of lateral fluid flow and solubility-controlled precipitation through the mineralizing system, possibly also due to a thermal gradient or a chemical gradient independent of temperature (e.g., pH, oxidation state, or chemical kinetic effect). The relatively high Cu and low Fe in the western part of the

deposit is inferred to represent the location of fluid entry or venting site into the Mehdiabad orebody.

Moreover, the Mehdiabad deposit also shows a strong vertical zonation, especially within the massive-replacement ore (Fig. 7). There is an upward increase of Zn, Pb, and Ba/(Zn + Pb) and a decrease of Cu/(Zn + Pb), Ag/(Pb + Ag), and Cu (Maghfouri, 2017) (Fig. 12) (Table 3). The vertical geochemical variations may reflect changes in the overall abundances of ore minerals, or alternatively, they could reflect the chemical composition of the sulfides. As an example, there is a greater abundance of iron sulfide in massive-replacement ore compared with vein sulfide ore, and sphalerite in vein ores is Fe-rich compared with that in other ore types.

5-3- Mineralizing process of the Mehdiabad deposit

5-3-1- Syn-sedimentary/diagenetic versus replacement

On the basis of mineralogical and textural studies, Maghfouri (2017) considered a three-stage (Fig. 22) mineralization model at Mehdiabad. Early sulfides and Ba sulfate (Py1, Sph1, Gn1, Cpy1 and Ba1) formed by exhalation and precipitated from an overlying water column during an early diagenetic stage (Stage 1), whereas late sulfides and Ba sulfate (Py2, Sph2, Gn2, Cpy2 and Ba2) resulted from a subsurface hydrothermal replacement event (Stage 2) (Fig. 19). The third stage is seen as a covering layer of barite (Ba3) that also contains pyrite (Py3), sphalerite (Sph3) and galena (Gn3) (Fig. 12).

The early syn-diagenetic stage is characterized by the presence of framboidal pyrite (Py1) (Fig.17G), colloform pyrite (Fig.17F) and laminated and disseminated sphalerite (Sph1), galena (Gn 1) and barite (Ba 1) (Figs. 9C, 17C,D, 16G, D and 17H), commonly found in the bedded ore

and rarely in the distal facies (Fig. 22). In addition to their shape, the small size of such pyrite grains rather supports a sedimentary-diagenetic origin. The sphalerite-galena-pyrite-rich bands are intricately interlayered with shale and siltstone beds and carbonate turbidites (Fig. 16D); the boundary between mineralized and non-mineralized siltstone beds is sharp and cannot be explained by a replacement process (Figs. 16 and 17C). In addition, there is a lack of significant sphalerite and galena in many of the siltstone, whereas this lithotype is favorable for a replacement model (Ghasemi, 2006). The relations between sulfide minerals and some sedimentary structures (Fig. 18) clearly indicate that sulfide layers of Stage 1 were already present on the seafloor when the turbidite units formed.

The second ore stage occurs as massive ore, veins, and replacements, located mainly in the massive-replacement ore and stockwork zones (Fig. 19, 14 and 22). This stage is also characterized by complex mineral assemblages (Fig. 22). Detailed mineralogical and textural observation revealed that replacement is the dominant process, with early sulfide minerals replaced by late and coarse-grained sulfide minerals (Figs. 14 and 19). Textural styles include bedding-parallel to cross-cutting extensional brecciation and veining, delicate to massive-replacement of host rock, and dissolution of limestones to form open spaces that were subsequently completely filled with sulfides (Figs. 18, 19 and 20). Based on geological, textural and mineralogical studies, Maghfouri, (2017) suggested that breccia's and open space filling mineralization (second stage) at Mehdiabad formed sub-seafloor epigenetically from mineralizing solutions that flowed laterally through the still permeable sediments along deep channel-ways from the Black-Hill Fault.

Second stage of mineralization (massive-replacement) is overlain by bedded barite (Ba₃) (third stage) (Fig. 22). The vertical transition between the massive-replacement ore facies and the

stratiform barite beds is gradual and is recognized by increasing of bedded barite mineralization. This stage of mineralization is characterized by massive to laminated barite and minor pyrite (Py₃), sphalerite (Sph₃), and galena (Gn₃) (Fig. 22).

5-3-2- The role of the Black-Hill Fault

Detailed studies of the distribution of sedimentary facies suggest that fault-related, hanging-wall subsidence began along the BH fault system in the Early Cretaceous with contemporaneous sedimentation leading to the deposition of the Taft Formation. In addition to these sedimentary processes, normal faulting also played a fundamental role for the localization of mineralization and hydrothermal alteration at Mehdiabad (Reichert et al., 2003; Reichert, 2007; Maghfouri et al., 2015; Pourfaraj et al., 2016; Maghfouri, 2017). The maximum thickness of sulfides is found close to the Black-Hill Fault, suggesting that this syn-sedimentary structure acted as the most important conduit for hydrothermal fluid flow (Fig. 12). This area, which is characterized by abundant tectonic and sedimentary breccias, displays evidence for the development of a vein network. Stockwork veins, similar to those at Mehdiabad have been interpreted as conduits through which mineralizing fluids vented to the seafloor (Goodfellow, 2004); therefore, the stockwork system likely represents the feeder zone. Based on fluid inclusion studies, Maghfouri and Hosseinzadeh (2018) suggested that the neighboring Mansourabad deposit formed from basinal, hydrothermal fluids rising through synsedimentary faults to the seafloor, which might be similar in the Mehdiabad deposit. Maghfouri (2017) noticed the presence of abundant clasts of sulfides, which are identical in texture and composition to the ore lens material, immediately underlying the mass flows (Fig. 20). This provides convincing evidence that base metal sulfide mineralization at Mehdiabad occur in the uppermost level of sea-floor sediments, where the laminated ores were susceptible to be affected by syn-depositional faulting and erosion caused

by mass flows. The relationship between ore lenses, organic matter-rich shale, and mass-flow breccias strongly suggests that the BH fault system controlled both ore fluid migration and sediment breccia deposition, with mineralization limited to the deepest part of the sub-basin adjacent to the active fault system. A variety of sulfide mineral (galena, sphalerite, and pyrite) clasts occur in the mass-flow breccias (Figs. 20 a,b,c,d). In SEDEX deposits, such syn-sedimentary faults are recognized as the pathways for metal-bearing, hydrothermal fluids to enter onto the seafloor (Logan, 1979; Large et al., 1996; Large et al., 2005). Tectono-sedimentary controls have been well documented in other SEDEX deposits (i.e. the Ikalukrok Unit in the Anarraaq deposit; Kelley et al., 2004a,b). Away from the fault, the outer northeastern margin of the deposit consists of banded silicic dolomite with pyrite, chert, and manganese and is interpreted to represent more distal hydrothermal exhalites (Fig. 15).

5-3-3- Processes involved in the formation of sulfide minerals

The $\delta^{34}\text{S}$ values found at Mehdiabad are much lower than expected if sulfur would have derived from a magmatic source. The source of the sulfur is probably seawater sulfate. The barite $\delta^{34}\text{S}$ values are comparable to the results reported by Maghfouri (2017) and are identical to the range of values (+15 to +21‰) for Lower Cretaceous seawater sulfate (Claypool et al., 1980; Bottrell and Newton, 2006). This suggests that barite precipitation at Mehdiabad is related to unmodified Cretaceous marine sulfates, as inferred for barite in the Carboniferous Red Dog deposit (Johnson et al., 2004; Kelley et al., 2004a) (Fig. 21C). The transformation of sulfate to sulfide can be achieved either by bacteriogenic sulfate reduction (BSR) or by thermochemical sulfate reduction (TSR), which implies isotopic fractionations greater than 20‰ (Ohmoto et al. 1990). As temperature increases, fractionation would decrease. Nevertheless, TSR processes usually reach metastability and therefore little or no isotopic fractionation occurs between sulfur

species (Cross and Bottrell, 2000). The Mehdiabad deposit is mineralogically and chemically zoned away from the feeder zone, reflecting steep temperature gradients. Silicic alteration found around the feeder zone suggests high-temperature conditions for this part of the deposit. However, the zonation, controlled mostly by zone refining and sub-sea floor replacement in the massive ore (Goodfellow and Lydon 2007; Wilkinson et al., 2005; Kelley et al., 2004a; Johnson et al., 2004), along with the $\delta^{34}\text{S}$ difference found between the bedded ore and the massive-replacement ore, suggest that, at the margins of the system (bedded ore and distal facies), temperatures would have been low enough to sustain BSR process. Maghfouri and Hosseinzadeh (2018) reported a similar zone-refining paragenetic succession in the vent complex (massive ore) of the Mansourabad deposit. If the temperature of ore-forming fluids is less than 150° C, microbial processes can play a significant role in sulfate reduction (Southam and Saunders, 2005). The presence of framboidal and colloform pyrite in the bedded ore facies within the lower parts of the massive-replacement ore supports this interpretation. Although the development of pyrite framboids can occasionally occur by means of abiotic processes (Berner, 1969; Prol-Ledesma et al., 2010), laminated and framboidal pyrites are usually associated with biogenic deposition (Alfonso et al., 2005).

Sulfides from the bedded ore and parts of massive-replacement ore and feeder zone have $\delta^{34}\text{S}$ values between -5.2 and -22‰. Those low values alone could account for a BSR process if it occurred in a closed or semi-closed system for sulfates, but open for sulfide following a Rayleigh distillation process (e.g. Canet et al. 2005). Microbial-mediated sulfate reduction is known to have played a prominent in situ role in the formation of SEDEX deposits (Southam and Saunders, 2005; Gadd et al., 2016a) (Fig. 21C). Compared to other sulfides, pyrite has generally lower $\delta^{34}\text{S}$ values, a feature typically reported in SEDEX and Irish type deposits (e.g., Anderson

et al., 1998; Large et al., 1996; Fallick et al., 2001; Ireland et al., 2004a,b; Kelley et al., 2004a; Leach et al., 2005; Gadd et al., 2016a). Colloform pyrite (stage 1) from the massive-replacement ore and feeder zone was replaced by galena and sphalerite (stage 2) with $\delta^{34}\text{S}$ values from +3.1 to +4.4 and from +2.3 to +4.6‰, respectively. These high isotopic values of late sulfides cannot be produced by BSR in the proposed closed system for sulfate and suggest at least a contribution of TSR to the sulfide sulfur that formed these minerals.

5-4- Type of Mehdiabad deposit

Different genetic types have been proposed for the Mehdiabad Zn-Pb-Ba-(Cu-Ag) deposit, including a MVT-type (Leach et al., 2010) and a sub-seafloor replacement SEDEX-type mineralization (Maghfouri et al., 2015; Maghfouri, 2017). Except for Leach et al. (2010) review paper, there is no mention of the Mehdiabad MVT-type deposit in any other article. However, our results are in not in agreement with the scenario proposed by Leach et al. (2010) for this deposit. Based on the location of mineralization in the Taft Formation, given that this formation is generally known as a carbonatic formation, Leach et al. (2010) considered Mehdiabad as a MVT-type deposit. The total thickness of the Taft Formation in the deposit area is about 400 meters (Figs. 5A and 23). However, Zn-Pb-Ba-(Cu-Ag) mineralization does not occur in the whole Taft Formation, but is located in 120 meters of the lower part of this formation (Figs. 5A, 6C, 7, 23 and 24). From these 120 meters, 80 meters consist of organic matter-rich shale, fine-grained black carbonate-rich siltstone, dolomite and the rest is carbonate (Figs. 5A, 23 and 24). Therefore, the main host rocks of mineralization in the Mehdiabad deposit are the clastic-carbonatic lithotypes: a feature similar to other SEDEX-type deposits (Figs. 23 and 24).

Most MVT Pb–Zn deposits are located in the regional thrust belts and along secondary reverse faults, and most researchers have considered that the Pb–Zn ore deposition was controlled by thrusting (Leach et al., 2005; 2010; Wilkinson, 2014). In fact, the ages of the carbonate hosted Pb–Zn mineralization stages in the MVT-type deposits are closely related to the timing of thrust faulting in the region of mineralization in the world. The MVT carbonate hosted Pb–Zn mineralization's are epigenetic (formed after the sedimentation of the host rock) (Leach et al., 2005; 2010; Wilkinson, 2014). Unlike most MVT-type deposits, Mehdiabad typically occurs within fine-grained marine sedimentary rocks (lower Unit of Taft Formation) (Figs. 23B and 24) overlying coarse-grained ones (Sangestan Formation) deposited in a continental rift basin, and are commonly spatially associated with the BH synsedimentary normal fault (Figs. 6A, 8B, 12 and 23A). The sharp contact between the Sangestan Formation and the Taft Formation near Mehdiabad (Fig. 4c) suggests the presence of a synsedimentary fault (BH), which may have acted as a pathway for the mineralizing fluids (Figs. 6A and 12). The hydrothermal alteration and the presence of mass flows observed in the proximal feeder zone at Mehdiabad (Figs. 12, 13, 20 and 23A) is reported in several SEDEX deposits (Tom, Jason, Vangorda, Rammelsberg, Century, Red Dog) and is generally interpreted to precede or overlap the mineralization event (Broadbent et al., 1998; Leach et al., 2005; Slack et al., 2004).

In the Mehdiabad deposit the ore minerals are fine-grained, and there are layered, banded and replacement textures (Figs. 16, 17 and 24). The chemistry of the ore minerals (e.g. higher Ag and Cu contents) reflects the high temperatures of the ore-forming fluid and is not typical of many MVT deposits. The feeder zone of the Mehdiabad deposit contains a copper mineralization (Figs. 13 and 24 (drill hole 7905)) that show a zonal pattern away from the main fault zone, with copper concentrating closest to the BH synsedimentary normal fault in the most altered rocks

(Figs. 12 and 24 (drill hole 7905)). Again this is unlike the numerous deposits typical of MVT districts.

Wilkinson (2014) defines sub-seafloor replacement deposits as deposits where the ores are hosted by carbonate and clastic rocks; they also have synsedimentary normal faults controlling the deposition of massive sulfides, formally known as sub-seafloor replacement SEDEX. These deposits (e.g., Anarraq) have many morphological and textural similarities to Irish-type ore bodies. In the sub-seafloor replacement SEDEX deposits, synsedimentary faults controlled the formation of seafloor depressions that may have trapped brines, acted as conduits for ore fluids, and generated debris flow breccias that can become mineralized (Figs. 6A, 12, 18, 19, 20, 23 and 24). Based on Wilkinson (2014), it is clear that the sub-seafloor replacement SEDEX-type deposits ore bodies are more akin to typical examples of the SEDEX family than to typical examples of the MVT clan. The principal similarity that the sub-seafloor replacement SEDEX-type deposits ores have with MVT deposits is that a part of the mineralization's can be carbonate-hosted but, as noted earlier, this is a poor classification criterion for sediment-hosted deposits given that classic shale-hosted SEDEX camps can include carbonate-hosted deposits, and many SEDEX deposits are hosted by carbonate-rich mudstones or siltstones (Wilkinson, 2014). Similar to SEDEX-type deposits, the Mehdiabad mineralization is hosted by organic matter-rich shale, fine-grained black carbonate-rich siltstone, and dolomite interlayered with a sandstone sequence (Figs. 9, 16, 17, 23A and 24). SEDEX deposits are generally tabular in shape and predominantly contain various amounts of Zn, Pb, Cu, Au, Ag, Ba, Fe and Mn oxides. The geometry of the Mehdiabad deposit, shown in Figs. 6A and 12, and its mineralogical, structural and textural features suggest that the deposit is a SEDEX-style. However, typically, in these sub-seafloor replacement SEDEX-type deposits, there can be some evidence of exhalation

of the fluids onto, or below the seafloor, to form laminated and stratabound ores (see Wilkinson, 2014). There is evidence of laminated ores, derived from exhalative processes at the Mehdiabad deposit (Figs. 16, 17 and 24). Mehdiabad has characteristics similar to those of the Anarraaq sub-seafloor replacement SEDEX deposit, located in the Red Dog basin, Canada. These are: the exhalative textures deposited on the seafloor, the sub-seafloor replacement textures and the synsedimentary fault that controls the mineralization parameters (Kelley et al., 2004).

We have refined the sub-seafloor replacement SEDEX genetic type for Mehdiabad (Maghfouri et al., 2015; Maghfouri, 2017), at a level of detail useful to explain the prevailing textural and chemical characteristics of the deposit (Figs. 17–18). In fact, the characteristics defining Mehdiabad as a sub-seafloor replacement SEDEX-type deposit are: 1) the marine extensional basin setting; 2) the clastic-carbonatic marine sedimentary host rock sequence (Figs. 5A, 8A, 9, 23 and 24); 3) the bedding-parallel Zn-Pb-Ba-(Cu-Ag) sulfide lenses (Figs. 8, 12, 15, 16 and 17); 4) the synsedimentary faults that controlled the formation of seafloor depressions that may have trapped the metal-rich brines, acted as conduits for ore fluids, and generated debris flow breccias that can become mineralized (Figs. 6A, 12 and 20); 5) the well-developed lateral and vertical geochemical and mineralogical zoned patterns, extending for kilometers to tens of meters from the BH synsedimentary fault (Fig. 12); 6) similarly to other SEDEX-type deposits, the barite content in the Mehdiabad deposit is variable and occurs above the sulfide mineralization (Figs. 6A, 12 and 14A); 7) the sulfides display a variety of textures; the most characteristic are fine-grained layered and banded lithotypes, but coarser-grained brecciated, veined and replacement textures are also commonly observed (Figs. 13, 14, 16, 17, 18, 19 and 24). Based on the above characteristics, we consider Mehdiabad as a vent-proximal sub-seafloor

replacement clastic-carbonate hosted SEDEX-type deposit, and not a typical Mississippi Valley-type deposit (MVT) (Leach et al., 2005).

5-5- Model for the formation of the hypogene mineralization in the Mehdiabad deposit

The Mehdiabad deposit has a prominent feeder zone and a well-developed massive-replacement ore, forming what has been interpreted as a vent complex, similar to many well-known shale-hosted massive-replacement sulfide deposits. Thus, the Mehdiabad deposit can be classified as a vent-proximal sub-seafloor replacement clastic-carbonate hosted SEDEX-type deposit (cf. Lydon, 1996; Sangster 2002; Large et al. 2005; Goodfellow and Lydon 2007).

The formation of the Mehdiabad deposit was strongly affected by the structural setting of the basin (Fig. 25). The stratiform and stratabound orebodies of this deposit occur in the thickest section of organic matter-rich shale and dolomite units, immediately adjacent to the synsedimentary Black-Hill normal fault (Maghfouri. 2017) (Fig. 25). This fault is considered to be a favorable conduit for hydrothermal fluid upflow and played an important role for the formation of the deposit, similar to other deposits described by Large et al. (2002), Leach et al. (2005), and Rajabi et al. (2014). In the Mehdiabad deposit, presumed feeder zones developed next to the syn-sedimentary normal fault (Fig. 25). The metal zonation patterns, the presence of synsedimentary breccias and debris flows, the increase of hydrothermal alteration and of the temperature of the ore fluids towards the faults, suggest that this fault was active during ore deposition and provided major conduits for the metalliferous hydrothermal fluids.

In contrast to the laminated shale-sulfide textures found in other SEDEX-type deposits (Sangster 2002; Large et al. 2005; Leach et al. 2005; Goodfellow and Lydon 2007; Wilkinson, 2014), textural evidence suggests that the Mehdiabad deposit formed both by deposition on the

832 sea floor and by subsequent further replacement of preexisting sediments and ore minerals.
 833 During the deposition of organic matter-rich shale (the Taft Formation), the water column in the
 834 southern Yazd Basin was likely stratified with deep anoxic waters (Fig. 25A). The regional
 835 anoxic conditions for the basin are indicated by physical evidence, such as well-preserved
 836 laminated sedimentary texture, the presence of organic matter, and the grain size of pyrite
 837 framboids. Early venting of the exhaling hydrothermal fluids in the basin led to the precipitation
 838 of an early assemblage of sulfide minerals in the bedded ore facies as a reaction of the mixing
 839 with seawater (Fig. 25B). Because the ore-forming fluids at Mehdiabad were probably slightly
 840 less dense than seawater (Maghfouri, 2017), they behaved as a buoyant plume (Sato, 1972), at
 841 least initially near and above the fluid vent. Subsequently mixing and cooling occurred, so that
 842 the fluid and the metals precipitating from the vent settled on the seafloor after exhalation (Fig.
 843 25B). The buoyant plume behavior of hydrothermal fluids in the Mehdiabad deposit is consistent
 844 with observations in modern back-arc sediment-covered rifts, such as the Okinawa trough
 845 (Halbach et al. 1991) and with similar ancient massive sulfide deposits, including the Heath
 846 Steele Belt in the Bathurst Mining Camp District (Goodfellow et al. 2003), the Red Dog deposit
 847 (Moore et al., 1986), and the Paleozoic Selwyn Basin deposits in Canada (Goodfellow, 2007).
 848 Fluid mixing with seawater produced cooling and alkalization, resulting in the destabilization
 849 of Pb and Zn chloride complexes and in the precipitation of sulfides. A significant proportion of
 850 reduced sulfur at the Mehdiabad deposit had a biogenic origin, which is supported by: 1) the
 851 broad range of $\delta^{34}\text{S}$ sulfide values, and 2) the textural evidence (framboidal and colloform
 852 textures) for microbial biota associated with sulfide deposition. This early stage of mineralization
 853 is economically insignificant at Mehdiabad, because the hydrothermal fluids, being less dense
 854 than seawater, dispersed metals away from the venting site within the sediments. The fine-

grained nature of the early stage minerals also suggests a rapid deposition (Herzig & Hannington 1995, Kelley et al. 2004b).

Continued up-flow of high-temperature, metal-bearing, dense hydrothermal fluids into and along the bedded ore facies (ore stage 1) and dolomite shows that early sulfide minerals, barite, and dolomitic rocks were later replaced by massive sulfides (ore stage 2) (Fig. 25C). Most of the Mehdiabad sulfide mineralization was deposited during this main ore stage due to subsurface replacement processes in the host rocks (Fig. 25C), due to favorable porosity and permeability conditions. Such a sulfide replacement can be ascribed to either sub-seafloor replacement of carbonate and sulfide lamina and bands, without ascent of fluids onto the seafloor (Fig. 25C), or by sinking of ore fluids into the seafloor sediments from a brine pool in a local depression (Sangster, 2002). In this model, metal-bearing fluids move up along the syn-sedimentary normal fault. Notwithstanding limited leakage of fluid onto the seafloor, these fluids mixed with saline seawater in the unconsolidated sediments and then spread laterally into the host rocks (Fig. 25C). Sulfide minerals formed by replacement of specific sedimentary laminae and bands, where metal chloride complexes in the fluid reacted with H_2S generated by the thermochemical reduction of sulfates at a temperature higher than about 178-238 °C (Maghfouri, 2017).

The ores from the second stage of mineralization (massive-replacement) are overlain by bedded barite (Fig. 12). The vertical transition between the massive-replacement ore facies and the stratiform barite beds is gradual and is characterized by increase of the bedded barite mineralization. These relations shown that brecciation and replacement of dolomite by sulfide minerals predates the closure of the basin. Stratiform barite mineralization forms the top of ore horizon I and conformably covers the sulfide mineralization (Fig. 25D). The barite ore body, which was deposited in oxic seawater, is well exposed in the Mehdiabad area. The transition

from the sulfide to sulfate ore implies major changes in the sedimentary regime, from dominantly anoxic seawater to oxic seawater (Figs. 25C and 25D). This stage of mineralization is characterized by massive to laminated barite and by minor framboidal pyrite, sphalerite, and galena. Barite mineralization in the Mehdiabad deposit formed predominantly on the seafloor (Fig. 18D). SO_4^{2-} required for the formation of barite in this deposit originated from Early Cretaceous seawater sulfate (Maghfouri, 2017) and H_2S requisite for sulfide mineralization was provided by bacterial reduction. Ultimately, the limestone rocks of the Taft Formation covered barite mineralization in the Mehdiabad deposit and the formation of Zn-Pb-Ba mineralization terminated in the ore horizon I (Fig. 25E).

Subsequently the area where the Mehdiabad ore deposit is located experienced inversion tectonics during Tertiary, leading to a modification of the initially Lower Cretaceous geometry. This late stage was accompanied by supergene oxidation and alteration (Reichert et al., 2003; Maghfouri et al., 2017) and a partial remobilization of the metals, especially in the BHO and ER areas. In the CM area, the origin of the upper Ore Horizon (OHII) is not clearly understood yet, as this orebody is completely oxidized.

6- Conclusions

The Mehdiabad deposit was formed in the early stages of a long-lived starved sedimentary basin. In the Mehdiabad sequence >1300 m of sediment accumulated in this basin during Lower Cretaceous. The deposit is hosted within the Taft Formation. The lower unit of the Taft Formation is a rhythmically banded, possibly distal turbiditic calcarenite and shale unit called the Sangestan Formation. The Sangestan Formation is the stratigraphic footwall to mineralization at Mehdiabad. The lower part of the Taft Formation, consisting of organic matter-rich shale,

900 siltstone and dolomite, contains the bulk of the sulfide mineralization. Sediments of the Taft
901 Formation were deposited in a dysoxic to anoxic water column favorable to sulfate-reduction.
902 Bacterial reduction of seawater sulfates provided reduced sulfur for the Mehdiabad sulfides,
903 although partially oxygenated bottom water conditions may have restricted the sulfate-reducing
904 bacteria to the underlying sediments.

905 Mehdiabad is a well-preserved vent-proximal SEDEX deposit consisting of (1) a stockwork
906 of sulfide veins with a hydrothermal alteration halo, discordantly underlying the main orebody;
907 (2) a massive-replacement sulfide orebody containing pyrite, sphalerite, barite and galena with
908 quartz and dolomite; (3) bedded ore that includes laminated sulfides with sedimentary structures;
909 (4) barite ore with minor pyrite, sphalerite and galena (5) distal exhalites made up of banded
910 chert with disseminated barite and minor pyrite. The Mehdiabad deposit has a prominent feeder
911 zone (stockwork) overlain by a well-developed vent complex (massive ore) that laterally gives
912 way to a bedded, stratiform ore body, similar as in the Mansourabad, Sullivan and Tom deposits
913 (Goodfellow and Lydon, 2007; Maghfouri and Hosseinzadeh, 2018). Sulfide minerals were
914 deposited from buoyant hydrothermal fluids vented into an anoxic basin that was affected by
915 synsedimentary faulting and developed in a submarine environment. A thick accumulation of
916 sulfide-bearing barite locally caps the sulfide deposit. The upwardly limiting factor for sulfide
917 precipitation at Mehdiabad, therefore, may not have been a failure at the vent but a change to
918 oxidizing conditions, which favored deposition of barite rather than sulfides.

919 The Mehdiabad deposit formed proximal to the source vent. Most of mineralization is a
920 massive-replacement or breccia textured to indistinctly bedded; facies changes from one ore
921 facies to another occur over short distances. Host-rock replacement textures are common, mainly

in massive ore facies. Mehdiabad deposit formed by a combination of sulfides and barite deposition on the sea floor, and sediment replacement within the sub-seafloor.

7- Acknowledgements

This paper is part of the senior author's PhD research study at University of Tabriz, Tabriz, Iran. The authors thank the University of Tabriz Research Grant Council and the Mehdiabad Mining Complex for funding fieldwork studies and providing access to exploration data and core logs. The paper benefited from detailed comments by Prof. Ebrahim Rastad. The authors thank the board of Mehdiabad Mining Complex (M. Hosseini and A. Safari) and Hajghasemi for allowing access to the deposit and their records. Particular thanks go to H. Pourfaraj for helping the first author during data collection.

References

- Aghanabati, A. (1998) Major sedimentary and structural units of Iran (map): Geosciences, v. 7, p. 29–30.
- Aghanabati, A. (2004) Geology of Iran. Geological Survey and Mineral Exploration of Iran, Tehran, 586 pp.

- 942 Agard, A., Omrani, J., Jolivet, L., Mouthereau, F. (2005) Convergence history across Zagros
943 (Iran): constraints from collisional and earlier deformation: *Int. J. Earth Sci.* v. 94, p. 401–419.
- 944 Alavi, M. 1991, Tectonic map of the Middle East: Tehran, Geological Survey of Iran, scale
945 1:5,000,000.
- 946 Alavi, M. (1994) Tectonics of the Zagros orogenic belt of Iran: New data and interpretations:
947 *Tectonophysics*, v. 229, p. 211–238.
- 948 Alfonso, P., Prol-Ledesma, R.M., Canet, C., Melgarejo, J.C., and Fallick, A.E (2005) Isotopic
949 evidence for biogenic precipitation as a principal mineralization process in coastal
950 gasohydrothermal vents, Punta Mita, Mexico: *Chemical Geology*, v. 224, p. 113–121.
- 951 Aistov, L., Melnikov, B., Krivyakin, B., Morozov, L., Kiristaev, V. (1984) Geology of the Khur
952 area (Central Iran). Explanatory Text of the Khur Quadrangle Map 1:250.000. – V/O
953 Technoexport, Report 20, pp. 1–130.
- 954 Anderson, I.K., Ashton, J.H., Boyce, A.J., Fallick, A.E., Russell, M.J. (1998) Ore depositional
955 processes in the Navan Zn+Pb deposit, Ireland. *ECONOMIC GEOLOGY*, v.93, p. 535–563.
- 956 Anger, G., Nielsen, H., Puchelt, H., Ricke, W. (1966) Sulfur isotopes in the Rammelsberg ore
957 deposit (Germany). *ECONOMIC GEOLOGY*, v. 61, p. 511–536.
- 958 Bagheri, S., Stampfli, G.M. (2008) The Anarak, Jandaq and Posht-e-Badam metamorphic
959 complexes in central Iran: new geological data, relationships and tectonic implications:
960 *Tectonophysics*, v. 451, p. 123–155.
- 961 Barrier, E., Vrielynck, B. (2008a) Map 6: Cenomanian (99.6–93.5 Ma). In: Barrier, E.,
962 Vrielynck, B. (Eds.), *Palaeotectonic Maps of the Middle East – Tectono- Sedimentary-*

- 963 Palinspastic Maps from the Late Norian to Pliocene. Commission for the Geological Map of
964 the World (CGMW/CCGM), Paris.
- 965 Barrier, E., Vrielynck, B. (2008b) Map 5: Middle Aptian (121.0–115.0 Ma). In: Barrier, E.,
966 Vrielynck, B. (Eds.), Palaeotectonic maps of the Middle East – tectonosedimentary–
967 palinspastic maps from the Late Norian to Pliocene. Commission for the Geological Map of
968 the World (CGMW/CCGM), Paris.
- 969 Berberian, M., King, G.C.P. (1981) Towards a palaeogeography and tectonic evolution of Iran:
970 Can. J. Earth Sci, v. 18, p. 210–265.
- 971 Berberian, F., Muir, I.D., Pankhurst, R.J., Berberian, M. (1982) Late Cretaceous and early
972 Miocene Andean-type plutonic activity in northern Makran and Central Iran: J. Geol. Soc.
973 London, v. 139, p. 605–614.
- 974 Berner, R.A, (1969) The synthesis of framboidal pyrite: ECONOMIC GEOLOGY, p. 64, p. 383
975 – 384.
- 976 Betts, P.G., Lister, G.S., O'dea, M.G., (1998) Asymmetric extension of the Middle Proterozoic
977 lithosphere, Mount Isa terrane, Queensland, Australia: Tectonophysics, v. 296, 293–316.
- 978 Betts, P.G., Lister, G.S., (2001) Comparison of the ‘strike-slip’ versus the ‘episodic rift-sag’
979 models for the origin of the Isa Superbasin: Aust. J. Earth Sci., v. 48, 265–280.
- 980 Boni, M., Gilg, H.A., Balassone, G., Schneider, J., Allen, C.R., Moore, F. (2007) Hypogene Zn
981 carbonate ores in the Angouran deposit, NW Iran. Mineralium Deposita: v. 42, p. 799-820.
- 982 Bottrell, S.R., Newton, R.J. (2006) Reconstruction of changes in global sulfur cycling from
983 marine sulfate isotopes: Earth-Science Reviews, v. 75, p. 59– 83.

- 984 Boviri, M., Rastad, E. (2016) Nature and origin of dolomitization associated with sulphide
985 mineralization: new insights from the Tappehsorkh Zn-Pb (-Ag-Ba) deposit, Irankuh Mining
986 District, Iran. Geological Journal. [http://dx.doi.org/ 10.1002/gj.2875](http://dx.doi.org/10.1002/gj.2875).
- 987 Boveiri, M., Rastad, E., Peter, J. (2017) A sub-seafloor hydrothermal syn-sedimentary to early
988 diagenetic origin for the Gushfil Zn-Pb-(Ag-Ba) deposit, south Esfahan, Iran: Neues Jahrbuch
989 für Mineralogie-Abhandlungen, Journal of Mineralogy and Geochemistry, v. 194 (1), p. 61-
990 90.
- 991 Brumsack, H.-J. (1986) The Inorganic Geochemistry of Cretaceous Black Shales (DSDP 41) in
992 Comparison to Modern Upwelling Sediments from the Gulf of California. In: Summerhayes,
993 C.P., Shackelton, N.J. (Eds.), North Atlantic Palaeoceanography, Geological Society, London,
994 Special Publication, v. 21, p. 447–462.
- 995 Bucur, I.I., Senowbari-Daryan, B., Majidifard, M.R. (2003) Neocomian microfossil association
996 from the Taft area near Yazd (Central Iran): Facies, v. 49, p. 217–222.
- 997 Bucur, I.I., Rashidi, K., Senowbari-Daryan, B. (2012) Early Cretaceous calcareous algae from
998 central Iran (Taft Formation, south of Alibad, near Yazd): Facies, v. 58, p. 605–636.
- 999 Buschendorf, F., Nielsen, H., Puchelt, H., Ricke, W. (1963) Schwefelisotopen-untersuchungen
1000 am Pyrit-Sphalerit-Baryt-Lager Meggen/Lenne (Deutschland) und an verschiedenen Devon-
1001 Evaporiten: Geochim Cosmochim Acta, v. 27, p. 501–523.
- 1002 Canet, C., Alfonso, P., Melgarejo, J.C., and Fallick, A.E. (2005) Stable isotope geochemistry of
1003 the Carboniferous Zn-Pb-Cu sediment-hosted sulfide deposits, northeastern Spain:
1004 International Geology Review, v. 47, p. 1298–1315.

- 2521
2522
2523 1005 Cifelli, F., Mattei, M., Rashid, H., Ghalamghash, J. (2013) Right-lateral transpressional tectonics
2524
2525 along the boundary between Lut and Tabas Blocks (Central Iran): *Geophys. J. Int.*, v. 193, p.
2526 1006
2527 1153–1165.
2528 1007
2529
2530 1008 Claypool, G.E., Holser, W.T., Kaplan, I.R., Sakai, H., Zak, I. (1980) The age curves for sulfur
2531
2532 and oxygen isotopes in marine sulfate and their mutual interpretation: *Chemical Geology*, v.
2533 1009
2534 28, p. 199–260.
2535 1010
2536
2537 1011 Cross, M.M., and Bottrell, S.H. (2000) Reconciling experimentally observed sulphur isotope
2538
2539 fractionation during thermochemical sulphate reduction (TSR) with field data: A ‘steadystate’
2540 1012
2541 model of isotopic behaviour: *Goldschmidt, Journal of Conference Abstracts*, v. 5, p. 325.
2542 1013
2543
2544 1014 Daliran, F., Pride, K., Walther, J., Berner, Z.A., Bakker, R.J. (2013) The Angouran Zn (Pb)
2545
2546 deposit, NW Iran: evidence for a two stage, hypogene zinc sulfide zinc carbonates
2547 1015
2548 mineralization: *Ore Geology Reviews*, v. 53, p. 373-402.
2549 1016
2550
2551 1017 Davoudzadeh, M. (1997) Iran. In: Moores, E.M., Fairbridge, R.M. (Eds.), *Encyclopedia of*
2552
2553 *European and Asian Regional Geology*, *Encyclopedia of Earth Sciences Series*. Chapman and
2554 1018
2555 Hall, London, p. 384–405.
2556 1019
2557
2558 1020 Davoudzadeh, M., Soffel, H., Schmidt, K. (1981) On the rotation of the Central-East Iran
2559
2560 microplate: *Neues Jb. Geol. Paläontol. Monat* 5, p. 180–192.
2561 1021
2562
2563 1022 Dercourt, J., Zonenshain, L.P., Ricou, L.-E., Kazmin, V.G., Le Pichon, X., Knipper, A.L.,
2564
2565 Grandjacquet, C., Sbertshikov, I.M., Geyssant, J., Lepvrier, C., Pechersky, D.H., Boulin, J.,
2566 1023
2567 Sibuet, J.-C., Savostin, L.A., Sorokhtin, O., Westphal, M., Bazhenov, M.L., Lauer, J.P., Bijou-
2568 1024
2569
2570
2571
2572
2573
2574
2575
2576

- 2577
2578
2579 1025 Duval, B. (1986) Geological evolution of the Tethys belt from the Atlantic to the Pamir since
2580
2581 1026 the Lias: Tectonophysics, v. 123, p. 241–315.
2582
2583
2584 1027 Ehya, F., Lotfi, M., Rasa, I. (2010) Emarat carbonate-hosted Zn–Pb deposit, Markazi Province,
2585
2586 1028 Iran: A geological, mineralogical and isotopic (S, Pb) study: Journal of Asian Earth Sciences,
2587
2588 1029 v. 37, p. 186–194.
2589
2590
2591 1030 Elliott, H. (2015) Pb-Zn mineralisation within the Limerick Basin (SW Ireland): A role for
2592
2593 1031 volcanism? PhD thesis, National Oceanography Centre, Southampton School of Ocean and
2594
2595 1032 Earth Science. Pp. 197.
2596
2597
2598 1033 Esmaeily, D., Bouchez, J.L., Siqueira, R. (2007) Magnetic fabrics and microstructures of the
2599
2600 1034 Jurassic Shah-Kuh granite pluton (Lut Block, Eastern Iran) and geodynamic inference:
2601
2602 1035 Tectonophysics, v. 439, p. 149–170.
2603
2604
2605 1036 Fallick, A.E., Ashton, J.H., Boyce, A.J., Ellam, R.J., Russell, M.J. (2001) Bacteria were
2606
2607 1037 responsible for the magnitude of the world-class hydrothermal base metal-sulfide orebody at
2608
2609 1038 Navan, Ireland: ECONOMIC GEOLOGY, v. 96, p. 885–890.
2610
2611
2612 1039 Gadd, M.G., Matthews, L.D., Peter, J.M. (2016a) Non-hydrothermal origin of apatite in SEDEX
2613
2614 1040 mineralization and host rocks of the Howard’s Pass district, Yukon, Canada: American
2615
2616 1041 Mineralogist, v. 101, p. 1061–1071.
2617
2618
2619 1042 Gadd, M.G., Matthews, L.D., Peter, J.M., Paradis, S., Jonsson, R.I. (2016b) The world-class
2620
2621 1043 Howard’s Pass SEDEX Zn-Pb district, Selwyn Basin, Yukon. Part II: the roles of
2622
2623 1044 thermochemical and bacterial sulfate reduction in metal fixation. Miner Deposita, Volume 52,
2624
2625 1045 Issue 3, pp 405–419.
2626
2627
2628
2629
2630
2631
2632

- 2633
2634
2635 1046 Ghasemi, M. (2006) Formation Mechanism of the Mehdi Abad Zn-Pb Deposit and its
2636
2637 1047 Comparison with Other Near Lead and Zinc Deposits (Unpublished M.Sc. Thesis): Research
2638
2639
2640 1048 Institute of Earth Sciences, Geological Survey of Iran, p. 238.
2641
2642
2643 1049 Ghazban, F., McNutt, R.H., Schwarcz, H.P. (1994) Genesis of sediment-hosted Zn-Pb-Ba
2644
2645 1050 deposits in the Iran Kouh district, Esfahan area, west-Central Iran: ECONOMIC GEOLOGY,
2646
2647 1051 v. 89, p. 1262–1278.
2648
2649
2650 1052 Gilg, H.A., Boni, M., Balassone, G., Allen, C.R., Banks, D., Moore, F. (2006) Marble hosted
2651
2652 1053 sulphide ores in the Angouran Zn- (Pb-Ag) deposit, NW Iran: interaction of sedimentary
2653
2654 1054 brines with a metamorphic core complex: Mineralium Deposita, v. 41, p. 1-16.
2655
2656
2657 1055 Goodfellow, W.D., Jonasson, I.R. (1987) Environment of formation of the Howards Pass (XY)
2658
2659 1056 Zn-Pb deposits, Selwyn Basin, Yukon, in: Morin, J.A., (eds.), Mineral deposits of Northern
2660
2661 1057 Cordillera: Canadian Institute of Mining and Metallurgy Spec, v. 37, p. 19–50.
2662
2663
2664 1058 Goodfellow, W.D., Mccutcheon, S.R., Peter, J.M. (2003) Massive sulfide deposits of the
2665
2666 1059 Bathurst Mining Camp, New Brunswick, and Northern Maine: introduction and summary of
2667
2668 1060 findings: Econ Geology Monograph, 11:1–1.
2669
2670
2671 1061 Goodfellow, W.D. (2004) Geology, genesis and exploration of SEDEX deposits, with emphasis
2672
2673 1062 on the Selwyn basin, Canada, in Deb, M., Goodfellow, W.D., (eds.), Sediment-hosted lead-
2674
2675 1063 zinc sulphide deposits: Attributes and models of some major deposits of India, Australia and
2676
2677 1064 Canada. Delhi, India, Narosa Publishing House, p. 24–99.
2678
2679
2680 1065 Goodfellow, W.D., and Lydon, J.W. (2007) Sedimentary exhalative (Sedex) deposits, in
2681
2682 1066 Goodfellow, W.D., eds., Mineral deposits of Canada: A synthesis of major deposit types,
2683
2684
2685
2686
2687
2688

2689
2690
2691 1067 district metallogeny, the evolution of geological provinces, and exploration methods:
2692
2693 1068 Geological Association of Canada, Mineral Deposits Division, Special Publication 5, p. 163–
2694
2695 183.
2696 1069
2697
2698 1070 Halbach, P.E., Marten, A., Schwanold, G. (1991) Present-day Kuroko type ore formation: results
2699
2700 from the central Okinawa trough. In: Pagel M, Leroy JL (eds) Source, transport and
2701 1071
2702 deposition of metals: Balkema, Rotterdam, p. 623–626.
2703 1072
2704
2705 1073 Haghipour, A., Pelissier, G. (1970) Geological Quadrangle Map of Iran, H 8, Ardekan
2706
2707 1:250,000. Geological Survey of Iran, Tehran.
2708 1074
2709
2710 1075 Hajmolla Ali, A., Majidifard, M.R. (2000) Geological Map of Yazd, Scale 1:100,000. Geological
2711
2712 Survey of Iran.
2713 1076
2714
2715 1077 Herzig, P.M., Hannington, M.D. (1995) Polymetallic massive sulfides at the modern seafloor, a
2716
2717 review: Ore Geology Review, v. 10, p. 95–115.
2718 1078
2719
2720 1079 Hoffman, D.L., Algeo, T.J., Maynard, J.B., Joachimski, M.M., Hower, J.C., Jaminski, J. (1998)
2721
2722 Regional stratigraphic variation in bottom water anoxia in offshore core shales of Upper
2723 1080
2724 Pennsylvanian cyclothems from Eastern Midcontinent Shelf (Kansas), U.S.A. in: Schieber, J.,
2725 1081
2726 Zimmerle, W., Sethi, P., (eds.), Shales and mudstones I.E Schweizerbart'sche
2727 1082
2728 Verlagsbuchhandlung, Stuttgart, p. 243–269.
2729 1083
2730
2731 1084 Huston, D.L., Power, M., Gemmell, J.B., Large, R.R. (1995) Design, calibration and geological
2732
2733 application of the first operational Australian laser ablation sulphur isotope microprobe: Aust.
2734 1085
2735 J. Earth Sci, v. 42, p. 549–555.
2736 1086
2737
2738
2739
2740
2741
2742
2743
2744

- 1087 Ireland, T. (2001) Lateral Variability in the HYC Zn-Pb-Ag deposit, Northern Territory,
1088 Australia; implications for ore genesis. Unpubl Honours thesis, University of Tasmania,
1089 Australia. pp. 189.
- 1090 Ireland, T., Bull, S.W., Large, R.R. (2004a) Mass flow sedimentology within the HYC Zn-Pb-Ag
1091 deposit, Northern Territory, Australia: Evidence for syn-sedimentary ore genesis: *Mineralium*
1092 *Deposita*, v. 39, p. 143–158.
- 1093 Ireland, T., Large, R.R., McGoldrick, P., Blake, M. (2004b) Spatial distribution patterns of sulfur
1094 isotopes, nodular carbonate, and ore textures in the McArthur River (HYC) Zn-Pb-Ag deposit,
1095 Northern Territory, Australia: *ECONOMIC GEOLOGY*, v. 99, p. 1687-1709.
- 1096 Jones, B., Manning, D.A.C (1994) Comparison of geochemical indices used for the interpretation
1097 of paleoredox conditions in ancient mudstone: *Chemical Geology*, v. 111, p. 111–129.
- 1098 Johnson, C.A., Kelley, K.D., Leach, D.L (2004) Sulfur and oxygen isotopes in barite deposits of
1099 the western Brooks Range, Alaska, and implications for the origin of the Red Dog massive
1100 sulfide deposits: *ECONOMIC GEOLOGY*, v. 99, p. 1435–1448.
- 1101 Kelley, K.D., Dumoulin, J.A., Jennings, S (2004a) The Anarraaq Zn-Pb- Ag and barite deposit,
1102 northern Alaska: Evidence for replacement of carbonate by barite and sulfides: *ECONOMIC*
1103 *GEOLOGY*, v. 99, p. 1577–1591.
- 1104 Kelley, K.D., Leach, D.L., Johnson, C.A., Clark, J.L., Fayek, M., Slack, J.F., Anderson, V.M,
1105 Ayuso, R.A, Ridley, W.I. (2004b) Textural, compositional, and sulfur isotope variations of
1106 sulfide minerals in the Red Dog Zn-Pb-Ag deposits, Brooks Range, Alaska: Implications for
1107 ore formation: *ECONOMIC GEOLOGY*, v. 99, p. 1509–1532.

- 1108 Large, R.R., Bull, S.W., Cooke, D.R., McGoldrick, P.J. (1996) Review of genetic models at
 1109 HYC: Constraints from new sedimentology, alteration halo studies and fluid chemical
 1110 modelling: Townsville, James Cook University of North Queensland: ECONOMIC
 1111 GEOLOGY Research Unit Contribution, v. 55, p. 72–74.
- 1112 Large, R.R., Bull, S.W., Yang, J., Cooke, D.R., Garven, G., McGoldrick, P.J., Selley, D. (2002)
 1113 Controls on the formation of giant stratiform sediment-hosted Zn-Pb-Ag deposits with
 1114 particular reference to the north Australian Proterozoic. University of Tasmania, Centre for
 1115 Special Ore Deposit and Exploration (CODES) Studies Publication 4, p. 107–149
- 1116 Large, R.R., Bull, S.W., McGoldrick, P.J., Walters, S., Derrick, G.M., Carr, G.R. (2005)
 1117 Stratiform and strata-bound Zn-Pb-Ag deposits in Proterozoic sedimentary basins, northern
 1118 Australia. ECONOMIC GEOLOGY 100th Anniversary, p. 931–963
- 1119 Leach, D.L., Sangster, D.F., Kelley, K.D., Large, R.R., Garven, G., Allen, C.R., Gutzmer, J.,
 1120 Walters, S. (2005) Sediment-hosted lead-zinc deposits: a global perspective. ECONOMIC
 1121 GEOLOGY 100th Anniversary, p. 561–607
- 1122 Leach, D.L., Bradley, D.C., Huston, D., Pisarevsky, S.A., Taylor, R.D., Gardoll, S.J (2010)
 1123 Sediment-hosted lead-zinc deposits in Earth history: ECONOMIC GEOLOGY, v. 105, p.
 1124 593–625.
- 1125 Lee, M.J., and Wilkinson, J.J. (2002) Cementation, hydrothermal alteration, and Zn-Pb
 1126 mineralization of carbonate breccias in the Irish Midlands: Textural evidence from the
 1127 Cooleen zone, near Silvermines, County Tipperary: ECONOMIC GEOLOGY, v. 97, p.
 1128 653–662.

- 2857
2858
2859 1129 Liaghat, S., Moore, F., Jami, M. (2000) The Kouh Sourmeh mineralization, a carbonate-hosted
2860
2861 Zn-Pb deposit in the Simply Folded Belt of the Zagros Mountains, SW Iran: Mineralium
2862 1130
2863 Deposita, v. 35, p. 72-78.
2864 1131
2865
2866 1132 Lindenberg, H.G., Görler, K., Ibbeken, H. (1983) Stratigraphy, structure and orogenic
2867
2868 evolution of the Sabzevar Zone in the area of Oryan Khorasan, NE Iran. Geological Survey of
2869 1133
2870 Iran Report, v. 51, p. 120–143.
2871 1134
2872
2873 1135 Logan, R.G. (1979) The Geology and Mineralogical Zoning of the H.Y.C. Ag-Pb-Zn Deposit,
2874
2875 McArthur River, Northern Territory, Australia. Unpublished Thesis. Australian National
2876 1136
2877 University, pp. 341.
2878 1137
2879
2880 1138 Lydon, J.W. (1996) Sedimentary exhalative sulphides (SEDEX), in Eckstrand, O.R., Sinclair,
2881
2882 W.D., Thorpe, R.I., (eds.), Geology of Canadian Mineral Deposit Types, Geology of Canada,
2883 1139
2884 No. 8, Geological Survey of Canada (also Geological Society of America, The Geology of
2885 1140
2886 North America P-1), p. 130-152.
2887 1141
2888
2889 1142 Lydon, W.J. (2004) Geology of the Beit-Purcell Basin and the Sullivan deposit, in Deb, M.
2890
2891 Goodfellow, W.D., (eds.), Sediment-hosted lead-zinc sulphide deposits: Attributes and models
2892 1143
2893 of some major deposits of India, Australia and Canada. Narosa Publishing House, Delhi,
2894 1144
2895 India, p. 100-148.
2896 1145
2897
2898 1146 Maghfouri, S., Hosseinzadeh, M.R., Rajabi, A., Azimzadeh, A.M., Choulet, F. (2015) Geology
2899
2900 and Origin of Mineralization in the Mehdiabad Zn-Pb-Ba (Cu) Deposit, Yazd Block, Central
2901 1147
2902 Iran, 13th SGA biennial meeting, Nancy-France.pp. 3.
2903 1148
2904
2905
2906
2907
2908
2909
2910
2911
2912

- Maghfouri, S., Hosseinzadeh, M.R., Rajabi, A., Azimzadeh, A.M. (2016a) Facies analysis and stratigraphy position of carbonate-clastic hosted Zn-Pb-Ba mineralization horizons in the Early Cretaceous sedimentary sequence, Southern Yazd Basin: Scientific Quarterly Journal, Geosciences, v. 26, p. 102-115.
- Maghfouri, S., Hosseinzadeh, M.R., Rajabi, A., Azimzadeh, A.M (2016b) Darreh-zanjir Deposit, a Typical Carbonate Hosted Zn-Pb Deposit (MVT) in Early Cretaceous Sedimentary Sequence, Southern Yazd Basin, Scientific Quarterly Journal, Geosciences, v. 26, No: 103.
- Maghfouri, S. (2017) Geology, Geochemistry, Ore Controlling Parameters and Genesis of Early Cretaceous Carbonate-clastic Hosted Zn-Pb Deposits in Southern Yazd Basin, with Emphasis on Mehdiabad Deposit (Unpublished Ph.D. Thesis). Tabriz University, Iran, p. 475.
- Maghfouri, S., and Hosseinzadeh, M.R. (2018) The early Cretaceous Mansourabad shale carbonate hosted Zn–Pb (-Ag) deposit, central Iran: An example of vent-proximal sub-seafloor replacement SEDEX mineralization. Ore Geology Reviews, v. 95, p. 20–39.
- Magnall, J.M., Stern, R.A., Gleeson, S.A., Paradis, S. (2015) Widespread euxinic conditions are not a prerequisite for sediment-hosted base metal (Pb-Zn-Ba) mineralization at MacMillan Pass, Yukon. In S. Paradis, Ed., Targeted Geoscience Initiative 4: Sediment-hosted Zn-Pb deposits: Processes and implications for exploration, p. 43–57. Geological Survey of Canada, Open File 7838.
- Magnall, M.J., Gleeson, S.A., Blamey, N.J.F., Paradis, S., Luo, Y. (2016) The thermal and chemical evolution of hydrothermal vent fluids in shale hosted massive sulphide (SHMS) systems from the MacMillan Pass district (Yukon, Canada): Geochimica et Cosmochimica Acta, v. 193, p. 251–273.

- 2969
2970
2971 1171 Majidifard, M.R. (1996) Stratigraphy, fossils and environment of Early Cretaceous rocks from
2972
2973 the northern hills of Shirkuh: Geological Survey of Iran, Earth Science 20, 2e31 (in Farsi).
2974 1172
2975
2976 1173 Majidifard, M.R., Vaziri, S.H. (2000) Geological Map of Bahadoran, Scale 1:100,000.
2977
2978 1174 Geological Survey of Iran.
2979
2980
2981 1175 Mattei, M., Cifelli, F., Muttoni, G., Zanchi, A., Berra, F., Mossavvari, F., Eshraghi, S.A. (2012)
2982
2983 Neogene block rotation in central Iran: evidence from paleomagnetic data: Geol. Soc, Am.
2984 1176
2985 Bull. v. 124, p. 943–956.
2986 1177
2987
2988 1178 Mohseni, M. (2011) Geology and mineralization of the Mehdi Abad Zn-Pb Deposit
2989
2990 (Unpublished M.Sc. Thesis). Dameghan University, Iran, 190 p.
2991 1179
2992
2993 1180 Momenzadeh, M. (1976) Stratabound Lead-Zinc Ores in the Early Cretaceous and Jurassic
2994
2995 Sediments in the Malayer-Esfahan District (West Central Iran), Lithology, Metal Content,
2996 1181
2997 Zonation and Genesis (Unpublished Ph.D. Thesis). University of Heidelberg, Heidelberg, 300
2998 1182
2999 p.
3000 1183
3001
3002 1184 Momenzadeh, M., Shafighi, S., Rastad, E., and Amstutz, G.S. (1979) The Ahangaran lead-silver
3003
3004 deposit, Se-Malayer, west Central Iran: Mineralium Deposita, v. 14, p. 323–341.
3005 1185
3006
3007 1186 Moore, D.E., Young, L.E., Modene, J.S., Plahuta, J.T. (1986) Geologic setting and genesis of the
3008
3009 Red Dog zinc-lead-silver deposit, Western Brooks Range, Alaska: ECONOMIC GEOLOGY,
3010 1187
3011 v. 81, p. 1696–1727.
3012 1188
3013
3014 1189 Muttoni, G., Mattei, M., Balini, M., Zanchi, A., Gaetani, M., Berra, F (2009a) The drift history
3015
3016 of Iran from the Ordovician to the Triassic. In: Brunet, M.-F., Wilmsen, M., Granath, J.W.
3017 1190
3018
3019
3020
3021
3022
3023
3024

- 1191 (Eds.), South Caspian to Central Iran Basins, v. 312. Geological Society London, Special
1192 Publications, p. 7–29.
- 1193 Muttoni, G., Gaetani, M., Kent, D.V., Sciunnach, D., Angiolini, L., Berra, F., Garzanti, E.,
1194 Mattei, M., Zanchi, A (2009b) Opening of the Neo-Tethys ocean and the Pangea B to Pangea
1195 A transformation during the Permian: GeoArabia, v. 14, p. 17–48.
- 1196 Nabavi, M (1972a) Lower Cretaceous deposits in the Taft-Yazd and Khur area. Geological
1197 Survey of Iran, Report 106, p. 1–127.
- 1198 Nabavi, M.H (1972b) Geological quadrangle map of Iran, H 9, Yazd 1:250,000. Geological
1199 Survey of Iran, Tehran.
- 1200 Nath, B. N., Bau, M., Ramlin Geswara Rao, B. & Rao, Ch.M (1997) Trace and rare earth
1201 elemental variation in Arabian Sea sediments through a transect across the oxygen minimum
1202 zone. Geochim: Cosmochim. Acta, v. 61, p. 2375 – 2388.
- 1203 Nosratian, J (1991) Geological Map of Mehdi Abad. Based on the Topographical Map Mehdi
1204 Abad-Yazd, 1:2000.
- 1205 Ohmoto, H., Kaiser, C.J., Geer, A (1990) Systematics of sulphur isotopes in recent marine
1206 sediments and ancient sediment-hosted base metal deposits, in: Herbert, H.K., Ho, S.E., (eds.),
1207 Stable isotopes and fluid processes in mineralization, v. 23, Perth, University of Western
1208 Australia Publishing, p. 70–120.
- 1209 Peter, J.M., Scott, S.D (1999) Windy Craggy, northwestern British Columbia: the world's largest
1210 Besshi deposit. Rev. ECONOMIC GEOLOGY. v. 8, p. 261–295.

- 1211 Pfaff, K., Hildebrandt, H., Leach, D., Jacob, D., Markl, G (2010) Formation of the Wiesloch
Mississippi Valley-type Zn-Pb-Ag deposit in the extensional setting of the Upper
Rhinegraben, SW Germany: *Miner Deposita*, v. 45, p. 647–666
- 1214 Philip, J., Floquet, M (2000) Late Cenomanian. In: Dercourt, J., Gaetani, M., Vrielynck, B.,
Barrier, E., Biju-Duval, B., Brunet, M.F., Cadet, J.P., Crasquin, S., Sandulescu, M. (Eds.),
Atlas Peri-Tethys Palaeogeographical Maps. CCGM/CGMW, Paris, p. 129–136.
- 1217 Pourfaraj, H., Maghfouri, S., Mohajjel, M., Rastad, E (2016) Kinematic change in syn-
sedimentary normal fault in Mehdi-Abad Mine of Yazd, 33rd National Geosciences
Symposium, Trhran.
- 1220 Prol-Ledesma, R.M., Canet, C., Villanueva-Estrada, R.E., and Ortega-Osorio, A (2010)
Morphology of pyrite in particulate matter from shallow submarine hydrothermal vents:
American Mineralogist, v. 95, p. 1500–1507.
- 1223 Quinby-Hunt, M.S., Wilde, P (1994) Thermodynamic zonation in the black shale facies based on
iron–manganese–vanadium content: *Chemical. Geology*, v. 113, p. 297–317.
- 1225 Rajabi, A., Rastad, E., Canet, C (2012) Metallogeny of Cretaceous carbonate hosted Zn–Pb
deposits of Iran: geotectonic setting and data integration for future mineral exploration,
International Geology Review, v. 54:14, p. 1649-1672.
- 1228 Rajabi A, Rastad E, Canet C (2013) Metallogeny of Permian–Triassic carbonate-hosted Zn–Pb
and F deposits of Iran: a review for future mineral exploration: *Aust J Earth Sci*, v. 60, p. 197–
216

- 1231 Rajabi, A., Rastad, E., Canet, C., Alfonso, P (2014) Basin evolution and Stratigraphic Correlation
1232 of Sedimentary-Exhalative Zn-Pb deposits, Zarigan-Chahmir Basin: Ore Geology Reviews, v.
1233 64, p. 328–353.
- 1234 Rasa, I (1987) Geologisch-petrographische untersuchungen in der Blei-lagerstatte Nakhlak,
1235 Zentraliran [Ph.D. thesis]: Heidelberg, Heidelberg University.
- 1236 Rastad, E (1981) Geological, Mineralogical, and Facies Investigations on the Early Cretaceous
1237 Stratabound Zn-Pb-(-Ba-Cu) Deposits of the Iran Kouh Mountain Range, Esfahan, West
1238 Central Iran (Unpublished Ph.D. Thesis). University of Heidelberg, Heidelberg, 334 p.
- 1239 Reed, C., Wallace, M (2004) Zn-Pb mineralisation in the Silvermines district, Ireland: a product
1240 of burial diagenesis: Mineralium Deposita, v. 39, p. 87–102.
- 1241 Reichert, J., Borg, G. & Rashidi, B. (2003) Mineralogy of calamine ore from the Mehdi Abad
1242 zinc-lead deposit, Central Iran. In: Eliopoulos, D.G. et al. (eds.) Mineral Exploration and
1243 Sustainable Development. vol. 1, 97-100, Millpress, Rotterdam.
- 1244 Reichert, J (2007) A Metallogenetic Model for Carbonatehosted Non-sulfide Zinc Deposits
1245 Based on Observations of Mehdi Abad and Iran Kouh, Central and Southwestern Iran
1246 (Unpublished Ph.D. Thesis). University of Martin Luther, Shillong, 129 p.
- 1247 Rossetti, F., Nasrabad, M., Vignaroli, G., Theye, T., Gerdes, A., Razavi, M., Moin Vaziri, H
1248 (2010) Early Cretaceous migmatitic mafic granulites from the Sabzevar range (NE Iran):
1249 implications for the closure of the Mesozoic peri-Tethyan oceans in central Iran: Terra Nova,
1250 v. 22, p. 26–34.

- 3193
3194
3195 1251 Sáez, R., Moreno, C., González, F., Almodóvar, G.R (2011) Black shales and massive sulfide
3196
3197 deposits: causal or casual relationships? Insights from Rammelsberg, Tharsis, and Draa Sfar:
3198 1252
3199 Mineralium Deposita, v. 46, no. 5–6, p. 585–614.
3200 1253
3201
3202 1254 Sangster, D.F (2002) The role of dense brines in the formation of vent-distal sedimentary
3203
3204 exhalative (SEDEX) lead-zinc deposits: field and laboratory evidence: Mineralium Deposita,
3205 1255
3206 v. 37, p. 149-157.
3207 1256
3208
3209 1257 Sato, T (1972) Behaviours of ore-forming solutions in seawater: Mining Geology, v. 22, p. 31–
3210
3211 42.
3212 1258
3213
3214 1259 Schlagintweit, F., Bucur, I.I., Rashidi, K., Hanifzadeh, R., Wilmsen, M (2013a) Torremiroella
3215
3216 hispanica Brun and Canérot, 1979 (benthic foraminifera) from the Lower Cretaceous of
3217 1260
3218 Central Iran and its palaeobiogeographic significance: Cretaceous. Research, v. 46, p. 272–279.
3219 1261
3220
3221 1262 Schlagintweit, F., Bucur, I.I., Rashidi, K., Saberzadeh, B (2013b) Praeorbitolina claveli n. sp.
3222
3223 (benthic foraminifera) from the Lower Aptian (Bedoulian) of Central Iran: Carnets de
3224 1263
3225 Géologie/Notebooks on Geology, v. 4, p. 255–272.
3226 1264
3227
3228 1265 Seyed-Emami, K., Bozorgnia, F., Eftekhari-Nezhad, J (1972) Der erste sichere Nachweis von
3229
3230 Valanginien im nordöstlichen Zentraliran (Sabzewar-Gebiet): Neues Jb. Geol. Paläontol.
3231 1266
3232 Monat, v.1, p. 52–67.
3233 1267
3234
3235 1268 Shanks, W.C., Woodruff, L.G., Jilson, G.A., Jennings, D.S., Modene, J.S., Ryan, B.D (1987)
3236
3237 Sulfur and lead isotope studies of stratiform Zn-Pb-Ag deposits, Anvil Range, Yukon: basinal
3238 1269
3239 brine exhalation and anoxic bottom-water mixing: ECONOMIC GEOLOGY, v. 82, p. 600–
3240 1270
3241 634.
3242 1271
3243
3244
3245
3246
3247
3248

3249
3250
3251 1272 Soffel, H., Davoudzadeh, M., Rolf, C., Schmidt, S (1996) New palaeomagnetic data from
3252
3253 Central Iran and a Triassic palaeo-reconstruction: *Geol. Rundsch*, v. 85, p. 293–302.
3254 1273
3255
3256 1274 Southam, G., Saunders, J.A (2005) The geomicrobiology of ore deposits: *ECONOMIC*
3257
3258 *GEOLOGY*, v. 100, p. 1067–1084.
3259 1275
3260
3261 1276 Stampfli, G.M., Borel, G.D (2002) A plate tectonic model for the Paleozoic and Mesozoic
3262
3263 constrained by dynamic plate boundaries and restored synthetic oceanic isochrones: *Earth*
3264 1277
3265 *Planet. Sci*, v. 196, p. 17–33.
3266 1278
3267
3268 1279 Takin, M (1972) Iranian geology and continental drift in the Middle East: *Nature*, v. 235, p. 147–
3269
3270 150.
3271 1280
3272
3273 1281 Taraz, H., Aghanabati, A (1983) Geological Quadrangle Map of Iran, G 9, Abadeh 1:250,000.
3274
3275 Geological Survey of Iran, Tehran.
3276 1282
3277
3278 1283 Tirrul, R., Bell, I.R., Griffis, R.J., Camp, V.E (1983) The Sistan suture zone of eastern Iran.
3279
3280 *Geol. Soc. Am. Bull*, v. 94, p. 134–150.
3281 1284
3282
3283 1285 Velasco, F., Herrero, J.M., Gil, P.P., Alvarez, L., Yusta, I (1994) Mississippi Valley-type,
3284
3285 SEDEX, and iron deposits in Lower Cretaceous rocks of the Basque-Cantabrian Basin,
3286 1286
3287 northern Spain. In: Fontbote, L., Boni, M., (eds.), *Sediment-hosted Zn-Pb Ores*. Society for
3288 1287
3289 *Geology Applied to Ore Deposits*, Springer-Verlag, Berlin-Heidelberg, v. 10, p. 246–270.
3290 1288
3291
3292 1289 Wilkinson JJ (2003) On diagenesis, dolomitisation and mineralisation in the Irish Zn-Pb orefield:
3293
3294 *Mineralium Deposita*, v. 38, p. 968–983.
3295 1290
3296
3297
3298
3299
3300
3301
3302
3303
3304

- 3305
3306
3307 1291 Wilkinson JJ, Eyre, S.L., Boyce, A.J. (2005) Ore-forming processes in Irish-type carbonate-
3308
3309 hosted Zn-Pb deposits: Evidence from mineralogy, chemistry and isotopic composition of
3310 1292
3311 sulfides at the Lisheen Mine: ECONOMIC GEOLOGY, v. 100, p. 63–86.
3312 1293
3313
3314 1294 Wilkinson, J.J. (2014) Sediment-Hosted Zinc–Lead Mineralization: Processes and Perspectives,
3315
3316 Treatise on Geochemistry 2nd Edition.
3317 1295
3318
3319 1296 Wilmsen, M., Fürsich, F.T., Seyed-Emami, K. (2003) Revised lithostratigraphy of the Middle
3320
3321 and Upper Jurassic Magu Group of the northern Tabas Block, Eastcentral Iran: Newsl.
3322 1297
3323 Stratigr, v. 39, p. 143–156.
3324 1298
3325
3326 1299 Wilmsen, M., Wiese, F., Seyed-Emami, K., Fürsich, F.T. (2005) First record and significance of
3327
3328 Cretaceous (Turonian) ammonites from the Shotori Mountains, east-central Iran: Cretaceous
3329 1300
3330 Research, v. 26, p. 181–195.
3331 1301
3332
3333 1302 Wilmsen, M., Fürsich, F.T., Seyed-Emami, K., Majidifard, M.R., Taheri, J (2009) The
3334
3335 Cimmerian orogeny in northern Iran: tectono-stratigraphic evidence from the foreland: Terra
3336 1303
3337 Nova, v. 21, p. 211–218.
3338 1304
3339
3340 1305 Wilmsen, M., Fürsich, F.T., Majidifard, J (2013) The Shah Kuh Formation, a latest Barremian –
3341
3342 early Aptian carbonate platform of Central Iran (Khur area, Yazd Block). Cretaceous
3343 1306
3344 Research, v. 39, p. 183–194.
3345 1307
3346
3347 1308 Wilmsen, M., Fürsich, F. T., Majidifard, M. R (2014) An overview of the Cretaceous
3348
3349 stratigraphy and facies development of the Yazd Block, western Central Iran, Journal of Asian
3350 1309
3351 Earth Sciences.pp. 18.
3352 1310
3353
3354
3355
3356
3357
3358
3359
3360

3361
3362
3363 1311 Whitehead, R.E. (1973) Environment of stratiform sulphide deposition; variation in Mn:Fe ratio
3364
3365 1312 in host rocks at Heath Steele Mine, New Brunswick, Canada: Mineral. Deposita, v. 8, p. 148–
3366
3367 160.
3368 1313
3369
3370 1314 Whittaker, S.G., Kyser, T.K. (1993) Variations in the neodymium and strontium isotopic
3371
3372 composition and REE content of molluscan shells from the Cretaceous Western Interior
3373 1315
3374 Seaway: Geochim. Cosmochim. Acta, v. 57, p. 4003 – 4014.
3375 1316
3376
3377 1317 Wingal, P.B., Twitchett, R.J. (1996) Oceanic anoxia and the end-Permian mass extinction:
3378
3379 Science, v. 272, p. 1155 –1158.
3380 1318
3381
3382 1319 Yaghubpur, A., Mehrabi, B. (1997) Kushk zinc-lead deposit: a typical black-shale-hosted deposit
3383
3384 in Yazd State, Iran: Journal of Sciences, Islamic Republic of Iran, v. 8, no. 2, p. 117–125.
3385 1320
3386
3387 1321 Yarmohammadi, A., Rastad, E., Rajabi, A. (2016) Geochemistry, fluid inclusion study and
3388
3389 genesis of the sediment-hosted Zn-Pb ($\pm$ Ag $\pm$ Cu) deposits of the Tiran basin, NW of
3390 1322
3391 Esfahan, Iran: N. Jb. Miner. Abh. (J. Min. Geochem.), v. 193/2, p. 183–203.
3392 1323
3393
3394
3395 1324
3396
3397
3398 1325
3399
3400
3401 1326
3402
3403
3404 1327
3405
3406
3407 1328
3408
3409 1329
3410
3411
3412
3413
3414
3415
3416

3417
3418
3419
3420
3421
3422
3423
3424
3425
3426
3427
3428
3429
3430
3431
3432
3433
3434
3435
3436
3437
3438
3439
3440
3441
3442
3443
3444
3445
3446
3447
3448
3449
3450
3451
3452
3453
3454
3455
3456
3457
3458
3459
3460
3461
3462
3463
3464
3465
3466
3467
3468
3469
3470
3471
3472

Table captions

Table 1. Major (%) and trace elements (ppm) contents of the Taft Formation shales hosting the mineralization of the Mehdiabad deposit. (BH: Black-Hill ore, CVOB: Central Valley Orebody, ER: East Ridge ore).

Table 2. Values of $\delta^{34}\text{S}_{\text{SDT}}$ in sulfide minerals and barite from the Mehdiabad deposit

Table 3. Assay analysis of ore minerals in the different drill holes of Mehdiabad deposits (Location of samples shown in the figure 12).

Figure captions

Fig.1. Distribution map of sediment hosted Zn-Pb deposits in the main tectonic units of Iran; Al: Alborz zone; CIGS: Central Iranian geological and structural gradual zone; E: East Iran ranges; K: Kopeh-Dagh; KR: Kermanshah Radiolarites subzone; KT: Khazar-Talesh-Ziveh structural zone; L: Lut Block; M: Makran zone; Oph: ophiolite belts; PB: Posht-e-Badam Block; SSZ: Sanandaj-Sirjan zone; T: Tabas Block; TM: tertiary magmatic rocks; UDMA: Urumieh-Dokhtar magmatic arc; Y: Yazd Block; Z: Zabol area; Za: Zagros ranges. Tectonic and structural map of Iran modified after Aghanabati (1998; 2004) and Alavi (1991).

Fig. 2. Simplified geological map of the YAMB in the CIEM, showing the location of the different types of ore deposits. Blue rectangle shows the location of the Southern Yazd Basin in the Fig. 3. (NDBF: Naein-Dehshir-Baft fault; ANF: Anar fault; PBF: Posht-e-Badam fault; CHF:

Chapedoni fault; BF: Biabanak fault; GKDF: Great Kavir Drounaeh fault) (modified after Haghypour and Pelissier, 1970; Nabavi, 1972b; Taraz and Aghanabati, 1983).

Fig. 3. Simplified geological map of the southern Yazd Basin, showing the location of Zn-Pb deposits within this basin (modified after Majidifard and Vaziri, 2000; Hajmolla Ali and Majidifard, 2000). Small rectangle shows the location of study area in the Fig. 4.

Fig. 4. Geological map of the Mehdiabad deposit showing the location of the barite deposit in the east of Black-Hill Fault and of the oxide Zn-Pb orebody in the northeast of the map. CVF: Central Valley fault; NTF: North Thrust fault; FF: Fourozandeh fault; BHO: Black-Hill ore; CVOB: Central Valley orebody; ER: East Ridge; CM: Calamine Mine (A-A', B-B' and C-C' sections are shown in Figs. 8 and 12) (modified after Nosratian, 1991).

Fig. 5. (A) Schematic columnar section of the Lower Cretaceous sedimentary sequence hosting the Mehdiabad deposit. The lower unit of the Taft Formation and the middle unit of the Abkuh Formation host the Zn-Pb-Ba mineralization. (B): General view of the Shir-Kuh granite, unconformably covered by the Sangestan Formation (view to the northeast). (C): View of the silicified silty limestone (distal facies) marking the limit between the Sangestan and Taft Formations (view to the northwest). (D): Hand specimen photograph of silty limestone with interbedded organic matter-rich shale host rock in the Mehdiabad deposit.

Fig. 6. (A): General view of the two ore horizons (OH I and OH II) with different ore facies, OH I is located within the Taft Formation, whereas OH II corresponding to the CM orebody that occurs within the Abkuh Formation. Mineralization in the Mehdiabad deposit occurs to the east of the Black-Hill Fault. Numbers show the location of cross-sections in the Fig. 7. (1 to 5 numbers of columnar sections and A-A', B-B', C-C' and D-D' sections are shown in Figs. 7, 8

and 12) (view to the north-northwest). (B): The location of the bedded ore facies (ER) and distal facies that are embedded with Sangestan and Taft Formations (view to the northwest).

Fig. 7. Lithostratigraphic correlation diagram of the Early Cretaceous sedimentary sequence in the Mehdiabad deposit from west (1) to east (5) showing the position of the ore horizons in the stratigraphic units. Numbers show the stratigraphical location of cross-sections in the Fig. 6A.

Fig. 8. Cross-sections and plan view of the ore sequence and extension of mineralization in the Mehdiabad deposit. (A): North-south cross section of the ore sequence with extension of sulfide and non-sulfide ore mineralization's in the CVOB. (B): Schematic cross section of the OH I in the Mehdiabad deposit, which include the BH orebody and ER mineralization. (C): Southeast-northwest section through the Mehdiabad deposit illustrating the geology, ore horizon, and structure of the area. The location of these sections is shown in Figs. 4 and 6A.

Fig. 9. Hand specimen (A, B, C, D and E) and microscopic (F) photographs of ore-bearing host rocks in the ore horizon I. (A): Drill core of organic matter-rich (Om) shale-siltstone (lower Unite) from the Taft Formation overlying the massive ore facies. (B): Hand specimen photograph of organic matter-rich (Om) shale with organic matter nodule (lower Unit of Taft Formation). (C): Alternation of silty limestone laminae (light-gray) and sphalerite (Sph)+galena (Gn)+pyrite (Py) sulfides (black gray) laminas. (D): Hand specimen photograph of shale. Gray laminas are shale and siltstone rich in pyrite (Py), whereas black laminas are rich in organic matter. (E): Typical laminated siltstone and silty sandstone hosting the ore horizon I. Note the precedence of normal micro-fractures, locally inversed in the lower part of specimen. (F): Silty shale laminae composed of feldspar, quartz, sericite, and clay minerals alternating with black organic matter (Om) laminae.

Fig. 10. Histograms showing the distribution of Fe and Mn contents of unaltered shale samples from the Taft Formation. These diagrams are used to discriminate the paleoenvironment and fields are taken from Quinby-Hunt and Wilde (1994). (A): the Mn content indicates the anoxic paleoconditions during the sedimentation of shale. (B): the Fe content is related to both anoxic sulfate reducing and nitrate-reducing conditions.

Fig. 11. Bivariate V/(V+Ni) vs. V/Cr plot indicating geochemical redox conditions at the time of the black shale deposition (modified after Sáez et al. 2011). Most samples plot in the anoxic conditions field. Field boundaries are from Jones and Manning (1994) and Hoffman et al. (1998).

Fig. 12. W-NE cross-section of the ore sequence and extension of ore facies in the Mehdiabad deposit (Maghfouri, 2017). The mineralization is located to the east of the Black-Hill syn-sedimentary fault. Location of this section is shown in Fig. 6A.

Fig. 13. Outcrop, hand specimen and microscopic photographs of main generations of veins and veinlets in the feeder zone of the Mehdiabad deposit.

Fig. 14. Outcrop (A), hand specimen (B, C, D, E, I and J) and microscopic (F, G and H) photographs of the massive-replacement ore facies in the Mehdiabad deposit. (A): General view of the relations between the massive-replacement ore facies and the barite ore. The massive oxide ore facies is capped by the barite ore mineralization. (B): Massive pyrite and sphalerite, showing pyrite (Py) replaced by sphalerite (Sph). (C): Barite (Ba) replaced by galena (Gn). (D): Massive sulfide ore facies consisting of sphalerite (Sph) and barite (Ba) minerals. (E): Dolomite host rock brecciated and replaced by galena (Gn) and sphalerite (Sph). (F): Early pyrite (Py1) replaced by secondary sphalerite (Sph2) and galena (Gn2). (G): Massive-replacement ore facies composed of galena (Gn2), sphalerite (Sph2), barite (Ba) and chalcopyrite (Cpy1). (H):

Replacement of early colloform pyrite (Py1) by late sphalerite (Sph2) and galena (Gn2). (I and J): Hand specimen photographs of massive-replacement sphalerite (Sph), galena (Gn) and pyrite (Py) sulfide mineralization in the lower unit of the Taft Formation.

Fig. 15. Geology map of the ER area showing the location of bedded ore and the distal facies at the boundary between the Taft and the Sangestan Formations.

Fig. 16. Outcrop (A and B), hand specimen (C, D, E and F) and microscopic (G) photographs of the bedded ore facies and distal facies in the Mehdiabad deposit in the lower Unit of the Taft Formation. (A and B): General view of the alternation of limy siltstone (light) and oxide Zn-Pb (black) layers in the ER orebody. (C): Photograph of typical pyritic (Py) laminae in the organic matter (Om) rich shale of the lower unit. The organic matter occurs as laminae and/or lenses within the sulfide laminae. (D): Inter-ore breccia resting on laminated sulfide-bearing shales. (Sphalerite (Sph), galena (Gn) and pyrite (Py)) laminae associated with black organic matter (Om) rich shale of lower Unit in the Taft Formation. (E): Alternation of shale (black-gray) and sulfides (light gray) laminae, including galena (Gn) and sphalerite (Sph) in the lower unit of the Taft Formation. (F): Laminated limy siltstone showing the three main types of laminae; organic matter-rich laminae (Type 1), siltstone laminae (Type 2), and pyritic rich laminae (Type 3). (G): Laminated sulfides including galena (Gn) and sphalerite (Sph1) from the bedded ore facies. (H): Photograph of alternation of chert laminae and Fe-oxide laminae in the distal facies of the Mehdiabad deposit. (I): Hand specimen photograph of chert laminae and Fe-oxide lamina in the diastal facies. (J): Reflected light image of dissaminated pyrite (Py) in the distal facies of the Mehdiabad deposit.

Fig. 17. Photographs (A, B, C, D and E) and photomicrographs (E, F, G and H) of sulfide-sediment textures in Mehdiabad deposit from ore horizon I in the lower unit of the Taft Formation. (A): Photograph of typical pyritic (Py) laminae associated with black organic matter (Om) rich shale of the lower unit. (B): Alternation of organic-matter (Om) laminae (black-gray) and siltstone host rock layers (light gray). (C): Alternation of shale laminae (black gray) and sulfides (sphalerite (Sph), galena (Gn) and pyrite (Py)) lamianas. (D): Sphalerite (Sph), galena (Gn) and pyrite (Py) sulfide layers assocaited with the lower unit black shale host rock. The sulfides occur as laminae and/or lenses within sulfide layer. (E): Siltstone laminae alternating with carbonate laminae. (F): Reflected light image of early colloform pyrite (Py1) crosscut by late sphalerite (Sph2) and galena (Gn2). (G): Early framboidal pyrite from the organic matter-rich shale. (H): Photomicrograph of laminated galena (Gn1) within the bedded ore facies in the lower unit of the Taft Formation.

Fig. 18. Photographs of sulfide-matrix breccia and sulfide-sediment textures in the Mehdiabad deposit from ore horizon I in the lower unit of the Taft Formation. (A and B): Sulfide-matrix breccia and Rounded-subrounded clasts of pyrite (Py) and galena (Gn) in a turbidite bed. In the Fig.A matrix is sphalerite (Sph), galena (Gn), pyrite (Py) and host rock and fragments of breccia is silty limestone and siltstone of the lower unit. In the Fig. B fragments and matrix both are pyrite (Py), galena (Gn) and sphalerite (Sph). (C): Photograph of sulfide-matrix breccia in the lower unit of the Taft Formation. (D): Load-cast texture, with a dolomite clast set on the thin-bedded shale. (E and F): Coarse-grained clasts, penetratating the underlying siltstone and shale bands. (G): Poorly sorted sulfide breccia layer with clasts of pyrite (Py), sphalerite (Sph) and galena (Gn). These layers are parallel to bedding.

Fig. 19. Textures of stage 2 (sub-seafloor replacement) sulfide mineralization in the ore horizon I. (A): Stage 1 (laminated sulfide and barite) crosscut and brecciated by stage 2 sulfide mineralization. Clasts of stage 1 relict within stage 2 sulfide matrix. (B and C): Stage 2 sulfides preferentially replacing matrix of dolomite breccia. (D): Replacement of dolomite (Dol) with stage 2 sphalerite (Sph) minerals is shown.

Fig. 20. Localization of the different mass flows textures, within the Mehdiabad deposit. Grain size of mass flows decreasing away from the BH syn-sedimentary fault. Sph: sphalerite; Gn: galena; Py: pyrite; Ba: barite; Om: organic matter. (See Fig. 12 for cross-section legend)

Fig. 21. (A): Histogram of the $\delta^{34}\text{S}$ values in the different ore facies of the Mehdiabad deposit . (B): Histogram of the $\delta^{34}\text{S}$ values for sulfide minerals and ore-associated barite from the Mehdiabad deposit. (C): Comparison between the $\delta^{34}\text{S}$ values of Mehdiabad sulfides and sulfates with other Zn-Pb deposits. References: Anvil District (Dy, Grum an Faro deposits), Canada (Shanks et al. 1987); Howards Pass, Canada (Goodfellow and Jonasson 1987); HYC, Australia (Ireland, 2001); Meggen, Germany (Buschendorf et al. 1963); Mt Isa, Australia (Large et al. 2005); Rammelsberg, Germany (Anger et al. 1966); Reocin and Troya, Spain (Velasco et al. 1994); Sullivan, Canada (Lydon, 2004); Tom and Jason, Canada (Goodfellow, 2004); Century, Australia (Wilkinson, 2014); George Fisher, Australia (Wilkinson, 2014); Navan and Silvermine, Ireland (Wilkinson, 2014); Red Dog, Canada (Wilkinson, 2014; Kelley et al., 2004a,b). Sulfur isotope composition of seawater sulfates is from Bottrell and Newton (2006).

Fig. 22. Paragenetic sequence of mineralization stages in the Mehdiabad deposit.

Fig. 23. (A): The six drill holes within the Sangestan, Taft and Abkuh Formations showing the sites of samples in the figure 24 and the boundaries of mineralization, BH syndimentary fault,

and different ore facies. (B): General view of the Sangestan Formation, conformably covered by Lower Unit of Taft Formation in the Farahabad area (view to the west). Lower Unit of Taft Formation is host rocks of main mineralization in the Mehdiabad deposit.

Fig. 24. Core photographs of Lower Unit of Taft Formation showing relation between ore minerals and rock forming minerals (location of samples shown in the figure 23A).

Fig. 25. Schematic diagram showing the inferred depositional environment of the Mehdiabad deposit. (A): Deposition of organic matter-rich shale (the Taft Formation) in the anoxic waters column. (B): Early venting of the exhaling hydrothermal fluid on the seafloor would have led to the precipitation of an early assemblage of the fine grain size minerals (stage 1) in the bedded ore facies. The temperature and salinity of hydrothermal fluids of this stage is 141 °C and 8.7% NaCl (Maghfouri, 2017). (C): Continued up-flow of high-temperature, metal-bearing, dense hydrothermal fluid into and along the bedded ore facies (ore stage I) and dolomite at a sub-sea floor level, show that the ore stage I minerals and dolomitic rocks were replaced by massive sulfide (ore stage II), normally towards their base. The temperature and salinity of hydrothermal fluids of this stage is 178-238 °C and 10.1% NaCl (Maghfouri, 2017). (D): This stage of mineralization consists of the precipitation of barite and minor sulfide minerals at and on the sea floor. Barite mineralization occurs from an oxic seawater column. (E): The upper units of the Taft Formation (limestone rocks) received the deposition of barite mineralization and finally the formation of Zn-Pb-Ba mineralization ended in the ore horizon I.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

Massive ore facies

Massive, replacement, brecciated, colloform, vein-veinlet

Chalcopyrite, galena, sphalerite, pyrite, barite, dolomite, calcite, minor quartz

Bedded ore facies

Laminated, disseminated, framboidal

Sphalerite, galena, pyrite, Fe-oxide, minor barite, dolomite, calcite, Mn-minerals

1 Vein-veinlet 1: chalcopryite,
2 quartz and
3 barite
4
5
6
7
8

9
10 Vein-veinlet 2: sphalerite,
11 galena,
12 pyrite and
13 barite
14
15
16
17
18

19
20 Vein-veinlet 3: sphalerite
21 and galena
22
23
24
25
26

27
28 Vein-veinlet 4: barite-rich
29 vein-veinlet's
30
31
32
33
34
35
36
37

38
39 Vein-veinlet 5: pyrite-bearing
40 calcite
41 vein-veinlet's
42
43
44
45
46
47
48
49
50
51
52
53
54
55

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

Reducing the grain size of mass flows with distance from the BH synsedimentary fault

Minerala		Mineralization			Post-mineralization		
		Stage I	Stage II	Stage III	(weathering/supergene)		
Pyrite I	Bedded ore (and distal facies)		Framboidal, colloform, laminated, disseminated				
Sphalerite I							
Galena I							
Quartz (chert)		Banded					
Barite I		Disseminated, banded					
Chalcopyrite I		Disseminated					
Dolomite (Db)		Replacement					
Pyrite II		Coarse grained, massive, vein					
Sphalerite II		Massive, vein, replacement					
Galena II		Massive, vein, replacement					
Chalcopyrite II		Vein, replacement					
Tetrahedrite		Inclusion					
Dolomite (Dc)		Replacement					
Barite II		Massive, vein, replacement					
Quartz (alteration)	Feeder zone and massive- replacement ore facies	Alteration					
Barite III		Banded barite cap				Banded, disseminated	
Pyrite III							
Sphalerite III							
Galena III							
Smithsonite							
Hemimorphite							
Hydrozincite							
Cerussite							
Anglesite							
Malachite							
Azurite							
Fe-oxide/hydroxides

Sample	SiO ₂	Al ₂ O ₃	Fe ₂ O ₃	MgO	CaO	Na ₂ O	K ₂ O	TiO ₂	P ₂ O ₅	MnO	LOI	Total	Ba	Ni
1	73.86	9.53	3.95	3.27	1.73	0.02	0.1	0.03	bdl	0.48	6.01	98.98	530	3.2
2	79.20	6.01	2.12	1.37	0.21	0.19	1.32	0.07	0.03	0.03	7.8	98.35	103	5.8
3	63.25	14.81	3.8	3.17	2.63	0.03	0.19	0.04	bdl	0.62	9.9	98.44	230	3.9
4	66.51	13.14	4.02	3.05	0.63	0.44	2.12	0.45	0.14	0.22	4.9	95.62	532	21.03
5	64.2	12.66	5.12	2.22	3.82	0.12	2.93	0.58	0.1	0.32	6.09	98.16	190	18.74
6	74.4	11.95	2.8	2.59	0.64	0.03	0.17	0.03	0.02	0.06	7.05	99.74	52	6.8
7	77.39	6.1	5.98	4.63	1.41	0.04	0.45	0.07	0.01	0.42	2.03	98.53	45	4.2
8	71.03	12.6	4.03	2.9	0.8	0.04	1.72	0.17	bdl	0.70	4.9	98.89	605	6.08
9	73.90	10.46	4.92	2.28	0.49	0.08	1.06	0.18	0.04	0.28	6.1	99.79	524	9.5
10	81.97	4	4.86	1.33	0.65	0.02	0.12	0.01	bdl	0.69	5.8	99.45	300	3.4
11	77.4	7.2	4.5	4.12	1.82	0.05	0.21	0.04	bdl	0.4	3.0	98.74	210	3.9
12	82.83	5.36	2.8	1.29	0.59	0.02	0.08	bdl	bdl	0.28	5.5	98.75	510	5.03
13	74.49	12.07	5.07	2.01	0.33	0.05	1.76	0.37	0.09	0.83	2.2	99.27	73	11.08
Sample	Co	Cs	Cr	Ga	Hf	Nb	Rb	Sr	Ta	Th	Tl	U	V	W
1	14.1	0.4	10	5.7	bdl	0.8	3.9	132	0.1	1.1	1.9	3.02	15	5.01
2	12.5	5.3	18	9.8	1.4	3.2	50.6	99	0.2	2.8	4.8	19	34	2.4
3	5.4	1.3	12	9.6	bdl	0.8	7.9	15.7	0.3	0.6	0.5	2.7	15	3.08
4	26.9	7.2	19	10	4.9	9.1	91.9	140.6	2	1.01	0.3	3.7	43	7.14
5	34.2	9.3	14	13.1	5.3	12.1	113	61.5	0.9	0.9	0.9	1.02	62.03	3.9
6	14.6	0.7	16	7.2	bdl	1	6.9	111	0.2	0.7	5.2	2.6	15	2.15
7	10.1	2.1	16	10.5	0.6	1.3	18.9	12.9	0.5	2.3	0.6	1.5	19	5.32
8	9.7	2.4	16	9.9	0.8	2.6	28.1	186.8	0.2	1.6	0.7	3.7	37	5.18
9	13.4	4	19	12.5	1.1	3.7	49	145	0.3	2.9	1.1	4.9	43	3.4
10	8	0.5	14	7.8	bdl	bdl	5	12.6	0.3	0.19	0.8	1	36	1.02
11	5.4	1	15	8.9	bdl	0.8	7.7	15.7	0.3	0.52	0.4	1.9	9	bdl
12	11.3	0.2	15	8.8	bdl	bdl	2.9	15.7	bdl	0.5	1.4	2.4	43	2.37
13	24.9	4.4	14	12.4	2.5	8.4	94.5	203	1.5	4.9	0.4	4.1	58	4.12
Sample	Zr	Y	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm
1	9.1	2.9	15.6	22.4	2.11	9.5	1.8	bdl	1.37	0.14	0.64	0.08	0.18	bdl
2	45.2	5.4	17.6	25.8	2.56	10.3	1.9	bdl	1.67	0.16	1.07	0.13	0.48	0.1
3	9.4	5	6.9	11.7	1.36	7.1	1.8	0.01	2.11	0.21	0.95	0.13	0.33	bdl
4	156.3	17.5	25.8	46.1	4.99	20.3	4.3	bdl	3.56	0.56	3.23	0.79	1.89	0.3
5	123.1	22.1	12.3	52.4	5.20	24.6	4.1	0.01	4.11	0.74	3.03	0.75	2.37	0.3
6	7.8	3.4	5.8	9.3	0.20	3.6	1	0.1	0.89	0.12	0.59	0.11	0.27	bdl
7	17.9	6.2	9.7	16.1	1.64	7.7	1.6	bdl	1.78	0.23	1.19	0.17	0.39	0.1
8	28.7	4.9	21.3	30.3	2.73	10.7	1.7	bdl	1.39	0.18	0.39	0.13	0.44	0.1
9	43.2	5.2	8.1	10.9	0.98	2.7	0.5	0.02	0.75	0.18	0.64	0.16	0.64	0.1
10	3.3	2.7	3	6	0.59	3.3	0.8	bdl	0.76	0.1	0.37	0.06	0.15	bdl
11	9.4	4.8	6.6	12	1.27	6.1	1.6	bdl	1.88	0.17	0.87	0.12	0.33	bdl
12	2.9	2.3	2.5	4.8	0.63	2.6	0.6	bdl	0.65	0.08	0.45	0.06	0.13	bdl
13	73	11.2	22.7	34.2	3.6	12.4	2	bdl	2.53	0.33	2.25	0.38	1.32	0.2
Sample	Yb	Lu	Mo	Cu	Pb	Zn	As	Cd	Sb	Bi	Ag	V/V+Ni	Th/U	V/Cr
1	0.2	0.03	0.7	34.5	890	602	48	1.1	35.8	0.1	0.08	0.824	0.364	1.5
2	0.7	0.09	2.5	63.8	302	112	173	2.3	10	0.2	84.4	0.854	0.147	1.888
3	0.3	0.05	1	1.2	99	102	11	0.01	6.4	0.1	0.1	0.793	0.222	1.25
4	1.7	0.2	0.5	8.4	64.6	78.23	15	bdl	1	0.2	0.2	0.671	0.272	2.263
5	2.1	0.34	0.4	6.6	20.5	16	26	0.1	5.7	0.4	0.4	0.767	0.882	4.430
6	0.3	0.04	1.1	53.6	74.02	23.7	179	0.15	30.4	0.1	0.02	0.688	0.269	0.937
7	0.4	0.06	1.4	1.4	53.9	46.2	5	0.3	4.4	0.1	0.1	0.818	1.533	1.187
8	0.4	0.07	2.7	5.4	202	144.1	27	2.3	14.6	0.1	1.3	0.858	0.432	2.312
9	0.7	0.11	4	64.1	503	233.1	20	2.9	16	1.5	3.5	0.819	0.591	2.263
10	0.1	0.02	0.7	37.1	12.05	8.3	43	0.89	34.5	0.1	13	0.913	0.19	2.571
11	0.3	0.04	1.1	1.4	402	47.3	11	0.5	6.2	0.1	0.1	0.697	0.273	0.6
12	0.2	0.03	0.7	74.3	83	76.4	65	1	55.8	0.1	1	0.895	0.208	2.866
13	1.4	0.22	1.6	1793	37	45.2	7	0.3	13.4	3.9	0.12	0.839	1.195	4.142

Table 1

Feeder zone			Massive ore facies			Bedded ore facies		
Sample No,	Mineralogy	$\delta^{34}\text{S}_{\text{SDT}}$	Sample No,	Mineralogy	$\delta^{34}\text{S}_{\text{SDT}}$	Sample No,	Mineralogy	$\delta^{34}\text{S}_{\text{SDT}}$
Meh-01	Cpy2	-1.6	Meh-17	Cpy1	-7.6	Meh-33	Gn1	-17.4
Meh-02	Cpy2	4.6	Meh-18	Cpy1	-9.6	Meh-34	Gn1	-17
Meh-03	Cpy1	-7.3	Meh-19	Cpy2	3.7	Meh-35	Gn1	-17
Meh-04	Cpy2	3.8	Meh-20	Cpy2	-1.2	Meh-36	Gn1	-17
Meh-05	Cpy1	-6.9	Meh-21	Gn2	3.2	Meh-37	Gn1	-17.3
Meh-06	Gn1	-10.5	Meh-22	Gn1	-11.5	Meh-38	Gn1	-17.7
Meh-07	Gn1	-7.9	Meh-23	Gn2	-11.2	Meh-39	Gn1	-18.1
Meh-08	Gn2	4.4	Meh-24	Gn1	-13.7	Meh-40	Gn1	-17.3
Meh-09	Gn2	3.1	Meh-25	Gn1	-13.5	Meh-41	Sph1	-17
Meh-10	Sph2	4.6	Meh-26	Sph1	-11.6	Meh-42	Sph1	-18
Meh-11	Sph1	-3.3	Meh-27	Sph2	2.3	Meh-43	Sph1	-17.6
Meh-12	Sph1	-5.2	Meh-28	Sph2	-6.5	Meh-44	Py1	-20.1
Meh-13	Sph2	-1.5	Meh-29	Sph1	-9.3	Meh-45	Py1	-20.5
Meh-14	Ba	22	Meh-30	Ba	20.1	Meh-46	Py1	-21.3
Meh-15	Ba	22.1	Meh-31	Ba	19.8	Meh-47	Py1	-21.9
Meh-16	Ba	22.4	Meh-32	Ba	19.65	Meh-48	Py1	-22
						Meh-49	Ba	22.3
						Meh-50	Ba	17.8
						Meh-51	Ba	19.9
						Meh-52	Ba	18.7

Cpy: chalcopyrite; Gn: Galena; Sph: sphalerite; Py: pyrite; Ba: barite

Table 2

Drill Hole	Sample no.	Depth (meter)	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Ba (%)	Drill Hole	Sample no.	Depth (meter)	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Ba (%)
8316 (300m)	1	30	613	3900	37800	26	67	7913 (340m)	1	45	54	324	4430	32	49
	2	37	402	11900	73500	10.8	59		2	59	102	457	4110	29	53
	3	46	982	5800	39600	13.9	58		3	70	108	299	3490	12	46
	4	56	598	2900	19700	19	54		4	87	203	6500	23700	62	42
	5	62	144	5900	48000	88	60		5	111	128	909	38300	19	31
	6	74	1658	113600	99500	56	63		6	124	542	1790	42900	42	36
	7	80	1983	79600	352200	18.8	50		7	130	987	1110	25300	62	35
	8	94	2900	54700	330100	39	54		8	146	985	19400	77300	53	29
	9	106	2645	130600	231300	61	45		9	159	1752	18700	76300	67	23
	10	120	2745	157900	175000	150	42		10	170	1047	22100	71700	96	29
	11	135	2900	30800	107600	107	33		11	185	1958	33100	65900	61	25
	12	146	2974	11400	68700	98	32		12	192	2901	38200	64500	78	14
	13	152	3098	24800	84200	140	26		13	202	2658	49200	62900	42	19
	14	163	3021	6700	48500	92	24		14	219	2814	15500	40600	49	8
	15	178	4002	6900	34400	84	27		15	236	3601	9200	58000	61	13
	16	182	4987	10500	58400	99	37		16	348	2589	6300	41900	35	14
	17	190	5237	11300	48500	62	14		17	260	3002	61000	35600	52	10
	18	195	5947	21500	31500	80	19		18	271	3298	48700	50800	62	12
	19	201	5387	21500	68700	58	5		19	286	4001	34800	53600	12	7
	20	225	6002	21100	69000	96	12		20	305	3628	31600	61200	29	2
	21	249	6284	7400	34300	84	6		21	321	4125	27400	56700	58	8
	22	261	5124	8000	35200	102	11		22	332	4852	12400	30700	46	3
	23	276	5400	4600	50600	84	8		-	-	-	-	-	-	-
Drill Hole	Sample no.	Depth (meter)	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Ba (%)	Drill Hole	Sample no.	Depth (meter)	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Ba (%)
6901 (360m)	1	142	102	812	2640	12	30	7008 (280m)	1	140	210	1600	2900	29	23
	2	158	309	1025	5320	23	42		2	153	720	762	5620	34	33
	3	162	402	982	4430	19	24		3	167	750	534	8750	29	27
	4	176	563	2310	6140	32	60		4	173	1970	700	11300	30	24
	5	185	1023	15600	79500	27	22		5	182	1700	34300	4400	36	13
	6	196	856	10900	50400	36	39		6	193	1200	2300	3800	65	20
	7	205	1001	1990	4570	42	24		7	203	654	307	6270	36	12
	8	216	984	5400	24500	40	12		8	211	2100	600	13300	78	14
	9	226	1023	15500	30100	41	19		9	215	3200	4600	8900	55	13
	10	239	2745	3700	98900	63	20		10	216	1300	11200	11800	38	14
	11	250	1964	7600	197000	39	12		11	217	3450	2900	6200	30	8
	12	259	2037	18600	244000	46	14		12	218	3120	11500	10600	49	14
	13	267	1029	37900	219000	28	12		13	219	700	2900	31300	52	10
	14	270	3014	64700	156000	35	13		14	220	3970	34300	24800	51	8
	15	275	2687	13600	82700	41	8		15	222	1205	18600	21700	62	7
	16	280	2158	1900	35100	42	13		16	227	6740	13100	28600	47	7
	17	289	2035	15800	41700	48	9		17	235	1470	15300	41900	39	4
	18	295	1984	177000	73000	52	10		18	239	1775	12200	41400	37	3
	19	301	1402	567	2700	19	9		19	245	1680	15900	30200	48	10
	20	309	987	1235	1105	26	8		20	246	1780	24200	37200	40	9
	21	316	1058	109	727	27	11		21	249	1700	36400	52200	84	8
	22	320	1023	244	856	31	5		22	250	1370	24400	53400	40	7
Drill Hole	Sample no.	Depth (meter)	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Ba (%)	Drill Hole	Sample no.	Depth (meter)	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Ba (%)
5102 (224m)	1	180	500	23500	20800	26	11	7004 (200m)	1	140	57	1900	5600	1	10
	2	182	700	42700	58900	10.8	9		2	145	8	1080	4210	1	9
	3	184	300	28600	55800	13.9	10		3	149	-5	800	8600	10	7
	4	186	700	35900	100500	19	9		4	153	691	1700	17700	12	9
	5	189	3400	34500	126500	88	12		5	156	53	3200	16200	11	8
	6	192	3300	27400	65600	56	4		6	160	70	9700	22500	11	12
	7	194	300	10000	35700	18.8	7		7	162	45	5500	19400	10	10
	8	196	400	10300	28400	19	12		8	170	10	1100	6000	16	7
	9	198	100	9700	34800	11	11		9	179	12	74	420	10	8
	10	201	200	13400	24200	30	7		-	-	-	-	-	-	-

Drill Hole	Sample no.	Depth (meter)	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Ba (%)	Drill Hole	Sample no.	Depth (meter)	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Ba (%)
5303 (191m)	1	135	34	27200	23600	8	8	5203 (206)	1	167	31	171	3330	1	6
	2	139	30	20600	18900	9	10		2	169	12	291	5000	2	3
	3	142	27	15400	19900	7	6		3	171	7	437	6100	2	5
	4	146	22	8200	21400	17	8		4	173	59	1110	25300	6	7
	5	150	30	13100	21500	22	9		5	175	115	15500	40600	4	4
	6	154	51	21700	22600	15	14		6	176	113	4380	18000	11	9
	7	158	18	7600	16600	22	12		7	178	72	3790	14700	8	3
	8	160	69	34300	26000	15	6		8	179	57	3240	14300	16	2
	9	164	53	27600	27800	17	8		9	180	41	7600	19900	5	6
	10	168	47	28200	30200	13	9								
	11	170	24	15500	20600	21	9								
	12	172	32	21000	26500	25	5								
	13	173	21	8000	18400	12	3								
	14	175	16	4400	14600	15	6								
	15	176	14	6400	18600	19	7								

Table 3