Caterpillars are Antimagic

Antoni Lozano* Mercè Mora† Carlos Seara‡ Joaquín Tey§

Abstract

An *antimagic labeling* of a graph G is a bijection from the set of edges $E(G)$ to $\{1, 2, \dots, |E(G)|\}$ such that all vertex sums are pairwise distinct, where the *vertex sum* at vertex u is the sum of the labels assigned to the edges incident to u . A graph is called *antimagic* when it has an antimagic labeling. Hartsfield and Ringel conjectured that every simple connected graph other than K_2 is antimagic and the conjecture remains open even for trees. Here we prove that caterpillars are antimagic by means of an $O(n \log n)$ algorithm.

1 Introduction

All graphs considered in this paper are finite, undirected, connected, and simple. Given a graph G , we denote its set of vertices by $V(G)$ and its set of edges by $E(G)$. For every vertex $v \in V(G)$, we denote by $E_G(v)$ the set of edges incident to v in G . The degree of a vertex $v \in V(G)$ is $d_G(v) = |E_G(v)|$ (we will just write $d(v)$ when G is clear from context). For undefined terminology about basic graph theory we refer the reader to [4].

An (*edge*) *labeling* of a graph G is an injection from $E(G)$ to the set of nonnegative integers. A labeling of G is called *antimagic* if it is a bijection $\phi : E(G) \rightarrow \{1, 2, \dots, |E(G)|\}$ such that all vertex sums are pairwise distinct, where the *vertex sum* at vertex u is $\sum_{e \in E_G(u)} \phi(e)$. A graph is called *antimagic* if it has an antimagic labeling. The following conjecture from Hartsfield and Ringel [13] is well known.

Conjecture 1. [13] *Every connected graph other than K_2 is antimagic.*

Classes of graphs which are known to be antimagic include: paths, stars, complete graphs, cycles, wheels, and complete bipartite graphs $K_{2,m}$ [13]; graphs of order n with maximum degree at least $n - 3$ [24]; dense graphs (i.e., graphs having minimum degree $\Omega(\log n)$) and complete partite graphs but K_2 [2, 11]; toroidal grid graphs [22]; lattice grids

*Computer Science Department, Universitat Politècnica de Catalunya, Spain, antoni@cs.upc.edu

†Mathematics Department, Universitat Politècnica de Catalunya, Spain, merce.mora@upc.edu

‡Mathematics Department, Universitat Politècnica de Catalunya, Spain, carlos.seara@upc.edu

§Math. Department, Universidad Autónoma Metropolitana-Iztapalapa, México, jtey@xanum.uam.mx

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 734922.

and prisms [6]; regular bipartite graphs [8]; odd degree regular graphs [9]; even degree regular graphs [5]; cubic graphs [21]; generalized pyramid graphs [3]; graph products [23]; and Cartesian product of graphs [7, 25] (see the dynamic survey [12] for more details on antimagic labelings). However, the conjecture is still open for the general class of trees.

Conjecture 2. *[13] Every tree other than K_2 is antimagic.*

One of the best known results for trees is due to Kaplan, Lev, and Roditty [16], who proved that any tree having more than two vertices and at most one vertex of degree two is antimagic (see also [20]). In this paper we focus on *caterpillars*, that is, trees of order at least 3 such that the removal of their leaves produces a path. In [19] the authors give sufficient conditions for a caterpillar to be antimagic and, recently, it has been shown that that caterpillars with maximum degree 3 are antimagic [10]. In this paper we take a step further proving that every caterpillar is antimagic.

Theorem 1. *Caterpillars are antimagic. Furthermore, there exists an algorithm that, given a caterpillar C of order n , produces an antimagic labeling for C in time $O(n \log n)$.*

Concretely, in Section 2 we provide the above mentioned algorithm, whose correctness and time bound are shown in Section 3. Therefore, our method follows a constructive approach, in contrast with the use of the Combinatorial NullStellenSatz method [1, 14, 17], which is the regular technique used in several of the references above.

2 Construction of an Antimagic Labeling

Suppose that C is a caterpillar with m edges and let $P = (v_0, \dots, v_p)$ be a longest path in C . Edges in P will be called *pathedges*, while edges not in P will be called *legs*. We use the notation $e_i = \{v_{i-1}, v_i\}$ ($1 \leq i \leq p$) for pathedges and f_i for the leg incident to a vertex v_i of degree 3. We define $E_0 = \{e_i : i \equiv p \pmod{2}\}$ and $E_1 = \{e_i : i \not\equiv p \pmod{2}\}$, so that $\{E_0, E_1\}$ is a partition of the set of pathedges such that $e_p \in E_0$ and any two incident edges of P belong to different sets. The size of these sets is $|E_0| = \lceil p/2 \rceil$ and $|E_1| = \lfloor p/2 \rfloor$.

Let $[a, b] = \{n \in \mathbb{N} : a \leq n \leq b\}$ and $[a, \infty) = \{n \in \mathbb{N} : n \geq a\}$, where a and b are positive integers. Split the set of available labels $L = [1, m]$ into the two subsets $L_0 = [1, m - \lfloor p/2 \rfloor]$ and $L_1 = [m - \lfloor p/2 \rfloor + 1, m]$ of sizes $m - \lfloor p/2 \rfloor$ and $\lfloor p/2 \rfloor$, respectively.

We now describe an algorithm in three steps to construct a labeling ϕ of C that will be shown to be antimagic in Section 3 (see the pseudocode in Page 4 and an example in Figure 1). Given a vertex u in C and a labeling ϕ of C , we denote the vertex sum at u by $s(u)$, that is, $s(u) = \sum_{e \in E_C(u)} \phi(e)$. When we describe the construction of the labeling, we also use the notation $s(u)$ to refer to a *partial sum* at u , that is, the sum of the values $\phi(e)$ for all edges e incident to u for which a label has already been assigned at that step of the algorithm. Similarly, we use $\phi(E(C))$ to denote the set of labels used up to that step.

In the first step, a label is assigned to all pathedges and a few legs incident to vertices of degree 3. Roughly speaking, the algorithm alternatively assigns consecutive labels from

the lists L_0 and L_1 to the pathedges with some exceptions, and taking into account that the label of edge e_p must belong to L_0 . Concretely, the edges of E_1 receive labels from L_1 in increasing order, and the edges of E_0 receive labels from L_0 in increasing order with the following exception. When assigning a label to the edge $e_i \in E_0$, $1 < i \leq p$, we check if the following condition holds:

$Q(i)$: $\phi(e_{i-1}) = \phi(e_k) + \phi(e_{k+1})$ for some $k \in \{1, \dots, i-3\}$ such that $d(v_k) = 3$.

If $Q(i)$ does not hold, then we assign the next unused label of L_0 to e_i . Otherwise, we assign the next unused label of L_0 to the leg f_k and the next one, to e_i . Such a vertex v_k of degree 3, whose leg f_k has been labeled at this step, will be called *light vertex*, while the rest of vertices of degree at least three will be called *heavy vertices*. The set of all light vertices obtained when finishing this step is denoted by U_p , while the set of heavy vertices is denoted by W .

In the second step, for each heavy vertex v , we randomly assign unused labels from L_0 to all but one of the legs incident to v .

Finally, it only remains to label one leg for each heavy vertex. To do that, in the third step we list the remaining legs in increasing order of the partial vertex sums at their corresponding incident heavy vertices. Then, we sort the remaining labels in L_0 in increasing order and assign them to the previous list of legs in the same order.

Figure 1: Example of a labeling of a caterpillar with a longest path of odd length. In this case, $L_0 = [1, 21]$ and $L_2 = [22, 27]$. Top, Step 1: Labeling the pathedges and legs incident to light vertices (circled vertices). Middle, Step 2: Labeling all legs but one for each heavy vertex (squared vertices). Bottom, Step 3: Labeling the last leg for each heavy vertex.

Algorithm 1 Antimagic Labeling of a Caterpillar.

Input: A caterpillar C of order n

Output: An antimagic labeling ϕ for C

STEP 1: LABELING THE PATHEDGES AND A FEW LEGS

```
1:  $\phi(e_1) \leftarrow 1$ 
2:  $\phi(e_2) \leftarrow m - \lfloor p/2 \rfloor + 1$ 
3:  $U_1, U_2 \leftarrow \emptyset$ 
4: if  $p$  is even then
5:   Exchange  $\phi(e_1)$  and  $\phi(e_2)$ 
6: for  $i = 3$  to  $p$  do
7:    $\phi(e_i) \leftarrow \phi(e_{i-2}) + 1$ 
8:    $U_i \leftarrow U_{i-1}$ 
9:   if  $i \equiv p \pmod{2}$  and  $\phi(e_{i-1}) = \phi(e_k) + \phi(e_{k+1})$ 
       for some  $k \in \{1, \dots, i-3\}$  s.t.  $d(v_k) = 3$  then
10:     $\phi(f_k) \leftarrow \phi(e_i)$ 
11:     $\phi(e_i) \leftarrow \phi(e_i) + 1$ 
12:     $U_i \leftarrow U_i \cup \{v_k\}$ 
13:  $W \leftarrow \{v \in V(C) \mid v \notin U_p \wedge d(v) \geq 3\}$ 
```

STEP 2: LABELING ALL LEGS EXCEPT ONE FOR EACH VERTEX IN W

```
14: for all  $v \in W$  do
15:   for all legs  $e$  incident to  $v$  except one do
16:      $\phi(e) \leftarrow$  a random label from  $L_0 \setminus \phi(E(C))$ 
```

STEP 3: LABELING THE LAST LEG OF EACH VERTEX IN W

```
17: Sort the vertices in  $W$  as  $w_1, \dots, w_t$  s.t.  $s(w_i) \leq s(w_{i+1})$  for all  $i < t$ 
18: Sort the labels in  $L_0 \setminus \phi(E(C))$  as  $\ell_1, \dots, \ell_t$  in increasing order
19: for  $i = 1$  to  $t$  do
20:    $\phi(w_i) = \ell_i$ 
```

3 Proof of Theorem 1

In this section we prove that the labeling produced by Algorithm 1 for a caterpillar of order n is antimagic and can be found in time $O(n \log n)$, thus proving Theorem 1. The following lemmas will be used in the proofs of correctness and efficiency of the algorithm.

We first introduce some notation. We denote by F_0 the set of legs incident to light vertices. Recall that the set U_i introduced in the algorithm contains light vertices, that is, vertices of degree 3 whose legs have been labeled up to iteration i of the *for* loop in Step 1. Hence, $|F_0| = |U_p|$. Let E_2 be the set of edges not belonging to $E_0 \cup E_1 \cup F_0$, that is, $\{E_0, E_1, E_2, F_0\}$ is a partition of the edge set $E(C)$.

Lemma 1. For every $i \in \{1, 2, \dots, p\}$ we have

$$\phi(e_i) = \begin{cases} \lceil \frac{i}{2} \rceil + |U_i|, & \text{if } e_i \in E_0; \\ m - \lfloor \frac{p}{2} \rfloor + \lceil \frac{i}{2} \rceil, & \text{if } e_i \in E_1. \end{cases} \quad (1)$$

Moreover,

- (1) $\phi(E_0 \cup F_0) = [1, \lceil p/2 \rceil + |U_p|] \subseteq L_0$;
- (2) $\phi(E_1) = L_1 = [m - \lfloor p/2 \rfloor + 1, m]$;
- (3) $\phi(E_2) = [\lceil p/2 \rceil + |U_p| + 1, m - \lfloor p/2 \rfloor] \subseteq L_0$.

Proof. First we check that the values $\phi(e_1)$ and $\phi(e_2)$ defined in Equation 1 are correct. Indeed, if p is odd, then $e_1 \in E_0$, $e_2 \in E_1$, and the labels assigned after the execution of lines 1–5 of the algorithm are $\phi(e_1) = 1 = \lceil 1/2 \rceil + |U_1|$ and $\phi(e_2) = m - \lfloor p/2 \rfloor + 1 = m - \lfloor p/2 \rfloor + \lceil 2/2 \rceil$. If p is even, then $e_1 \in E_1$, $e_2 \in E_0$, and the labels assigned in lines 1–5 are $\phi(e_1) = m - \lfloor p/2 \rfloor + 1 = m - \lfloor p/2 \rfloor + \lceil 1/2 \rceil$ and $\phi(e_2) = 1 = \lceil 2/2 \rceil + |U_2|$.

Now, we show that the value $\phi(e_i)$, $3 \leq i \leq p$, defined in Equation 1 corresponds to the same value calculated in the *for* loop at lines 6–12. Suppose that $3 \leq i \leq p$ and all values given in Equation 1 up to $\phi(e_{i-1})$ are correct. Then, either $e_i \in E_0$ or $e_i \in E_1$. Suppose first that $e_i \in E_1$. Then, lines 10–12 are not executed and $\phi(e_i) = \phi(e_{i-2}) + 1$. Since $e_{i-2} \in E_1$ and, by hypothesis, $\phi(e_{i-2}) = m - \lfloor p/2 \rfloor + \lceil \frac{i-2}{2} \rceil$, we have

$$\phi(e_i) = m - \lfloor \frac{p}{2} \rfloor + \lceil \frac{i-2}{2} \rceil + 1 = m - \lfloor \frac{p}{2} \rfloor + \lceil \frac{i-2}{2} + 1 \rceil = m - \lfloor \frac{p}{2} \rfloor + \lceil \frac{i}{2} \rceil.$$

Now suppose that $e_i \in E_0$. In this case, $e_{i-2} \in E_0$ and, by hypothesis, $\phi(e_{i-2}) = \lceil \frac{i-2}{2} \rceil + |U_{i-2}|$. Note that $e_{i-1} \in E_1$ and, then, the iteration for $i-1$ produces $U_{i-1} = U_{i-2}$. If condition $Q(i)$ does not hold, then lines 10–12 are not executed. Thus, $|U_i| = |U_{i-1}|$ and $\phi(e_i) = \phi(e_{i-2}) + 1$, and we get

$$\phi(e_i) = \lceil \frac{i-2}{2} \rceil + |U_{i-2}| + 1 = \lceil \frac{i-2}{2} + 1 \rceil + |U_{i-2}| = \lceil \frac{i}{2} \rceil + |U_i|.$$

Otherwise, after the execution of lines 10–12 we have that $\phi(e_i) = \phi(e_{i-2}) + 2$ and $|U_i| = |U_{i-1}| + 1$. Since $e_{i-2} \in E_0$ and, by hypothesis, $\phi(e_{i-2}) = \lceil \frac{i-2}{2} \rceil + |U_{i-2}|$, we obtain

$$\phi(e_i) = \lceil \frac{i-2}{2} \rceil + |U_{i-2}| + 2 = \lceil \frac{i-2}{2} + 1 \rceil + |U_{i-2}| + 1 = \lceil \frac{i}{2} \rceil + |U_i|.$$

In both of the above cases, the value $\phi(e_i)$ computed by the algorithm coincides with that defined in Equation 1.

Now, we prove the first item. On the one hand, the values obtained for the edges in E_0 range between 1 and $\lceil p/2 \rceil + |U_p|$ by Equation 1, and the value of a leg $f_k \in F_0$ (i.e., a leg incident to a light vertex) is less than the value of some edge belonging to E_0 . Hence, $\phi(E_0 \cup F_0) \subseteq [1, \lceil p/2 \rceil + |U_p|]$. Since ϕ is an injection and $|E_0 \cup F_0| = \lceil p/2 \rceil + |U_p|$, we

have $\phi(E_0 \cup F_0) = [1, \lceil p/2 \rceil + |U_p|]$. On the other hand, $|E_0 \cup F_0| \leq m - |E_1| = m - \lfloor p/2 \rfloor$. Thus, $\phi(E_0 \cup F_0) = [1, \lceil p/2 \rceil + |U_p|] \subseteq L_0$.

As for the second item, since $|E_1| = \lfloor p/2 \rfloor = |L_1|$, $\phi(E_1) \subseteq L_1$ and ϕ is an injection, we have that $\phi(E_1) = L_1 = [m - \lfloor p/2 \rfloor + 1, m]$. Finally, in Steps 2 and 3, the algorithm assigns the remaining labels to the remaining edges. Therefore, $\phi(E_2) = [1, m] \setminus \phi(E_0 \cup F_0 \cup E_1) = [\lceil p/2 \rceil + |U_p| + 1, m - \lfloor p/2 \rfloor] \subseteq L_0$ and the third item is true. $\square$

Lemma 2. *For any i, j such that $1 \leq i \leq j < p$, we have:*

- (1) $m - \lfloor p/2 \rfloor + 2 \leq \phi(e_i) + \phi(e_{i+1}) \leq m + \lceil p/2 \rceil + |U_p|$;
- (2) If $i < j$, then $\phi(e_i) + \phi(e_{i+1}) < \phi(e_j) + \phi(e_{j+1})$;
- (3) If v_i and v_j are light vertices and $i < j$, then $\phi(f_i) < \phi(f_j)$.

Proof. We begin by proving the first item. Since one of the edges from $\{e_i, e_{i+1}\}$ belongs to E_0 and the other one to E_1 , according to Equation 1 in Lemma 1, we have

$$\phi(e_i) + \phi(e_{i+1}) = m - \left\lfloor \frac{p}{2} \right\rfloor + \left\lceil \frac{i}{2} \right\rceil + \left\lceil \frac{i+1}{2} \right\rceil + h,$$

for some h , $0 \leq h \leq |U_p|$. Hence, the lower bound trivially holds. For the upper bound, note that

$$m - \left\lfloor \frac{p}{2} \right\rfloor + \left\lceil \frac{i}{2} \right\rceil + \left\lceil \frac{i+1}{2} \right\rceil + h \leq m - \left\lfloor \frac{p}{2} \right\rfloor + \left\lceil \frac{p-1}{2} \right\rceil + \left\lceil \frac{p}{2} \right\rceil + |U_p| \leq m + \left\lceil \frac{p}{2} \right\rceil + |U_p|.$$

To prove the second item, let $g : [1, p-1] \rightarrow \mathbb{N}$ be the function

$$g(i) = \phi(e_i) + \phi(e_{i+1}).$$

Now, it is enough to show that function g is strictly increasing. Note that for any i such that $1 \leq i \leq p-2$, proving $g(i) < g(i+1)$ is equivalent to proving the inequality:

$$\phi(e_i) < \phi(e_{i+2}) \tag{2}$$

whenever $1 \leq i \leq p-2$. Hence, we consider two cases according to the value of ϕ for pathedges given in Equation 1. If $e_i \in E_0$, then

$$\phi(e_i) < \phi(e_{i+2}) \Leftrightarrow \left\lceil \frac{i}{2} \right\rceil + |U_i| < \left\lceil \frac{i+2}{2} \right\rceil + |U_{i+2}| \Leftrightarrow |U_i| < 1 + |U_{i+2}|,$$

which is true. Now, if $e_i \in E_1$, then

$$\phi(e_i) < \phi(e_{i+2}) \Leftrightarrow m - \left\lfloor \frac{p}{2} \right\rfloor + \left\lceil \frac{i}{2} \right\rceil < m - \left\lfloor \frac{p}{2} \right\rfloor + \left\lceil \frac{i+2}{2} \right\rceil \Leftrightarrow \left\lceil \frac{i}{2} \right\rceil < \left\lceil \frac{i+2}{2} \right\rceil,$$

which is also true. This concludes the proof of the second item.

Let us now prove the third item. Suppose that $1 \leq i < j \leq p-1$. Notice that labels $\phi(f_i)$ and $\phi(f_j)$ are assigned in the first step of the algorithm if $Q(i')$ and $Q(j')$

hold when assigning the label of some pathedges $e_{i'}$ and $e_{j'}$ of E_0 , with $i' > i$ and $j' > j$, and such that $\phi(e_{i'-1}) = \phi(e_i) + \phi(e_{i+1})$ and $\phi(e_{j'-1}) = \phi(e_j) + \phi(e_{j+1})$. On the one hand, since $i < j$, we deduce by the preceding item that $\phi(e_{i'-1}) < \phi(e_{j'-1})$. Besides, $e_{i'-1}, e_{j'-1} \in E_1$, and from Inequality 2 we deduce $i' < j'$. On the other hand, the labels assigned to the edges f_i and f_j are $\phi(f_i) = \phi(e_{i'-2}) + 1$ and $\phi(f_j) = \phi(e_{j'-2}) + 1$, where $e_{i'-2}, e_{j'-2} \in E_0$ and $i' - 2 < j' - 2$. Again, from Inequality 2, we conclude that $\phi(f_i) = \phi(e_{i'-2}) + 1 < \phi(e_{j'-2}) + 1 = \phi(f_j)$. $\square$

3.1 Proof of Correctness

Now we show the correctness of the algorithm, that is, we show that, given a caterpillar C , Algorithm 1 produces an antimagic labeling of C .

Since $E(C) = E_0 \cup E_1 \cup E_2 \cup F_0$, we have by Lemma 1 that ϕ is a bijection from $E(C)$ onto $[1, m]$. To prove that ϕ is an antimagic labeling, it remains to show that all vertex sums are pairwise different.

Consider the partition $\{V_1, V_2, V_3\}$ of $V(C)$, where V_1 is the set of vertices of degree 1, V_2 is the set of vertices of degree 2 and light vertices, and V_3 is the set of heavy vertices. Let $s(V_i) = \{s(v) : v \in V_i\}$, for $1 \leq i \leq 3$. It is enough to prove that for each i , $1 \leq i \leq 3$, all sums in $s(V_i)$ are pairwise distinct and the sets $s(V_1)$, $s(V_2)$, $s(V_3)$ are pairwise disjoint.

We begin by checking that the vertices of V_1 have pairwise distinct vertex sums. The vertex sum at a vertex of degree one is the value of the label of the pendant edge incident to it. Since all edges receive different labels, vertex sums at vertices of degree one are pairwise distinct.

Now let us establish an interval of possible values in $s(V_1)$. Notice that a pendant edge is either a leg, or the first or the last edge of the path P . According to Lemma 1, legs are always labeled with elements from L_0 . Regarding the first and last edges of the path P , if p is odd, their labels are from L_0 , whereas if p is even, these pathedges are labeled with the smallest values of L_1 and L_0 , respectively, that is, $m - \lfloor p/2 \rfloor + 1$ and 1. Hence, $s(V_1) \subseteq L_0 \cup \{m - \lfloor p/2 \rfloor + 1\} = [1, m - \lfloor p/2 \rfloor + 1]$.

In order to check that all vertices of V_2 have pairwise distinct sums, consider two distinct vertices $v_j, v_k \in V_2$. Now, we distinguish three cases. In the first case, suppose that v_j and v_k are vertices of degree 2. If $1 \leq j < k < p$, then their sums can be expressed as

$$s(v_j) = \phi(e_j) + \phi(e_{j+1}), \quad s(v_k) = \phi(e_k) + \phi(e_{k+1}),$$

and by Lemma 2, $s(v_j) < s(v_k)$. In the second case, suppose that both v_j and v_k are light vertices. If $1 \leq j < k < p$, then, similarly to the first case, their sums are

$$s(v_j) = \phi(e_j) + \phi(e_{j+1}) + \phi(f_j), \quad s(v_k) = \phi(e_k) + \phi(e_{k+1}) + \phi(f_k)$$

and, by Lemma 2 we conclude that $s(v_j) < s(v_k)$. In the third case, suppose that one of the vertices, say v_j , has degree 2 and the other one, that is v_k , is a light vertex. In such a case, we know that $Q(i)$ holds for some i , with $k + 2 < i < p$, that is,

$$\phi(e_k) + \phi(e_{k+1}) = \phi(e_{i-1}), \quad \phi(f_k) = \phi(e_{i-2}) + 1, \quad \text{and} \quad \phi(e_i) = \phi(e_{i-2}) + 2.$$

Besides,

$$s(v_k) = \phi(e_k) + \phi(e_{k+1}) + \phi(f_k) \text{ and } s(v_j) = \phi(e_j) + \phi(e_{j+1}).$$

Now, by applying Lemma 2, if $j \leq k$, then

$$\begin{aligned} s(v_j) &= \phi(e_j) + \phi(e_{j+1}) \leq \phi(e_k) + \phi(e_{k+1}) \\ &< \phi(e_k) + \phi(e_{k+1}) + \phi(f_k) = s(v_k); \end{aligned}$$

if $k < j \leq i - 2$, then

$$\begin{aligned} s(v_j) &= \phi(e_j) + \phi(e_{j+1}) \leq \phi(e_{i-2}) + \phi(e_{i-1}) \\ &< \phi(e_{i-1}) + \phi(e_{i-2}) + 1 \\ &= \phi(e_k) + \phi(e_{k+1}) + \phi(f_k) = s(v_k); \end{aligned}$$

and if $k < i - 1 \leq j$, then

$$\begin{aligned} s(v_j) &= \phi(e_j) + \phi(e_{j+1}) \geq \phi(e_{i-1}) + \phi(e_i) \\ &= \phi(e_{i-1}) + \phi(e_{i-2}) + 2 \\ &= \phi(e_k) + \phi(e_{k+1}) + \phi(f_k) + 1 > s(v_k). \end{aligned}$$

Hence, all vertices in V_2 have different vertex sums.

We now determine the interval of possible values in $s(V_2)$. Let $v_k \in V_2$. For the lower bound, we have $s(v_k) \geq \phi(e_k) + \phi(e_{k+1}) \geq m - \lfloor p/2 \rfloor + 2$ by Lemma 2. For the upper bound, again by Lemma 2, the vertex sum at any vertex of degree two can be bounded by $m + \lceil p/2 \rceil + |U_p|$, while if v_k is a light vertex, then for some i with $k + 2 < i \leq p$ and $i \equiv p \pmod{2}$ we have:

$$s(v_k) = \phi(e_k) + \phi(e_{k+1}) + \phi(f_k) = \phi(e_{i-1}) + \phi(f_k) = \phi(e_{i-1}) + \phi(e_{i-2}) + 1 < m + \left\lceil \frac{p}{2} \right\rceil + |U_p| + 1.$$

Therefore, $s(V_2) \subseteq [m - \lfloor p/2 \rfloor + 2, m + \lceil p/2 \rceil + |U_p|]$.

Finally, recall that V_3 is the set of heavy vertices. Due to the way labels are assigned in Steps 2 and 3, all vertex sums of heavy vertices will be pairwise distinct.

Let us calculate the interval of possible values of their vertex sums. Let v_i be a heavy vertex and suppose, in the first place, that $d(v_i) = 3$. Since, by construction, the leg of v_i has not been labeled in Step 1, the partial sum of v_i after Step 1 must be at least $m + 1$. Indeed, suppose on the contrary that $\phi(e_i) + \phi(e_{i+1}) \leq m$. By Lemma 2, we would have $\phi(e_i) + \phi(e_{i+1}) = \ell \in [m - \lfloor p/2 \rfloor + 2, m] \subseteq L_1$, so that after assigning the label ℓ to some edge e_{j-1} of E_1 in Step 1, condition $Q(j)$ would have held for some j , with $j > i$, implying that the leg incident to v_i would have been labeled in Step 1, a contradiction. Hence, $\phi(e_i) + \phi(e_{i+1}) \geq m + 1$. By Lemma 1, the labels for its leg are at least $\lceil p/2 \rceil + |U_p| + 1$. Therefore,

$$s(v_i) \geq m + \left\lceil \frac{p}{2} \right\rceil + |U_p| + 2.$$

Suppose now that $d(v_i) \geq 4$. Then, since there are at least two legs incident with v_i , we have

$$\begin{aligned} s(v_i) &\geq \phi(e_i) + \phi(e_{i+1}) + \left(\left\lceil \frac{p}{2} \right\rceil + |U_p| + 1\right) + \left(\left\lceil \frac{p}{2} \right\rceil + |U_p| + 2\right) \\ &\geq m - \left\lfloor \frac{p}{2} \right\rfloor + 2 + 2 \cdot \left\lceil \frac{p}{2} \right\rceil + 2|U_p| + 3 \geq m + \left\lceil \frac{p}{2} \right\rceil + 2|U_p| + 5. \end{aligned}$$

Hence, $s(V_3) \subseteq [m + \lceil p/2 \rceil + |U_p| + 2, \infty)$.

Notice that $s(V_1)$, $s(V_2)$, and $s(V_3)$ have been shown to be included in pairwise disjoint intervals and, as a consequence, they must also be pairwise disjoint.

3.2 Proof of Efficiency

Finally, we show that Algorithm 1 runs in time $O(n \log n)$.

Assignments in Step 1 of the algorithm can be done in constant time, except for the one at line 13, which requires computing a set difference and can be done in time $O(m)$. Condition in line 9 (which is equivalent to the fact that both $e_i \in E_0$ and $Q(i)$ hold) can be checked in linear time globally due to the fact that partial sums are increasing as variable i increases, as shown in Lemma 2. Therefore, the cost of Step 1 is the result of the linear loop at lines 6–12 and the assignment at line 13, that is, $O(m)$. Step 2 visits at most m edges and assigns a random label to each of them, but labels can be chosen increasingly from the unused labels in L_0 , thus giving a cost $O(m)$. Step 3 requires time $O(m \log m)$ due to fact that the partial vertex sums must be sorted (line 17).

The total cost of the algorithm is, then, $O(m \log m)$, but since C is a tree, $n = m + 1$ and the cost can be expressed as $O(n \log n)$.

Theorem 1 now follows from the proofs of correctness and efficiency contained, respectively, in Subsections 3.1 and 3.2.

4 Conclusions and Open Problems

We consider a consequence of our main result regarding oriented graphs. Define a *labeling* of a directed graph D with m arcs as a bijection from the set of arcs of D to $[1, m]$. A labeling of D is said to be *antimagic* if all oriented vertex sums are pairwise distinct, where the *oriented vertex sum* of a vertex in D is the sum of labels of all incoming arcs minus that of all outgoing arcs. A graph is said to have an *antimagic orientation* if it has an orientation which admits an antimagic labeling. Hefetz, Mütze, and Schwartz [15] formulate the following conjecture.

Conjecture 3. [15] *Every connected graph admits an antimagic orientation.*

As the authors point out in the paper, every bipartite antimagic undirected graph G admits an antimagic orientation by simply orienting all edges in the same direction between the two stable sets of G . Therefore, we can derive the following corollary from Theorem 1, which was originally proved in [18].

Corollary 1. [18] *Caterpillars have antimagic orientations.*

Lobsters, defined as trees such that the removal of their leaves produces a caterpillar, are a natural class of trees to which one can try to apply the techniques presented in this paper. A first question in the line of the above corollary would be the following.

Open problem 1. *Do lobsters have antimagic orientations?*

More generally, we state the following question.

Open problem 2. *Are lobsters antimagic?*

5 Acknowledgments

Antoni Lozano is supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement ERC-2014-CoG 648276 AUTAR). Mercè Mora is supported by projects Gen. Cat. DGR 2017SGR1336, MINECO MTM2015-63791-R, and H2020-MSCA-RISE project 734922-CONNECT. Carlos Seara is supported by projects Gen. Cat. DGR 2017SGR1640, MINECO MTM2015-63791-R, and H2020-MSCA-RISE project 734922-CONNECT. Joaquín Tey is supported by project PRODEP-12612731.

References

- [1] N. Alon. Combinatorial Nullstellensatz. *Combinatorics, Probability and Computing*, 8, (1999), 7–29.
- [2] N. Alon, G. Kaplan, A. Lev, Y. Roditty, and R. Yuster. Dense graphs are antimagic. *Journal of Graph Theory*, 47(4), (2004), 297–309.
- [3] S. Arumugam, M. Miller, O. Phanalasy, and J. Ryan. Antimagic labeling of generalized pyramid graphs. *Acta Mathematica Sinica*, 30(2), (2014), 283–290.
- [4] G. Chartrand, L. Lesniak, and P. Zhang. *Graphs and Digraphs*, fifth edition. CRC Press, Boca Raton, 2011.
- [5] F. Chang, Y.-Ch. Liang, Z. Pan, and X. Zhu. Antimagic labeling of regular graphs. *Journal of Graph Theory*, 82(4), (2016), 339–349.
- [6] Y. Cheng. Lattice grids and prisms are antimagic. *Theoretical Computer Science*, 374(1-3), (2007), 66–73.
- [7] Y. Cheng. A new class of antimagic Cartesian product graphs. *Discrete Mathematics*, 308(24), (2008), 6441–6448.
- [8] D. W. Cranston. Regular bipartite graphs are antimagic. *Journal of Graph Theory*, 60(3), (2009), 173–182.

- [9] D. W. Cranston, Y.-Ch. Liang, and X. Zhu. Regular graphs of odd degree are antimagic. *Journal of Graph Theory*, 80(1), (2015), 28–33.
- [10] K. Deng and Y. Li. Caterpillars with maximum degree 3 are antimagic. *Discrete Mathematics*, 342, (2019), 1799–1801.
- [11] T. Eccles. Graphs of large linear size are antimagic. *Journal of Graph Theory*, 81(3), (2016), 236–261.
- [12] J. A. Gallian. A dynamic survey of graph labeling. *The Electronic Journal of Combinatorics*, 5, (2018), #DS6.
- [13] N. Hartsfield and G. Ringel. *Pearls in Graph Theory: A Comprehensive Introduction*. Academic Press, INC., Boston, 1990 (revised version, 1994), 108–109.
- [14] D. Hefetz. Antimagic graphs via the Combinatorial NullStellenSatz. *Journal of Graph Theory*, 50(4), (2005), 263–272.
- [15] D. Hefetz, T. Mütze, and J. Schwartz. On antimagic directed graphs. *Journal of Graph Theory*, 64 (2010), 219–232.
- [16] G. Kaplan, A. Lev, and Y. Roditty. On zero-sum partitions and antimagic trees. *Discrete Mathematics*, 309, (2009), 2010–2014.
- [17] A. Lladó and M. Miller. Approximate results for rainbow labelings. *Periodica Mathematica Hungarica*, 74(1), (2017), 11–21.
- [18] A. Lozano. Caterpillars Have Antimagic Orientations. *An. St. Univ. Ovidius Constanta Ser. Mat.*, 26(3), (2018), 171–180.
- [19] A. Lozano, M. Mora, and C. Seara. Antimagic labelings of caterpillars. *Applied Mathematics and Computation*, 347, (2019), 734–740.
- [20] Y.-Ch. Liang, T.-L. Wong, and X. Zhu. Antimagic labeling of trees. *Discrete Mathematics*, 331, (2014), 9–14.
- [21] Y.-Ch. Liang and X. Zhu. Antimagic labeling of cubic graphs. *Journal of Graph Theory*, 75 (1), (2014), 31–36.
- [22] T.-M. Wang. Toroidal grids are antimagic. *Proc. 11th Annual International Computing and Combinatorics Conference, COCOON'2005*, LNCS 3595, (2005), 671–679.
- [23] T.-M. Wang and C.-C. Hsiao. On anti-magic labeling for graph products. *Discrete Mathematics*, 308(16), (2008), 3624–3633.
- [24] Z. B. Yilma. Antimagic properties of graphs with large maximum degree. *Journal of Graph Theory*, 72(4), (2013), 367–373.
- [25] Y. Zhang and X. Sun. The antimagicness of the Cartesian product of graphs. *Theoretical Computer Science*, 410(8–10), (2009), 727–735.