

**Biological activity as an indicator of the effect of thermally dried and composted sewage sludges over the reclamation process of two residual soils from a limestone quarry**

**Biological effect of the addition of thermally dried and composted sewage sludges to residual soils from a limestone quarry**

**Effects on biological activity of thermally dried and composted sewage sludge addition to residual soils from a limestone quarry**

**Graciela Marando, Patricia Jiménez, Manuel Bonmatí**, *Department of Agri-Food Engineering and Biotechnology. Technical University of Catalonia. Campus Baix Llobregat. Av. Canal Olímpic, s.n., 08860 Castelldefels (Barcelona). Spain, Phone: 34-93-55221218, E-mail:manuel.bonmati@upc.edu*

**Marta Ginovart**, *Department of Applied Mathematics III. Technical University of Catalonia. Campus Baix Llobregat. Av. Canal Olímpic, s.n., 08860 Castelldefels (Barcelona). Spain.*

**Sewage sludge, Thermally dried, Composting, Enzymatic activities, Microbial biomass**

#### **Abstract**

The differences of the effects of three composted and three thermally dried sewage sludges on the biological fertility of two different residual soils (MNS and MLS, containing respectively 35% and 20% of fine earth) from a limestone quarry were evaluated under field conditions. Each sludge was mixed with each waste to reach an organic carbon content (C) of 1% (d. m) in the less than 2 mm fraction. Mixtures and control soils were sown and, after 13 months, sampling was performed after harvest.  $\beta$ -glucosidase activity,  $\beta$ -galactosidase activity, total (TCH) and extracted (ECH) carbohydrate content, microbial biomass (MB), extracted C (EC), basal respiration (BR) and cumulated  $\text{CO}_2$  ( $\text{CCO}_2$ ) were determined and their ratios with respect to C were calculated. Values of all the assayed absolute parameters were higher in the mixtures than in their relative residual soils. Sludge type had a significant effect in MB,  $\beta$ -glucosidase activity and ECH. TCH and EC were significantly higher in mixture samples containing MNS.  $\text{CCO}_2$  and BR/C were generally higher in mixtures containing thermally dried sludges. EC/C was the index better detecting differences among sludge type effects on the soils.