

Note on the number of obtuse angles in point sets

Ruy Fabila-Monroy¹ Clemens Huemer² Eulàlia Tramuns³

Abstract

In 1979 Conway, Croft, Erdős and Guy proved that every set S of n points in general position in the plane determines at least $\frac{n^3}{18} - O(n^2)$ obtuse angles and also presented a special set of n points to show the upper bound $\frac{2n^3}{27} - O(n^2)$ on the minimum number of obtuse angles among all sets S . We prove that every set S of n points in convex position determines at least $\frac{2n^3}{27} - o(n^3)$ obtuse angles, hence matching the upper bound (up to sub-cubic terms) in this case. Also on the other side, for point sets with low rectilinear crossing number, the lower bound on the minimum number of obtuse angles is improved.

Introduction

A point set S in the plane is in general position if no three points of the set lie on a common straight line. Throughout this paper, all considered point sets S will be in general position in the plane and $|S| = n$. An angle abc at b determined by three points $\{a, b, c\}$ of S is obtuse if it is greater than $\frac{\pi}{2}$. Prominent problems and results on obtuse and acute angles in point sets can be found in Ref. 3. For example, 2^n points in the plane ($n \geq 3$) always determine an angle greater than $\pi(1 - \frac{1}{n})$. Another classical result is that $2^d + 1$ points in d -dimensional Euclidean space E_d always determine at least one obtuse angle. An open problem (for higher dimension d) is to determine the largest integer $f(d)$ for which there are $f(d)$ points in E_d all the angles of which are less than $\frac{\pi}{2}$.

Here we are interested in the number of obtuse angles determined by point sets S in the plane. Conway et al., see Ref. 2, proved that the minimum number

¹Departamento de Matemáticas, Cinvestav-IPN, Av. IPN 2508, Col. San Pedro Zacatenco 07360, D.F., Mxico. ruyfabila@math.cinvestav.edu.mx

²Departament de Matemàtica Aplicada IV, Universitat Politècnica de Catalunya, C. Esteve Terradas, 5, 08860 Castelldefels. clemens@ma4.upc.edu

³Departament de Matemàtica Aplicada IV, Universitat Politècnica de Catalunya, C/ Jordi Girona 1-3, Mòdul C3, Campus Nord, 08034 Barcelona. etramuns@ma4.upc.edu

Preprint submitted to International Journal of Computational Geometry and Applications

of obtuse angles among all point sets S with n points is between $\frac{n^3}{18} - O(n^2)$ and $\frac{2n^3}{27} - O(n^2)$. In this note we prove that point sets S in convex position determine at least $\frac{2n^3}{27} - o(n^3)$ obtuse angles. Interestingly, this matches (up to sub-cubic terms) the upper bound example from Ref. 2, which also is a point set in convex position. We conjecture that $\frac{2n^3}{27}$ is indeed the right order of magnitude for the minimum number of obtuse angles (that is to say, every set of n points in general position in the plane determines at least $\frac{2n^3}{27} - o(n^3)$ obtuse angles).

We also relate the number of obtuse angles in a point set S with the *rectilinear crossing number* $\overline{cr}(S)$, which is defined as the number of crossings of edges in a drawing of the complete graph K_n with vertex set S and edges drawn as straight line segments. Four points of S define a crossing if and only if they are in convex position. Therefore, $\overline{cr}(S)$ equals the number of convex quadrilaterals with vertices in S . Point sets in convex position are characterized as the point sets that maximize $\overline{cr}(S)$, and an upper bound of $\binom{n}{4}$ on $\overline{cr}(S)$ is obvious. The rectilinear crossing number of K_n , $\overline{cr}(n)$, is the minimum of $\overline{cr}(S)$ among all sets S of n points. The current best lower bound on $\overline{cr}(n)$ is $\frac{277}{729} \binom{n}{4} + \Theta(n^3)$, see Ref. 1, and there are point sets S that only have $\overline{cr}(S) = 0.380473 \binom{n}{4} + \Theta(n^3)$, see Ref. 4. We show that point sets S whose crossing number is not too large, at most $\frac{2}{3} \binom{n}{4}$, have more than $\frac{n^3}{18}$ obtuse angles.

Results and Proofs

Theorem 0.1. *Every set S of n points in convex position in the plane determines at least $\frac{2n^3}{27} - o(n^3)$ obtuse angles.*

Proof. First we consider the case when n is a prime number; the case when n is not a prime number will be treated at the end of the proof. We label the points of S from 0 to $n - 1$ in counter-clockwise order. For three points $a, b, c \in S$ in counter-clockwise order, we say that the angle abc at point b is of class (i, j) if the open halfplane bounded by the line through points a and b , and not containing point c contains i points of S , and if the open halfplane bounded by the line through points b and c , and not containing point a contains j points of S ; see Figure 1. (Then ab is an i -edge and bc is a j -edge.) Hence each angle defined by S belongs to some class (i, j) , where $0 \leq i + j \leq n - 3$. For i, j fixed, $i \neq j$, we consider the polygon P that starts at point 0, visits points of S in counter-clockwise order, alternately skipping i points and j points of S , until it returns to point 0 the second time. Three steps of such a polygonal path of P are shown in Figure 1. Note that the polygon P is self-intersecting and can visit vertices more than once. An example for $n = 7$ points is shown in Figure 2. The following claim implies that P is well-defined, that is, P indeed returns to point 0.

- *Claim:* Each angle of class (i, j) and each angle of class (j, i) are encountered exactly once in P .




Figure 1: Alternately skipping i and j points in the polygonal path for class (i, j) .




Figure 2: Left: The polygon P for $n = 7, i = 1$, and $j = 2$. P visits the vertices in order $0, 2, 5, 0, 3, 5, 1, 3, 6, 1, 4, 6, 2, 4, 0$. Right: The corresponding polygon P' visits the vertices in order $0, 5, 3, 1, 6, 4, 2, 0$.

We modify P to obtain a new polygon P' by pairing two consecutive steps of P which skip i points and j points respectively; that is, we now move from a point m to point $m + i + j + 2 \pmod n$. Since n is a prime number, each non-zero element of the additive group \mathbb{Z}_n is a generator of the group; in particular also $i + j + 2$. This implies that P' returns to the starting point 0 after it visited each point of $S \setminus \{0\}$ exactly once. We now retrieve the original polygon P by splitting the paired steps into steps skipping alternately i points and j points. It follows that each point of S is visited twice in P , and each angle of class (i, j) and each angle of class (j, i) are encountered exactly once in P .

- *Claim:* The rotation number, see Ref. 5, of the polygon P is $i + j + 2$.
The rotation number measures how many times the polygon turns around. Note that the underlying point set is in convex position and all steps are done in counter-clockwise order. The polygon visits each vertex m twice; from a point m the polygonal path continues once to point $m + i + 1$

mod n , and once to point $m + j + 1 \pmod n$; in total the path advances $i + j + 2$ points from m . Hence, summing over all n vertices, we count $(i + j + 2)n$ steps between consecutive points of the point set in counter-clockwise order. n steps between consecutive points describe one full turn. Thus the rotation number is $i + j + 2$.

- *Claim:* At least $2n - 3(i + j + 2)$ angles of the $2n$ angles of classes (i, j) and (j, i) encountered in P are obtuse.

For the sake of contradiction, suppose that P contains less than $2n - 3(i + j + 2)$ obtuse angles. Then, P contains more than $3(i + j + 2)$ acute or right angles. By an averaging argument, at least one of the $i + j + 2$ full turns of the polygon contains more than three acute or right angles. But this is not possible, unless P contains four right angles forming a 4-cycle, which contradicts the fact that n is a prime number.

Hence, each pair of classes (i, j) and (j, i) of angles, with $i \neq j$, contains at least $2n - 3(i + j + 2)$ obtuse angles. Summing over all possible values i, j we thus get the lower bound on the number of obtuse angles in S

$$\frac{1}{2} \sum_{i=0}^{\lfloor \frac{2n}{3} - 2 \rfloor} \sum_{j=0, j \neq i}^{\lfloor \frac{2n}{3} - 2 - i \rfloor} (2n - 3(i + j + 2)) = \frac{2n^3}{27} - O(n^2).$$

It remains to consider the case when n is not a prime number. In this case it suffices to only count the number of obtuse angles in a subset S' of S consisting of n_p points, where n_p is the largest prime number smaller than n . Since $n_p > n - o(n)$, see e.g. Ref. 6, we get the lower bound on the number of obtuse angles in S

$$\frac{2(n - o(n))^3}{27} - O(n^2) = \frac{2n^3}{27} - o(n^3).$$

□

Proposition 0.2. *Every set S of n points in general position in the plane with rectilinear crossing number $\overline{cr}(S)$ determines at least $\frac{n^3}{12} - \frac{\overline{cr}(S)}{n-3} - O(n^2)$ obtuse angles.*

Proof. We first remark that the number of right angles formed by S is negligible for our purpose. In fact, it is enough to observe that no edge spanned by S is incident to more than two right angles, due to the general position assumption. Hence we upper bound the number of right angles by $2\binom{n}{2}$. Each 4-tuple of points in convex position forms at least one obtuse angle or four right angles; and each 4-tuple of points not in convex position forms at least two obtuse angles. Thus, the total number of obtuse angles in S is at least $\frac{(\overline{cr}(S) - 2\binom{n}{2}/4) \cdot 1 + (\binom{n}{4} - \overline{cr}(S)) \cdot 2}{n-3}$, where we divide by $n - 3$ because each obtuse angle is counted $n - 3$ times. Simplifying gives the claimed bound. □

Acknowledgements

Ruy Fabila-Monroy supported by Conacyt of Mexico, Grant 153984. Clemens Huemer supported by projects MEC MTM2012-30951 and Gen. Cat. DGR 2009SGR1040, 2014SGR46 and ESF EUROCORES programme EuroGIGA, CRP ComPoSe: grant EUI-EURC-2011-4306 for Spain. Eulàlia Tramuns supported by MTM2011-28800-C02-01 from Spanish MEC.

References

- [1] B.M. Ábrego, M. Cetina, S. Fernández-Merchant, J. Leaños, G. Salazar, On $(\leq k)$ -edges, crossings, and halving lines of geometric drawings of K_n , *Discrete and Computational Geometry* **48** (2012), 192–215.
- [2] J.H. Conway, H.T. Croft, P. Erdős, M.J.T. Guy, On the distribution of values of angles determined by coplanar points, *Journal of the London Mathematical Society* (2) **19** (1979), 137–143.
- [3] P. Erdős, Z. Füredi, The greatest angle among n points in the d -dimensional Euclidean space, *Annals of Discrete Mathematics* **17** (1983), 275–283.
- [4] R. Fabila-Monroy, J. López, Computational search of small point sets with small rectilinear crossing number, arXiv:1403.1288, (2014).
- [5] B. Grünbaum, G.C. Shephard, Rotation and winding numbers for planar polygons and curves, *Transactions of the American Mathematical Society* **322** (1990), 169–187.
- [6] R. Guy, Gaps between primes. Twin primes, in *Unsolved problems in number theory*, eds. K. Bencsáth and P.R. Halmos (Third Edition, Springer, New York, 2004), 31–39.