

BASS NUMBERS OF LOCAL COHOMOLOGY OF COVER IDEALS OF GRAPHS

JOSEP ÀLVAREZ MONTANER AND FATEMEH SOHRABI

ABSTRACT. We develop splitting techniques to study Lyubeznik numbers of cover ideals of graphs which allow us to describe them for large families of graphs including forests, cycles, wheels and cactus graphs. More generally we are able to compute all the Bass numbers and the shape of the injective resolution of local cohomology modules by considering the connected components of all the induced subgraphs. Indeed our method gives us a very simple criterion for the vanishing of these local cohomology modules in terms of the number of connected components of the induced subgraphs.

1. INTRODUCTION

Let $G = (V_G, E_G)$ be a finite graph in the set of vertices $V_G = \{x_1, \dots, x_n\}$ and the set of edges E_G . We will assume that the graph is simple so no multiple edges between vertices or loops are allowed. In order to study such a graph from an algebraic point of view one may associate a monomial ideal in the polynomial ring $R = \mathbb{K}[x_1, \dots, x_n]$, where \mathbb{K} is a field. There are several ways to do so but the most used in the literature are:

- **Edge ideal:** $I(G) = (x_i x_j \mid \{x_i, x_j\} \in E_G)$.
- **Cover ideal:** $J(G) = \bigcap_{\{x_i, x_j\} \in E_G} (x_i, x_j)$.

That is, the edges of the graph describe the generators of the edge ideal and the primary components of the cover ideal. Another approach would be through the Stanley-Reisner theory that associates a squarefree monomial ideal to a simplicial complex (see [11], [32]). In our case, $I(G)$ is the Stanley-Reisner ideal associated with the simplicial complex

$$\Delta(I(G)) := \{F \subseteq V_G \mid e \not\subseteq F, \forall e \in E_G\}$$

of independent sets of vertices. On the other hand, $J(G)$ is the Stanley-Reisner ideal associated with the simplicial complex $\Delta(J(G))$ whose facets are complements of edges and we have that $\Delta(I(G))$ is the Alexander dual of $\Delta(J(G))$. Although we are just going to use the terminology from cover and edge ideals, we encourage the interested reader to follow their own preferences.

2010 *Mathematics Subject Classification.* Primary 13D45, 13C11, 05C38, 05C40.

Key words and phrases. Local cohomology, injective resolution, graphs, connectivity.

The first author was partially supported by Generalitat de Catalunya 2017SGR-932 project and Spanish Ministerio de Economía y Competitividad MTM2015-69135-P. He is also a member of the Barcelona Graduate School of Mathematics (BGSMath, MDM-2014-0445).

A common theme in combinatorial commutative algebra has been to understand graph theoretic properties of G from the algebraic properties of the associated ideal and vice versa. For instance, a lot of attention has been paid to the study of free resolutions of edge ideals and its associated invariants such as Betti numbers, projective dimension or Castelnuovo-Mumford regularity. Notice that edge ideals are a very particular class of squarefree monomial ideals so one can use some of the techniques available in this context such as Hochster's formula [20], splitting techniques [14, 15] or discrete Morse theory [10]. Despite these efforts, a full description of these algebraic invariants is only known for a few families of graphs.

The aim of this paper is to study Bass numbers of local cohomology modules supported on cover ideals of graphs. The choice of cover ideals instead of edge ideals is because free resolutions and local cohomology modules of any squarefree monomial ideal are related via Alexander duality (see [31], [28],[29], [8]) and, in the particular case we are considering, edge and cover ideals are Alexander dual to each other. Moreover it seems more natural to use the primary decomposition of an ideal if we want to use the Mayer-Vietoris sequence to study local cohomology modules. We remark that the study of Lyubeznik numbers of edge ideals of graphs should be of independent interest because no relations are known between the Lyubeznik numbers of a squarefree monomial ideal and its Alexander dual.

In order to compute the Bass numbers of local cohomology modules of any squarefree monomial ideal we may refer to the work of K. Yanagawa [34] or the work of the first author with his collaborators in [1], [2], [8]. Indeed, one can use the computational algebra system `Macaulay 2` [18] to compute them as it has been shown in [6]. We point out that, using the restriction functor, we may just reduce to the case of studying Bass numbers with respect to the homogeneous maximal ideal, which are also known as Lyubeznik numbers [27].

The methods presented in [8] may seem quite appropriate for the case of cover ideals of graphs. Namely, in order to compute Lyubeznik numbers, one has to describe the linear strands of the free resolution of the corresponding edge ideal and compute the homology groups of a complex of \mathbb{K} -vector spaces associated to these linear strands. However, even though one may find some explicit free resolutions of edge ideals of graphs in the literature, it seems quite complicated to give closed formulas for the Lyubeznik numbers even for simple families of graphs.

In this paper we shift gears and we present some splitting techniques that would allow us to compute the Lyubeznik numbers of large families of graphs without any previous description of its local cohomology modules or equivalently, the free resolution of the corresponding edge ideals. The idea behind these splitting techniques is to relate the Lyubeznik numbers of our initial graph to the Lyubeznik numbers of the subgraph obtained by removing a vertex. Indeed, the Lyubeznik table remains invariant when we remove a whisker or even a 3 or 4-cycle. Moreover we can control the Lyubeznik table when we remove degree two vertices or a dominating vertex. To compute all the Lyubeznik numbers of any given graph in a fixed number of vertices is out of the scope of this work but we can reduce enormously the number of cases that we have to consider by a simple inspection of the shape of the graph. More generally, we can compute all the Bass numbers of local cohomology modules just considering subgraphs of our initial graph. In particular we can describe the linear strands of the injective resolution of these modules. The structure of these injective resolutions depend on the number of connected components of the corresponding subgraphs. Quite nicely, we

deduce a vanishing criterion for local cohomology modules depending of these connected components of the subgraphs.

We should mention that, in general, Lyubeznik numbers depend on the characteristic of the base field. However, all the methods we develop here are independent of the characteristic, meaning that the Lyubeznik numbers of a graph will depend on the characteristic if and only if the Lyubeznik numbers of the graph obtained after removing a vertex also depend on the characteristic.

The organization of this paper is as follows. In Section 2 we introduce all the basics on local cohomology supported on squarefree monomial ideals and its injective resolution. In particular we introduce Bass numbers and how to describe them using the graded pieces of the composition of local cohomology modules. Since we can always reduce to the case of Lyubeznik numbers we briefly recall in Subsection 2.1 its definition and the main properties we are going to use throughout this work. In Subsection 2.2 we review the relation between Lyubeznik numbers and linear strands of the Alexander dual ideal. Finally, in Subsection 2.3 we propose the notion of MV-splitting (see Definition 2.8) together with an application of the long exact sequence of local cohomology modules (see Discussion 2.12) that will be crucial later on. The reason of working in the general framework of squarefree monomial ideals is that, even though we want to study cover ideals of graphs, we will have to leave this context when applying these techniques. Moreover, all these splitting methods could be also applied for any squarefree monomial ideal.

In Section 3 we focus on the study of Lyubeznik numbers of cover ideals of graphs. Our first result is Theorem 3.3 in which we describe the Lyubeznik table associated to the cover ideal of a simple connected graph for which the MV-splitting satisfies some extra conditions. These conditions are naturally satisfied when we consider splitting vertices and thus we specialize to this case. In Proposition 3.5 we prove that the Lyubeznik table remains invariant after removing vertices of degree one. In Proposition 3.6 we prove that the Lyubeznik table is also invariant if we remove a handle, which is a 3 or 4-cycle having a degree two vertex. More generally, we describe in Proposition 3.7 and Corollary 3.8 the Lyubeznik table of any graph as long as we find degree two splitting vertices. In this way we can apply recursion to reduce the computation to the case of a smaller graph.

In Subsection 3.2 we apply these splitting techniques to compute the Lyubeznik table of some families of examples. We prove that trees have trivial Lyubeznik table and we deduce a formula for the case of forests. Any cone of a graph, for example a wheel, also has trivial Lyubeznik table. The results on degree two vertices allow us to compute the case of cycles and, more generally, the family of graphs obtained by joining cycles in such a way that we can still find degree two vertices that we can remove in order to simplify the graph. Indeed, after removing whiskers and handles we may consider cycles joined by paths or sharing edges. Such an example would be the case of cactus graphs or cycles with chords.

In Section 4 we study all the Bass numbers of the cover ideal of a graph by considering the Lyubeznik numbers of the corresponding subgraphs. In Subsection 4.1 we pay attention to a class of graphs (that include forests and Cohen-Macaulay graphs) whose local cohomology modules have a linear injective resolution. In particular we give a closed formula for these Bass numbers in Theorem 4.4. Quite surprisingly we provide in Proposition 4.6 a vanishing criterion

for the local cohomology modules in terms of the number of connected components of the subgraphs. Using Alexander duality it also gives a formula for the projective dimension of the edge ideal of such a graph. We also study the injective resolution of local cohomology modules of graphs obtained by joining cycles in Subsection 4.2. Finally, we also provide a vanishing criterion for the local cohomology modules associated to the corresponding subgraphs in Proposition 4.12 and Proposition 4.13.

2. BASS NUMBERS OF LOCAL COHOMOLOGY MODULES

Throughout this section we will assume the general framework of a squarefree monomial ideals in a polynomial ring $R = \mathbb{K}[x_1, \dots, x_n]$ with coefficients over a field \mathbb{K} . Namely, a squarefree monomial ideal $J \subseteq R$ is generated by monomials of the form $\mathbf{x}^\alpha := x_1^{\alpha_1} \cdots x_n^{\alpha_n}$, where $\alpha = (a_1, \dots, a_n) \in \{0, 1\}^n$. Its minimal primary decomposition is given in terms of face ideals $\mathfrak{p}_\alpha := \langle x_i \mid a_i \neq 0 \rangle$, $\alpha \in \{0, 1\}^n$. For simplicity we will denote the homogeneous maximal ideal $\mathfrak{m} := \mathfrak{p}_\mathbf{1} = (x_1, \dots, x_n)$, where $\mathbf{1} = (1, \dots, 1)$. As usual, we denote $|\alpha| = a_1 + \cdots + a_n$ and $\varepsilon_1, \dots, \varepsilon_n$ will be the standard basis of \mathbb{Z}^n . The **Alexander dual** of the ideal J is the squarefree monomial ideal $J^\vee \subseteq R$ defined as $J^\vee = (\mathbf{x}^{\alpha_1}, \dots, \mathbf{x}^{\alpha_s})$ associated to the minimal primary decomposition $J = \mathfrak{p}_{\alpha_1} \cap \cdots \cap \mathfrak{p}_{\alpha_s}$.

Let $\mathbb{Z}^\alpha \subseteq \mathbb{Z}^n$ be the coordinate space spanned by $\{\varepsilon_i \mid a_i = 1\}$, $\alpha \in \{0, 1\}^n$. The **restriction** of R to the face ideal $\mathfrak{p}_\alpha \subseteq R$ is the \mathbb{Z}^α -graded \mathbb{K} -subalgebra of R

$$R_{\mathfrak{p}_\alpha} := \mathbb{K}[x_i \mid a_i = 1].$$

Let $J = \mathfrak{p}_{\alpha_1} \cap \cdots \cap \mathfrak{p}_{\alpha_s}$ be the minimal primary decomposition of a squarefree monomial ideal $J \subseteq R$. Then, the restriction of J to the face ideal \mathfrak{p}_α is the squarefree monomial ideal

$$J_{\mathfrak{p}_\alpha} = \bigcap_{\alpha_j \leq \alpha} \mathfrak{p}_{\alpha_j} \subseteq R_{\mathfrak{p}_\alpha}.$$

Moreover, the restriction of a local cohomology module is

$$[H_J^r(R)]_{\mathfrak{p}_\alpha} = H_{J_{\mathfrak{p}_\alpha}}^r(R_{\mathfrak{p}_\alpha})$$

Roughly speaking, restriction gives us a functor that plays the role of the localization functor. For details and further considerations we refer to [29].

A key fact in its study is that local cohomology modules $H_J^r(R)$ supported on monomial ideals are \mathbb{Z}^n -graded modules. Indeed, these modules satisfy some nice properties since they fit, modulo a shifting by $\mathbf{1}$, into the category of **straight** (resp. **1-determined**) modules introduced by K. Yanagawa [34] (resp. E. Miller [29]). In what follows we are going to introduce the basic notions that we are going to use in this work. Most of them can be found in textbooks such as [11] and [30] or the lecture notes [5].

In order to give a module structure to the straight module $H_J^r(R)$ we have to describe:

- The graded pieces $[H_J^r(R)]_{-\alpha}$ for all $\alpha \in \{0, 1\}^n$.
- The multiplication morphisms: $\cdot x_i : [H_J^r(R)]_{-\alpha} \longrightarrow [H_J^r(R)]_{-(\alpha - \varepsilon_i)}$.

This structure has been described by N. Terai [33] and M. Mustařă [31] in terms of some simplicial complexes associated to the monomial ideal J . The approach considered in [7] gives an interpretation in terms of the components appearing in the minimal primary decomposition of J which will be more convenient for our purposes.

Let \mathcal{P}_J be the partially ordered set consisting of the sums of ideals in the minimal primary decomposition of J ordered by reverse inclusion. Namely, if $J = \mathfrak{p}_{\alpha_1} \cap \cdots \cap \mathfrak{p}_{\alpha_s}$ is the minimal primary decomposition we have that any ideal $J_p \in \mathcal{P}_J$ is a certain sum $J_p = \mathfrak{p}_{\alpha_{i_1}} + \cdots + \mathfrak{p}_{\alpha_{i_j}}$ and, since the sum of face ideals is a face ideal we have that $J_p = \mathfrak{p}_\alpha$ for some $\alpha \in \{0, 1\}^n$. In what follows we will just denote by \mathfrak{p}_α , or simply α , the elements of \mathcal{P}_J .

Let $1_{\mathcal{P}_J}$ be a terminal element that we add to the poset. To any $\alpha \in \mathcal{P}_J$ we may consider the **order complex** associated to the subposet $(\alpha, 1_{\mathcal{P}_J}) := \{z \in \mathcal{P}_J \mid \alpha < z < 1_{\mathcal{P}_J}\}$ and the dimensions of the reduced simplicial homology groups

$$m_{r,\alpha} := \dim_{\mathbb{K}} \tilde{H}_{|\alpha|-r-1}((\alpha, 1_{\mathcal{P}_J}); \mathbb{K}).$$

Then, the graded pieces of the local cohomology modules of J can be described as follows:

$$(2.1) \quad [H_J^r(R)]_{-\alpha} = \bigoplus_{\alpha \in \mathcal{P}_J} [H_{\mathfrak{p}_\alpha}^{|\alpha|}(R)^{m_{r,\alpha}}]_{-\alpha}$$

The category of straight modules is a category with enough injective modules. Indeed, the indecomposable injective objects are the shifted injective envelopes $E_\alpha := {}^*E_R(R/\mathfrak{p}_\alpha)(\mathbf{1})$, $\alpha \in \{0, 1\}^n$, and every graded injective module is isomorphic to a unique (up to order) direct sum of indecomposable injectives. It follows that the **minimal \mathbb{Z}^n -graded injective resolution** of a local cohomology module $H_J^r(R)$, is an exact sequence:

$$\mathbb{I}_\bullet(H_J^r(R)) : \quad 0 \longrightarrow H_J^r(R) \longrightarrow I_0 \xrightarrow{d^0} I_1 \xrightarrow{d^1} \cdots \longrightarrow I_m \xrightarrow{d^m} 0,$$

where the p -th term is

$$I_p = \bigoplus_{\alpha \in \mathbb{Z}^n} E_\alpha^{\mu_p(\mathfrak{p}_\alpha, H_J^r(R))}$$

and the invariants defined by $\mu_p(\mathfrak{p}_\alpha, H_J^r(R))$ are the **Bass numbers** of $H_J^r(R)$. Given an integer ℓ , the **ℓ -linear strand** of $\mathbb{I}_\bullet(H_J^r(R))$ is the complex:

$$\mathbb{I}_\bullet^{<\ell>}(H_J^r(R)) : \quad 0 \longrightarrow I_0^{<\ell>} \longrightarrow I_1^{<\ell>} \longrightarrow \cdots \longrightarrow I_m^{<\ell>} \longrightarrow 0,$$

where

$$I_p^{<\ell>} = \bigoplus_{|\alpha|=p+\ell} E_\alpha^{\mu_p(\mathfrak{p}_\alpha, H_J^r(R))},$$

Remark 2.1. The (\mathbb{Z}^n -graded) Bass numbers coincide with the usual Bass numbers in the minimal injective resolution of $H_J^r(R)$ as it was proved by S. Goto and K. I. Watanabe in [17]. Indeed they provided a method to compute the Bass numbers with respect to any prime ideal. Namely, given any prime ideal $\mathfrak{p} \in \text{Spec}R$, let \mathfrak{p}_α be the largest face ideal contained in \mathfrak{p} . If $\text{ht}(\mathfrak{p}/\mathfrak{p}_\alpha) = s$ then $\mu_p(\mathfrak{p}_\alpha, H_J^r(R)) = \mu_{p+s}(\mathfrak{p}, H_J^r(R))$.

The Bass numbers of straight modules, and local cohomology modules in particular, were already studied and described in [34]. The approach that we will use in this work is using the

graded pieces of the composition of local cohomology modules. Namely, using [8, Corollary 3.6] (see also [4]), we have:

Proposition 2.2. *Let $J \subseteq R$ be a squarefree monomial ideal and $\mathfrak{p}_\alpha \subseteq R$ be a face ideal, $\alpha \in \{0, 1\}^n$. Then, the Bass numbers of the local cohomology module $H_J^r(R)$ with respect to \mathfrak{p}_α are*

$$\mu_p(\mathfrak{p}_\alpha, H_J^r(R)) = \dim_{\mathbb{K}}[H_{\mathfrak{p}_\alpha}^p(H_J^r(R))]_{-\alpha}.$$

In particular, the Bass numbers with respect to the homogeneous maximal ideal \mathfrak{m} are

$$\mu_p(\mathfrak{m}_\alpha, H_J^r(R)) = \dim_{\mathbb{K}}[H_{\mathfrak{m}}^p(H_J^r(R))]_{-1}.$$

Remark 2.3. Bass numbers behave well with respect to the restriction functor so we may always assume that the face ideal \mathfrak{p}_α is the maximal ideal. Namely, we have

$$\mu_p(\mathfrak{p}_\alpha, H_J^r(R)) = \mu_p(\mathfrak{p}_\alpha R_{\mathfrak{p}_\alpha}, [H_J^r(R)]_{\mathfrak{p}_\alpha}).$$

2.1. Lyubeznik numbers. In the seminal works of C. Huneke and R. Y. Sharp [22] and G. Lyubeznik [27] it is proven that the Bass numbers of local cohomology modules are all finite. This prompted G. Lyubeznik to introduce a new set of invariants defined as follows:

Let A be a noetherian local ring that admits a surjection from an n -dimensional regular local ring (R, \mathfrak{m}) containing its residue field \mathbb{K} , and $J \subseteq R$ be the kernel of the surjection. Then, the Bass numbers

$$\lambda_{p,i}(A) := \mu_p(\mathfrak{m}, H_J^{n-i}(R))$$

depend only on A , i and p , but not on the choice of R or the surjection $R \rightarrow A$. More generally, all the Bass numbers $\mu_p(\mathfrak{p}, H_J^{n-i}(R))$ are invariants of the local ring A as it was proved later on in [3]. Bass numbers behave well with respect to completion so we may always assume that A is a quotient of a formal power series ring R . Considering a squarefree monomial ideal as an ideal in the polynomial or the formal power series ring makes no difference since the Bass numbers of the corresponding local cohomology modules coincide. It is for this reason that we will keep considering, for simplicity, just the case of R being a polynomial ring.

Lyubeznik numbers satisfy $\lambda_{d,d}(A) \neq 0$ and $\lambda_{p,i}(A) \neq 0$ implies $0 \leq p \leq i \leq d$, where $d = \dim A$. A way to collect these invariants is by means of the so-called **Lyubeznik table**:

$$\Lambda(A) = \begin{pmatrix} \lambda_{0,0} & \cdots & \lambda_{0,d} \\ & \ddots & \vdots \\ & & \lambda_{d,d} \end{pmatrix}$$

and we say that the Lyubeznik table is **trivial** if $\lambda_{d,d} = 1$ and the rest of these invariants vanish.

The highest Lyubeznik number $\lambda_{d,d}(A)$ has an interesting interpretation in terms of the so-called **Hochster-Huneke graph** defined in [21], which is the graph whose vertices are the top-dimensional minimal primes of A and we have an edge between two vertices \mathfrak{p} and \mathfrak{q} if and only if $\text{ht}(\mathfrak{p} + \mathfrak{q}) = 1$. The following result was proved by G. Lyubeznik [26] when \mathbb{K} is a positive characteristic field, and a characteristic-free proof was given by W. Zhang [35]. To avoid technicalities in the statement of the result we will restrict ourselves to the case of squarefree monomial ideals in a polynomial ring.

Theorem 2.4. *Let $I \subseteq R = \mathbb{K}[x_1, \dots, x_n]$ be a squarefree monomial ideal and $A = R/I$. The highest Lyubeznik number $\lambda_{d,d}(A)$ equals the number of connected components of the Hochster-Huneke graph of A .*

Another property that we are going to use in this work is the following Thom-Sebastiani type formula for the case of squarefree monomial ideals that was proved in [9].

Proposition 2.5. *Let $I \subseteq R = \mathbb{K}[x_1, \dots, x_m]$ and $J \subseteq S = \mathbb{K}[y_1, \dots, y_n]$ be squarefree monomial ideals in two disjoint sets of variables and set $T = \mathbb{K}[x_1, \dots, x_m, y_1, \dots, y_n]$. Then, the Lyubeznik numbers of $T/IT \cap JT$ have the following form:*

i) If either the height of I or the height of J is 1, then $T/IT \cap JT$ has trivial Lyubeznik table.

ii) If both the height of I and the height of J are ≥ 2 , then we have:

$$\begin{aligned} \lambda_{p,i}(T/IT \cap JT) &= \lambda_{p,i}(T/IT) + \lambda_{p,i}(T/JT) + \sum_{\substack{q+r=p+\dim T \\ j+k=i+\dim T-1}} \lambda_{q,j}(T/IT) \lambda_{r,k}(T/JT) \\ &= \lambda_{p-n,i-n}(R/I) + \lambda_{p-m,i-m}(S/J) + \sum_{\substack{q+r=p \\ j+k=i-1}} \lambda_{q,j}(R/I) \lambda_{r,k}(S/J). \end{aligned}$$

The following particular case will be very useful later on.

Corollary 2.6. *Let $I \subseteq R = \mathbb{K}[x_1, \dots, x_n]$ be a squarefree monomial ideal admitting a decomposition $I = I_1 \cap \dots \cap I_c$ in disjoint sets of variables such that $\dim R/I_j = d$ and $\Lambda(R/I_j)$ is trivial for $j = 1, \dots, c$. Then*

$$\lambda_{d-2k,d-k}(R/I) = \binom{c}{k+1} \quad \text{for } k = 0, \dots, c-1$$

and the rest of Lyubeznik numbers are zero.

Proof. First we notice that the matrices $\Lambda(R/I_i)$ have the same size for all i . In the case that $c = 2$ we have, using Proposition 2.5, $\lambda_{d,d} = 2$ and $\lambda_{d-2,d-1} = 1$. Then we proceed using induction on the number of components. \square

A general formula for the case of c disjoint sets of variables could be worked out but we will just focus on finding the smallest integer i for which there exist p such that $\lambda_{p,i}(R/I) \neq 0$.

Corollary 2.7. *Let $I \subseteq R = \mathbb{K}[x_1, \dots, x_n]$ be a squarefree monomial ideal admitting a decomposition $I = I_1 \cap \dots \cap I_c$ in disjoint sets of variables. Let i_j be the smallest integer for which there exist p such that $\lambda_{p,i_j}(R/I_j) \neq 0$ for $j = 1, \dots, c$. Then, the smallest i for which there exist p such that $\lambda_{p,i}(R/I) \neq 0$ is $i = (i_1 + \dots + i_c) + (c-1)$.*

Proof. In the case that $c = 2$, let q and r be integers such that $\lambda_{q,i_1}(R/I_1) \neq 0$, $\lambda_{r,i_2}(S/I_2) \neq 0$. Then $\lambda_{q+r,i_1+i_2+1}(T/I_1T + I_2T) \neq 0$ using Proposition 2.5 and it gives the smallest integer i satisfying this property. Then we proceed using induction on the number of components. \square

2.2. Local cohomology modules and free resolutions. A way to interpret Lyubeznik numbers for the case of squarefree monomial ideals is in terms of the linear strands of the free resolution of the Alexander dual of the ideal. This approach was given in [8] and further developed in [9] and we will briefly recall it here.

Let J^\vee be the Alexander dual of a squarefree monomial ideal $J \subseteq R$. Its minimal \mathbb{Z} -graded **free resolution** is an exact sequence of free \mathbb{Z} -graded R -modules:

$$\mathbb{L}_\bullet(J^\vee) : 0 \longrightarrow L_m \xrightarrow{d_m} \cdots \longrightarrow L_1 \xrightarrow{d_1} L_0 \longrightarrow J^\vee \longrightarrow 0$$

where the j -th term is of the form

$$L_j = \bigoplus_{\ell \in \mathbb{Z}} R(-\ell)^{\beta_{j,\ell}(J^\vee)},$$

and the matrices of the morphisms $d_j : L_j \rightarrow L_{j-1}$ do not contain invertible elements. The \mathbb{Z} -graded **Betti numbers** of J^\vee are the invariants $\beta_{j,\ell}(J^\vee)$. Given an integer r , the r -**linear strand** of $\mathbb{L}_\bullet(J^\vee)$ is the complex:

$$\mathbb{L}_\bullet^{\langle r \rangle}(J^\vee) : 0 \longrightarrow L_{n-r}^{\langle r \rangle} \xrightarrow{d_{n-r}^{\langle r \rangle}} \cdots \longrightarrow L_1^{\langle r \rangle} \xrightarrow{d_1^{\langle r \rangle}} L_0^{\langle r \rangle} \longrightarrow 0,$$

where

$$L_j^{\langle r \rangle} = R(-j-r)^{\beta_{j,j+r}(J^\vee)},$$

and the differentials $d_j^{\langle r \rangle} : L_j^{\langle r \rangle} \rightarrow L_{j-1}^{\langle r \rangle}$ are the corresponding components of d_j .

We point out that these differentials can be described using the so-called **monomial matrices** introduced by E. Miller in [29] (see also [30]). These are matrices with scalar entries that keep track of the degrees of the generators of the summands in the source and the target. Now we construct a complex of \mathbb{K} -vector spaces

$$\mathbb{F}_\bullet^{\langle r \rangle}(J^\vee)^* : 0 \longleftarrow \underbrace{\mathbb{K}^{\beta_{n-r,n}(J^\vee)}}_{\text{deg } 0} \longleftarrow \cdots \longleftarrow \underbrace{\mathbb{K}^{\beta_{1,1+r}(J^\vee)}}_{\text{deg } n-r-1} \longleftarrow \underbrace{\mathbb{K}^{\beta_{0,r}(J^\vee)}}_{\text{deg } n-r} \longleftarrow 0$$

where the morphisms are given by the transpose of the corresponding monomial matrices and thus we reverse the indices of the complex. Then, the Lyubeznik numbers are described by means of the homology groups of these complexes. Namely, the result given in [8, Corollary 4.2] is the following characterization

$$(2.2) \quad \lambda_{p,n-r}(R/J) = \dim_{\mathbb{K}} H_p(\mathbb{F}_\bullet^{\langle r \rangle}(J^\vee)^*).$$

2.3. Mayer-Vietoris splitting. A successful technique used in the study of free resolutions of monomial ideals was developed by S. Eliahou and M. Kervaire in [14] and refined by C. Francisco, H. T. Hà and A. Van Tuyl in [15] under the terminology of **splittings** of monomial ideals and **Betti splittings** respectively.

An analogous technique can be used to study local cohomology modules.

Definition 2.8. Let $J \subseteq R$ be a squarefree monomial ideal. We say that the decomposition $J = L \cap K$ is a MV-splitting if the Mayer-Vietoris sequence

$$\cdots \longrightarrow H_{L+K}^r(R) \longrightarrow H_L^r(R) \oplus H_K^r(R) \longrightarrow H_J^r(R) \longrightarrow H_{L+K}^{r+1}(R) \longrightarrow \cdots$$

splits into short exact sequences

$$0 \longrightarrow H_L^r(R) \oplus H_K^r(R) \longrightarrow H_J^r(R) \longrightarrow H_{L+K}^{r+1}(R) \longrightarrow 0$$

for all r .

Remark 2.9. Using Alexander duality, we have that this notion is equivalent to the fact that J^\vee admits a Betti splitting $J^\vee = L^\vee + K^\vee$ in the sense of [15], which means that the \mathbb{Z}^n -graded Betti numbers satisfy

$$\beta_{i,\alpha}(J^\vee) = \beta_{i,\alpha}(L^\vee) + \beta_{i,\alpha}(K^\vee) + \beta_{i-1,\alpha}(L^\vee \cap K^\vee)$$

Certainly we have a MV-splitting if the \mathbb{Z}^n -graded morphisms $H_{L+K}^r(R) \longrightarrow H_L^r(R) \oplus H_K^r(R)$ are zero for all r . Sufficient conditions for this vanishing can be given in terms of the posets of sums of ideals associated to L , K and $L+K$. The following result, which uses the terminology of Equation (2.1), can be understood as a reinterpretation of [15, Theorem 2.3].

Proposition 2.10. *Let $J = L \cap K$ be a decomposition of a squarefree monomial ideal $J \subseteq R$. Consider the posets $\mathcal{P}_L, \mathcal{P}_K$ and \mathcal{P}_{L+K} associated to the primary decompositions of the ideals L, K and $L+K$ respectively. Assume that $m_{r,\alpha}(L+K) \neq 0$ implies $m_{r,\alpha}(L) = m_{r,\alpha}(K) = 0$ for any r and any $\alpha \in \{0, 1\}^n$. Then the decomposition $J = L \cap K$ is a MV-splitting.*

Proof. The assumptions we are considering are telling us that $[H_{L+K}^r(R)]_{-\alpha} \neq 0$ implies $[H_L^r(R)]_{-\alpha} = [H_K^r(R)]_{-\alpha} = 0$ by means of Equation (2.1), and thus the \mathbb{Z}^n -graded morphisms $H_{L+K}^r(R) \longrightarrow H_L^r(R) \oplus H_K^r(R)$ are zero for all r . \square

Corollary 2.11. *Let $J = L \cap K$ be a decomposition of a squarefree monomial ideal $J \subseteq R$. Assume that the posets $\mathcal{P}_L, \mathcal{P}_K$ and \mathcal{P}_{L+K} associated to the primary decompositions of the ideals L, K and $L+K$ have no face ideal in common. Then the decomposition $J = L \cap K$ is a MV-splitting.*

We want to apply these splitting techniques to the study of the composition of local cohomology modules. The following discussion will be crucial in the rest of this work.

Discussion 2.12. The degree - 1 part of the long exact sequence of local cohomology associated to the short exact sequences

$$(2.3) \quad 0 \longrightarrow H_L^r(R) \oplus H_K^r(R) \longrightarrow H_J^r(R) \longrightarrow H_{L+K}^{r+1}(R) \longrightarrow 0$$

obtained in a MV-splitting is

$$(2.4) \quad \begin{aligned} \dots \longrightarrow [H_m^{p-1}(H_{L+K}^{r+1}(R))]_{-1} &\xrightarrow{\partial_{p-1}^r} [H_m^p(H_L^r(R))]_{-1} \oplus [H_m^p(H_K^r(R))]_{-1} \longrightarrow [H_m^p(H_J^r(R))]_{-1} \longrightarrow \\ &\longrightarrow [H_m^p(H_{L+K}^{r+1}(R))]_{-1} \xrightarrow{\partial_p^r} [H_m^{p+1}(H_L^r(R))]_{-1} \oplus [H_m^{p+1}(H_K^r(R))]_{-1} \longrightarrow \dots \end{aligned}$$

Equivalently, it is the long exact sequence of \mathbb{K} -vector spaces whose dimensions are the corresponding Lyubeznik numbers. Namely,

$$(2.5) \quad \begin{aligned} \dots \longrightarrow \mathbb{K}^{\lambda_{p-1,n-r-1}(R/L+K)} &\xrightarrow{\partial_{p-1}^r} \mathbb{K}^{\lambda_{p,n-r}(R/L)} \oplus \mathbb{K}^{\lambda_{p-1,n-r}(R/K)} \longrightarrow \mathbb{K}^{\lambda_{p-1,n-r}(R/J)} \longrightarrow \\ &\longrightarrow \mathbb{K}^{\lambda_{p,n-r-1}(R/L+K)} \xrightarrow{\partial_p^r} \mathbb{K}^{\lambda_{p+1,n-r}(R/L)} \oplus \mathbb{K}^{\lambda_{p+1,n-r}(R/K)} \longrightarrow \dots \end{aligned}$$

Therefore, if we want to compute the Lyubeznik numbers of R/J in terms of the Lyubeznik numbers of R/L , R/K and $R/L + K$, we need to control the connecting morphisms ∂_p^r 's.

Using the methods considered in [8] we may give an interpretation of these differentials in terms of linear strands. First, the short exact sequence (2.3) corresponds to the short exact sequence of complexes of \mathbb{K} -vector spaces

$$0 \longleftarrow \mathbb{F}_{\bullet}^{\langle r \rangle}(L^\vee)^* \oplus \mathbb{F}_{\bullet}^{\langle r \rangle}(K^\vee)^* \longleftarrow \mathbb{F}_{\bullet}^{\langle r \rangle}(J^\vee)^* \longleftarrow \mathbb{F}_{\bullet}^{\langle r+1 \rangle}((L+K)^\vee)^* \longleftarrow 0$$

and the long exact sequence (2.4) corresponds to

$$\begin{aligned} \cdots \longleftarrow H_{p-1}(\mathbb{F}_{\bullet}^{\langle r+1 \rangle}((L+K)^\vee)^*) \xleftarrow{\partial_{p-1}^r} H_p(\mathbb{F}_{\bullet}^{\langle r \rangle}(L^\vee)^*) \oplus H_p(\mathbb{F}_{\bullet}^{\langle r \rangle}(K^\vee)^*) \longleftarrow H_p(\mathbb{F}_{\bullet}^{\langle r \rangle}(J^\vee)^*) \longleftarrow \\ \longleftarrow H_p(\mathbb{F}_{\bullet}^{\langle r+1 \rangle}((L+K)^\vee)^*) \xleftarrow{\partial_p^r} H_{p+1}(\mathbb{F}_{\bullet}^{\langle r \rangle}(L^\vee)^*) \oplus H_{p+1}(\mathbb{F}_{\bullet}^{\langle r \rangle}(K^\vee)^*) \longleftarrow \cdots \end{aligned}$$

3. LYUBEZNIK TABLES OF COVER IDEALS OF GRAPHS

Let $G = (V_G, E_G)$ be a simple finite graph in the set of vertices $V_G = \{x_1, \dots, x_n\}$ and the set of edges E_G . For simplicity we will also assume that G is connected. For a vertex x_i , we consider its **neighbour set** $N(x_i) = \{x_j \in G \mid \{x_i, x_j\} \in E_G\}$. The **degree** of a vertex is the cardinal of its neighbour set.

Let $J(G) \subseteq R$ be the cover ideal of G where $R = \mathbb{K}[x_1, \dots, x_n]$ is a polynomial ring with coefficients in a field \mathbb{K} . In this section we will develop MV-splitting techniques to study the Lyubeznik numbers of $R/J(G)$. To start with, we recall that since $J(G)$ is a pure height two ideal, the dimension of the support of $H_{J(G)}^r(R)$ is strictly smaller than $n - r$ for $r > 2$ (see [1, Theorem 3.8]). Then it follows from [1, Theorem 4.4] that all the entries in the main diagonal of the Lyubeznik table are zero except for the highest Lyubeznik number.

Lemma 3.1. *Let $J(G)$ be the cover ideal of a simple connected graph G . Then, the highest Lyubeznik number is $\lambda_{d,d}(R/J(G)) = 1$*

Proof. The vertices of the Hochster-Huneke graph of $J(G)$ correspond to the edges of G , and the edges of the Hochster-Huneke graph correspond to adjacent edges of G . Therefore, the Hochster-Huneke graph has just one connected component since the graph G is connected. \square

Under these restrictions, the shape of the Lyubeznik table is

$$\Lambda(R/J(G)) = \begin{pmatrix} 0 & \lambda_{0,1} & \cdots & \lambda_{0,d-1} & \lambda_{0,d} \\ & 0 & \cdots & \lambda_{1,d-1} & \lambda_{1,d} \\ & & \ddots & \vdots & \vdots \\ & & & 0 & \lambda_{d-1,d} \\ & & & & 1 \end{pmatrix}$$

In the case that $R/J(G)$ is Cohen-Macaulay we have that the Lyubeznik table is trivial (see [1, Remark 4.2]). Recall that, combining the results in [13] with [16], we have the following characterization of this property.

Proposition 3.2. *Let G be a simple graph. Then the following are equivalent:*

- i) The cover ideal $J(G)$ is Cohen-Macaulay.*
- ii) The edge ideal $I(G)$ has a linear resolution.*
- iii) The complement graph G^c is chordal.*

Free resolutions of edge ideals have been extensively studied over the last years and we may find in the literature several families of Cohen-Macaulay graphs. For example,

- **Complete graphs** K_n [23].
- **Complete bipartite graphs** $K_{n,m}$ and in particular **star graphs** $K_{1,m}$ [23].
- **Ferrers graphs** [12].

The simplest examples of ideals with non-trivial Lyubeznik table are minimal non-Cohen-Macaulay squarefree monomial ideals (see [25]). The unique minimal non-Cohen-Macaulay squarefree monomial ideal of pure height two in $R = \mathbb{K}[x_1, \dots, x_n]$ is the cover ideal of the **complement of a cycle**:

$$J(C_n^c) = (x_1, x_3) \cap \dots \cap (x_1, x_{n-1}) \cap (x_2, x_4) \cap \dots \cap (x_2, x_n) \cap (x_3, x_5) \cap \dots \cap (x_{n-2}, x_n).$$

Its Lyubeznik table is of the form (see [8])

$$\Lambda(R/J(C_n^c)) = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 1 & 0 \\ & 0 & 0 & \cdots & 0 & 0 & 0 \\ & & 0 & & 0 & 0 & 1 \\ & & & \ddots & & 0 & 0 \\ & & & & & \vdots & \vdots \\ & & & & & 0 & 0 \\ & & & & & & 1 \end{pmatrix}$$

To provide a full description of all the possible Lyubeznik tables of cover ideals of graphs is completely out of the scope of this work. Our aim is to introduce some Mayer-Vietoris splitting techniques that will allow us to compute large families of examples. To such purpose we will follow the ideas considered in Discussion 2.12. To start with, we consider the case where $J(G) = L \cap K$ is a MV-splitting with the extra assumption that the Lyubeznik table of R/K is trivial.

Theorem 3.3. *Let $J(G) \subseteq R$ be the cover ideal of a simple connected graph G . Let $J(G) = L \cap K$ be a MV-splitting such that $\Lambda(R/K)$ is trivial. Then:*

- i) If $\Lambda(R/L)$ and $\Lambda(R/L + K)$ are trivial, then $\Lambda(R/J(G))$ is trivial.*
- ii) If $\Lambda(R/L + K)$ is trivial, then $\Lambda(R/J(G)) = \Lambda(R/L)$.*

iii) If $\Lambda(R/L)$ is trivial and

$$\Lambda(R/L + K) = \begin{pmatrix} \lambda'_{0,0} & \cdots & \lambda'_{0,d-1} \\ & \ddots & \vdots \\ & & \lambda'_{d-1,d-1} \end{pmatrix},$$

then the Lyubeznik table of $R/J(G)$ is

$$\Lambda(R/J(G)) = \begin{pmatrix} 0 & \lambda'_{0,0} & \cdots & \lambda'_{0,d-2} & \lambda'_{0,d-1} \\ & 0 & \cdots & \lambda'_{1,d-2} & \lambda'_{1,d-1} \\ & & \ddots & \vdots & \vdots \\ & & & 0 & \lambda'_{d-1,d-1} - 1 \\ & & & & 1 \end{pmatrix}.$$

Proof. Assume that $\Lambda(R/K)$ is trivial and recall that, using Lemma 3.1, the highest Lyubeznik number of the cover ideal of a graph is one. In particular we have $[H_m^{n-2}(H_K^2(R))]_{-1} = \mathbb{K}$ and $[H_m^{n-2}(H_L^2(R))]_{-1} = \mathbb{K}$. Then, for $r = 2$, the long exact sequence 2.4 considered in Discussion 2.12

$$\cdots \longrightarrow [H_m^{n-3}(H_{L+K}^3(R))]_{-1} \xrightarrow{\partial_{n-3}^2} [H_m^{n-2}(H_L^2(R))]_{-1} \oplus [H_m^{n-2}(H_K^2(R))]_{-1} \longrightarrow [H_m^{n-2}(H_{J(G)}^2(R))]_{-1} \longrightarrow 0$$

turns out to be

$$\cdots \longrightarrow \mathbb{K}^{\lambda_{n-3,n-2}(R/L)} \longrightarrow \mathbb{K}^{\lambda_{n-3,n-2}(R/J(G))} \longrightarrow \mathbb{K}^{\lambda_{n-3,n-3}(R/L+K)} \xrightarrow{\partial_{n-3}^2} \mathbb{K} \oplus \mathbb{K} \longrightarrow \mathbb{K} \longrightarrow 0$$

Moreover, for $r > 2$ and any p , the long exact sequence becomes

$$\cdots \longrightarrow \mathbb{K}^{\lambda_{p-1,n-(r+1)}(R/L+K)} \xrightarrow{\partial_{p-1}^r} \mathbb{K}^{\lambda_{p,n-r}(R/L)} \longrightarrow \mathbb{K}^{\lambda_{p,n-r}(R/J(G))} \longrightarrow \mathbb{K}^{\lambda_{p,n-(r+1)}(R/L+K)} \xrightarrow{\partial_p^r} \cdots$$

Now we are ready to consider all the cases:

- i) If $\Lambda(R/L)$ and $\Lambda(R/L + K)$ are trivial then the Lyubeznik table of $R/J(G)$ is trivial as well. Notice that for $r = 2$ we have

$$\cdots \longrightarrow \mathbb{K}^{\lambda_{n-3,n-2}(R/J(G))} \longrightarrow \mathbb{K} \xrightarrow{\partial_{n-3}^2} \mathbb{K} \oplus \mathbb{K} \longrightarrow \mathbb{K} \longrightarrow 0$$

and thus $\lambda_{n-3,n-2}(R/J(G)) = 0$ and the vanishing of the rest of Lyubeznik numbers follow immediately.

- ii) If $\Lambda(R/L + K)$ is trivial then we have

$$0 \longrightarrow \mathbb{K}^{\lambda_{n-3,n-2}(R/L)} \longrightarrow \mathbb{K}^{\lambda_{n-3,n-2}(R/J(G))} \longrightarrow \mathbb{K} \xrightarrow{\partial_{n-3}^2} \mathbb{K} \oplus \mathbb{K} \longrightarrow \mathbb{K} \longrightarrow 0$$

and thus $\lambda_{n-3,n-2}(R/L) = \lambda_{n-3,n-2}(R/J(G))$. The rest of Lyubeznik numbers also coincide so we get $\Lambda(R/J(G)) = \Lambda(R/L)$.

- iii) If $\Lambda(R/L)$ is trivial then we have

$$0 \longrightarrow \mathbb{K}^{\lambda_{n-3,n-2}(R/J(G))} \longrightarrow \mathbb{K}^{\lambda_{n-3,n-3}(R/L+K)} \xrightarrow{\partial_{n-3}^2} \mathbb{K} \oplus \mathbb{K} \longrightarrow \mathbb{K} \longrightarrow 0$$

and thus $\lambda_{n-3,n-2}(R/J(G)) = \mathbb{K}^{\lambda_{n-3,n-3}(R/L+K)} - 1$. The rest of Lyubeznik numbers satisfy $\lambda_{p,n-r}(R/J(G)) = \lambda_{p,n-(r+1)}(R/L + K)$ and the result follows.

□

3.1. Splitting vertices. Let $J(G) \subseteq R$ be the cover ideal of a simple connected graph G . The easiest way to provide a MV-splitting $J(G) = L \cap K$ satisfying that the Lyubeznik table of R/K is trivial is by means of a **splitting vertex**. Namely, we fix a vertex, say x_n , and we decompose the ideal $J(G)$ depending on the edges that contain this vertex.

$$J(G) = \underbrace{\left(\bigcap_{x_i, x_j \notin N_G(x_n)} (x_i, x_j) \right)}_L \cap \underbrace{\left(\bigcap_{x_k \in N_G(x_n)} (x_k, x_n) \right)}_K$$

Observe that $J(G) = K$ when G is a star graph with x_n being the internal vertex so in this case we will pick up a different vertex. In general we have:

- $L = J(G \setminus \{x_n\})$ is the cover ideal of the subgraph obtained removing the vertex x_n .
- $K = J(K_{1,g})$ is the cover ideal of a star graph with $g = \deg(x_n)$.
- $L + K$ is a height 3 monomial ideal which admits a (non-necessarily minimal) primary decomposition of the form:

$$L + K = \bigcap_{x_k \in N_G(x_n)} \left[\left(\bigcap_{x_i, x_j \notin N_G(x_n), N_G(x_k)} (x_i, x_j, x_k, x_n) \right) \cap \left(\bigcap_{x_l \in N_G(x_k)} (x_l, x_k, x_n) \right) \right].$$

Of course we can make it minimal removing conveniently the extra components. Notice that $\Lambda(R/K)$ is trivial. In order to check that this decomposition indeed provides a MV-splitting we only need to invoke [19, Theorem 4.2] where it is proved that every vertex is a splitting vertex except for some limit cases where the vertex is isolated or its complement consists of isolated vertices.

Remark 3.4. The operations of sum and intersection distribute over each other in the case of squarefree monomial ideals. This gives us a very simple method to find (non-necessarily minimal) primary decompositions of sums $L + K$ of squarefree monomial ideals in terms of the primary decompositions of L and K respectively. Namely,

$$\underbrace{(I_1 \cap \dots \cap I_c)}_L + \underbrace{(J_1 \cap \dots \cap J_d)}_K = (I_1 + J_1) \cap \dots \cap (I_c + J_d).$$

In order to make it minimal we just remove the extra components. Throughout the rest of this work we will use this fact without further reference.

3.1.1. Splitting vertices of degree one. Let x_n be a splitting vertex of a graph G . Assume that its degree is one and, for simplicity, we will take x_{n-1} as the unique vertex in its neighbourhood. We can rephrase it by saying that we are adding a **whisker** to the vertex x_{n-1} of the graph $G \setminus \{x_n\}$.

Proposition 3.5. *Let $J(G) \subseteq R$ be the cover ideal of a simple connected graph G . Let $x_n \in V_G$ be a vertex of degree one. Then $\Lambda(R/J(G)) = \Lambda(R/J(G \setminus \{x_n\}))$.*

Proof. Let x_{n-1} be the unique vertex in the neighbourhood of x_n . Then, the MV-splitting $J_G = L \cap K$ given by x_n has $K = (x_{n-1}, x_n)$ and

$$\begin{aligned} L + K &= \left(\bigcap_{x_i, x_j \notin N_G(x_{n-1})} (x_i, x_j, x_{n-1}, x_n) \right) \cap \left(\bigcap_{x_l \in N_G(x_{n-1})} (x_l, x_{n-1}, x_n) \right) \\ &= \underbrace{\left[\left(\bigcap_{x_i, x_j \notin N_G(x_{n-1})} (x_i, x_j) \right) \cap \left(\bigcap_{x_l \in N_G(x_{n-1})} (x_l) \right) \right]}_M + (x_{n-1}, x_n) \end{aligned}$$

The ideal M is a height one ideal in two sets of disjoint variables. Therefore, its Lyubeznik table is trivial because of Proposition 2.5. Given the isomorphism

$$\mathbb{K}[x_1, \dots, x_n]/L + K \cong \mathbb{K}[x_1, \dots, x_{n-2}]/M$$

we get that $\Lambda(R/L + K)$ is trivial as well. Then the result follows using Theorem 3.3 and the fact that $L = J(G \setminus \{x_n\})$. \square

3.1.2. Splitting vertices of degree two. Let x_n be a splitting vertex of a graph G . Assume that its degree is two and the vertices in its neighbourhood are x_{n-1} and x_{n-2} . In this case we also have the invariance of the Lyubeznik table after removing the splitting vertex under certain extra conditions.

Proposition 3.6. *Let $J(G) \subseteq R$ be the cover ideal of a simple connected graph G . Let $x_n \in V_G$ be a vertex of degree 2 with $N_G(x_n) = \{x_{n-2}, x_{n-1}\}$. If any of the following conditions hold:*

- i) $\{x_{n-1}, x_{n-2}\} \in E_G$,*
- ii) there exists $x_c \in N_G(x_{n-1}) \cap N_G(x_{n-2})$,*

then $\Lambda(R/J(G)) = \Lambda(R/J(G \setminus \{x_n\}))$.

Proof. We have a MV-splitting $J(G) = L \cap K$ where $K = (x_{n-1}, x_n) \cap (x_{n-2}, x_n)$ and

$$\begin{aligned} L &= \left(\bigcap_{x_i, x_j \notin N_G(x_n)} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_a \in N_G(x_{n-1}) \\ x_a \notin N_G(x_{n-2})}} (x_a, x_{n-1}) \right) \cap \left(\bigcap_{\substack{x_b \in N_G(x_{n-2}) \\ x_b \notin N_G(x_{n-1})}} (x_b, x_{n-2}) \right) \\ &\quad \cap \left(\bigcap_{\substack{x_c \in N_G(x_{n-1}) \\ x_c \in N_G(x_{n-2})}} (x_c, x_{n-2}) \cap (x_c, x_{n-1}) \right) \cap (x_{n-2}, x_{n-1}) \end{aligned}$$

Under the assumptions we are considering, at least one of the last components in this decomposition must appear. Therefore

$$\begin{aligned}
 L + K &= \underbrace{\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j, x_{n-1}, x_n) \right) \cap \left(\bigcap_{\substack{x_a \in N_G(x_{n-1}) \\ x_a \notin N_G(x_{n-2})}} (x_a, x_{n-1}, x_n) \right)}_M \\
 &\quad \cap \underbrace{\left(\bigcap_{\substack{x_c \in N_G(x_{n-1}) \\ x_c \in N_G(x_{n-2})}} (x_c, x_{n-1}, x_n) \right) \cap (x_{n-2}, x_{n-1})}_M \\
 &\quad \cap \underbrace{\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j, x_{n-2}, x_n) \right) \cap \left(\bigcap_{\substack{x_b \in N_G(x_{n-2}) \\ x_b \notin N_G(x_{n-1})}} (x_b, x_{n-2}, x_n) \right)}_N
 \end{aligned}$$

Notice that we can rephrase the ideals M and N as

$$\begin{aligned}
 M &= \left[\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_a \in N_G(x_{n-1}) \\ x_a \notin N_G(x_{n-2})}} (x_a) \right) \cap \left(\bigcap_{\substack{x_c \in N_G(x_{n-1}) \\ x_c \in N_G(x_{n-2})}} (x_c) \right) \cap (x_{n-2}) \right] + (x_{n-1}, x_n) \\
 N &= \left[\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-2})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_b \in N_G(x_{n-2}) \\ x_b \notin N_G(x_{n-1})}} (x_b) \right) \right] + (x_{n-2}, x_n)
 \end{aligned}$$

so both $\Lambda(R/M)$ and $\Lambda(R/N)$ are trivial using Proposition 2.5. Moreover, in the case that condition $i)$ is satisfied, we have

$$M + N = \left[\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_b \in N_G(x_{n-2}) \\ x_b \notin N_G(x_{n-1})}} (x_b) \right) \cap \left(\bigcap_{\substack{x_c \in N_G(x_{n-1}) \\ x_c \in N_G(x_{n-2})}} (x_c) \right) \right] + (x_{n-2}, x_{n-1}, x_n)$$

Under condition $ii)$ we have

$$M + N = \left[\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_a \in N_G(x_{n-1}) \\ x_a \notin N_G(x_{n-2}) \\ x_b \in N_G(x_{n-2}) \\ x_b \notin N_G(x_{n-1})}} (x_a, x_b) \right) \cap \left(\bigcap_{\substack{x_c \in N_G(x_{n-1}) \\ x_c \in N_G(x_{n-2})}} (x_c) \right) \right] + (x_{n-2}, x_{n-1}, x_n)$$

In any case the Lyubeznik table of $R/M + N$ is trivial since we are dealing with a height one ideal in a disjoint set of variables so we can apply Proposition 2.5 once again. As a consequence of Theorem 3.3, the Lyubeznik table $\Lambda(R/L + K)$ is trivial and the result follows applying Theorem 3.3 once more. To finish the proof we need to check that the decomposition $L + K = M \cap N$ is, indeed, a MV-splitting.

If $\{x_{n-2}, x_{n-1}\}$ is not an edge of G we have that the variable x_{n-1} appears in all the components of the primary decomposition of M but not in N . We also have that x_{n-2} appears in all the components of N but not in M and both x_{n-2}, x_{n-1} appear in $M + N$. In particular the posets associated to M , N and $M + N$ do not have common ideals. Then the result follows from Corollary 2.11.

When $\{x_{n-2}, x_{n-1}\}$ is an edge of G we have to be more careful. The variable x_{n-1} appears in all the components of $M + N$ but not in N and thus its corresponding posets do not have common ideals so we only have to compare $M + N$ with M . Indeed, since the variables x_a 's and x_b 's do not belong to both ideals and x_{n-1}, x_n do so we may just assume that the ideals are

$$M = \underbrace{\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_c \in N_G(x_{n-1}) \\ x_c \in N_G(x_{n-2})}} (x_c) \right)}_Q \cap (x_{n-2})$$

$$M + N = \left[\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_c \in N_G(x_{n-1}) \\ x_c \in N_G(x_{n-2})}} (x_c) \right) \right] + (x_{n-2})$$

and thus we have a MV-splitting

$$0 \longrightarrow H_Q^r(R) \oplus H_{(x_{n-2})}^r(R) \longrightarrow H_M^r(R) \longrightarrow H_{M+N}^{r+1}(R) \longrightarrow 0$$

so the graded pieces of $H_{M+N}^{r+1}(R)$ are related to the graded pieces of $H_M^r(R)$ instead of those of $H_M^{r+1}(R)$. \square

We can still say something about the Lyubeznik numbers of $J(G)$ in the event that the hypothesis of the previous proposition do not hold. In this case, the Lyubeznik table of $R/L + K$ is not trivial so we need to control the connecting morphisms

$$\mathbb{K}^{\lambda_{p-1, n-(r+1)}(R/L+K)} \xrightarrow{\partial_{p-1}^r} \mathbb{K}^{\lambda_{p, n-r}(R/L)}$$

considered in Discussion 2.12. If the Lyubeznik table of $L = J(G \setminus \{x_n\})$ is trivial or, more generally, the connecting morphisms are zero, we can give a formula for the Lyubeznik numbers

of $J(G)$ in terms of those of $J(G \setminus \{x_n\})$ and the Lyubeznik numbers associated to the graph

$$H := (G \setminus \{x_n\}) \cup \bigcup_{\substack{x_a \in N_G(x_{n-1}) \\ x_b \in N_G(x_{n-2})}} \{x_a, x_b\}$$

whose cover ideal is

$$J(H) = \left(\bigcap_{\substack{x_i, x_j \notin N_G(x_{n-2}) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_a \in N_G(x_{n-2}) \\ x_b \in N_G(x_{n-1})}} (x_a, x_b) \right)$$

In other words, the graph H is obtained by adjoining to $G \setminus \{x_n\}$ a complete bipartite graph in the set of vertices $N_G(x_{n-2})$ and $N_G(x_{n-1})$.

Proposition 3.7. *Let $J(G) \subseteq R$ be the cover ideal of a simple connected graph G . Let $x_n \in V_G$ be a vertex of degree 2 with $N_G(x_n) = \{x_{n-1}, x_{n-2}\}$. Assume that conditions i) and ii) of Proposition 3.6 no not hold and that the connecting morphisms ∂_p^r are zero for all $r > 2$ and for all p . Then we have*

$$\lambda_{d,d}(R/J(G)) = 1, \quad \lambda_{d-1,d}(R/J(G)) = \lambda_{d-1,d}(R/J(G \setminus \{x_n\})) + 1,$$

$$\lambda_{p,r}(R/J(G)) = \lambda_{p,r}(R/J(G \setminus \{x_n\})) + \lambda_{p,r-2}(R/J(H))$$

for $r = 2, \dots, d-1$ and $p = 0, \dots, r-2$ and the rest of Lyubeznik numbers satisfy

$$\lambda_{p,r}(R/J(G)) = \lambda_{p,r}(R/J(G \setminus \{x_n\})) .$$

That is,

$$\Lambda(R/J(G)) = \begin{pmatrix} \lambda_{0,0} & \lambda_{0,1} & \lambda_{0,2} + \lambda'_{0,0} & \lambda_{0,3} + \lambda'_{0,1} & \cdots & \lambda_{0,d-1} + \lambda'_{0,d-3} & \lambda_{0,d} \\ & \lambda_{1,1} & \lambda_{1,2} & \lambda_{1,3} + \lambda'_{1,1} & \cdots & \lambda_{1,d-1} + \lambda'_{1,d-3} & \lambda_{1,d} \\ & & \lambda_{2,2} & \lambda_{2,3} & \cdots & \lambda_{2,d-1} + \lambda'_{2,d-3} & \lambda_{2,d} \\ & & & & \ddots & \vdots & \vdots \\ & & & & & \lambda_{d-3,d-1} + \lambda'_{d-3,d-3} & \lambda_{d-3,d} \\ & & & & & \lambda_{d-2,d-1} & \lambda_{d-2,d} \\ & & & & & \lambda_{d-1,d-1} & \lambda_{d-1,d} + 1 \\ & & & & & & 1 \end{pmatrix}$$

where

$$\Lambda(R/J(G \setminus \{x_n\})) = \begin{pmatrix} \lambda_{0,0} & \cdots & \lambda_{0,d} \\ & \ddots & \vdots \\ & & \lambda_{d,d} \end{pmatrix}, \quad \Lambda(\mathbb{K}[x_1, \dots, x_{n-3}]/J(H)) = \begin{pmatrix} \lambda'_{0,0} & \cdots & \lambda'_{0,d-3} \\ & \ddots & \vdots \\ & & \lambda'_{d-3,d-3} \end{pmatrix}$$

Proof. Following the same approach as in the proof of Proposition 3.6 we have a MV-splitting $J(G) = L \cap K$ with $K = (x_{n-1}, x_n) \cap (x_{n-2}, x_n)$ but in this case we also have

$$L = \left(\bigcap_{x_i, x_j \notin N_G(x_n)} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_a \in N_G(x_{n-1}) \\ x_a \notin N_G(x_{n-2})}} (x_a, x_{n-1}) \right) \cap \left(\bigcap_{\substack{x_b \in N_G(x_{n-2}) \\ x_b \notin N_G(x_{n-1})}} (x_b, x_{n-2}) \right)$$

$$L + K = \underbrace{\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j, x_{n-1}, x_n) \right) \cap \left(\bigcap_{\substack{x_a \in N_G(x_{n-1}) \\ x_a \notin N_G(x_{n-2})}} (x_a, x_{n-1}, x_n) \right)}_M$$

$$\cap \underbrace{\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j, x_{n-2}, x_n) \right) \cap \left(\bigcap_{\substack{x_b \in N_G(x_{n-2}) \\ x_b \notin N_G(x_{n-1})}} (x_b, x_{n-2}, x_n) \right)}_N$$

Once again we rewrite the ideals M and N as

$$M = \left[\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_a \in N_G(x_{n-1}) \\ x_a \notin N_G(x_{n-2})}} (x_a) \right) \right] + (x_{n-1}, x_n)$$

$$N = \left[\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_n) \\ x_i, x_j \notin N_G(x_{n-2})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_b \in N_G(x_{n-2}) \\ x_b \notin N_G(x_{n-1})}} (x_b) \right) \right] + (x_{n-2}, x_n)$$

so both $\Lambda(R/M)$ and $\Lambda(R/N)$ are trivial. Moreover

$$M + N = \underbrace{\left[\left(\bigcap_{\substack{x_i, x_j \notin N_G(x_{n-2}) \\ x_i, x_j \notin N_G(x_{n-1})}} (x_i, x_j) \right) \cap \left(\bigcap_{\substack{x_a \in N_G(x_{n-1}) \\ x_b \in N_G(x_{n-2})}} (x_a, x_b) \right) \right]}_{J(H)} + (x_{n-2}, x_{n-1}, x_n).$$

We have that the variable x_{n-1} appears in all the components of the primary decomposition of M but not in N . We also have that x_{n-2} appears in all the components of N but not in M and both x_{n-2}, x_{n-1} appear in $M + N$. In particular the posets associated to M , N and $M + N$ do not have common ideals. Therefore $L + K = M \cap N$ is a MV-splitting by using Corollary 2.11.

Applying the long exact sequence of local cohomology modules to the short exact sequence

$$0 \longrightarrow H_M^r(R) \oplus H_N^r(R) \longrightarrow H_{L+K}^r(R) \longrightarrow H_{M+N}^{r+1}(R) \longrightarrow 0$$

we get, for $r = 3$,

$$\dots \longrightarrow H_m^{n-4}(H_{M+N}^4(R)) \xrightarrow{\partial_{n-4}^3} H_m^{n-3}(H_M^3(R)) \oplus H_m^{n-3}(H_N^3(R)) \longrightarrow H_m^{n-3}(H_{L+K}^3(R)) \longrightarrow 0$$

Since $\Lambda(R/M)$ and $\Lambda(R/N)$ are trivial and taking into account that $\text{ht } M = \text{ht } N = 3$ and $\text{ht } M + N = 5$ we get

$$\begin{aligned} [H_m^{n-3}(H_{L+K}^3(R))]_{-1} &\cong \mathbb{K}^2 \\ [H_m^p(H_{L+K}^3(R))]_{-1} &\cong 0 \quad \forall p < n-3 \\ [H_m^p(H_{L+K}^r(R))]_{-1} &\cong [H_m^p(H_{M+N}^{r+1}(R))]_{-1} \quad \forall p, \forall r > 3. \end{aligned}$$

Now, if we go back to the long exact sequence 2.4 considered in Discussion 2.12 to compute the Lyubeznik table of $R/J(G)$ we get

$$0 \longrightarrow \mathbb{K}^{\lambda_{n-3,n-2}(R/L)} \longrightarrow \mathbb{K}^{\lambda_{n-3,n-2}(R/J(G))} \longrightarrow \mathbb{K}^{\lambda_{n-3,n-3}(R/L+K)} \xrightarrow{\partial_{n-3}^3} \mathbb{K} \oplus \mathbb{K} \longrightarrow \mathbb{K} \longrightarrow 0$$

and $\lambda_{p,n-2}(R/L) = \lambda_{p,n-2}(R/J(G))$ for all $p < n-3$. Moreover, for $r > 2$ and any p , the long exact sequence becomes

$$\dots \longrightarrow \mathbb{K}^{\lambda_{p-1,n-(r+1)}(R/L+K)} \xrightarrow{\partial_{p-1}^r} \mathbb{K}^{\lambda_{p,n-r}^r(R/L)} \longrightarrow \mathbb{K}^{\lambda_{p,n-r}(R/J(G))} \longrightarrow \mathbb{K}^{\lambda_{p,n-(r+1)}(R/L+K)} \xrightarrow{\partial_p^r} \dots$$

Therefore we have:

$$\begin{aligned} \lambda_{d,d}(R/J(G)) &= 1, \\ \lambda_{d-1,d}(R/J(G)) &= \lambda_{d-1,d}(R/L) + 1, \\ \lambda_{p,d}(R/J(G)) &= \lambda_{p,d}(R/L) \quad \forall p < d-1 \end{aligned}$$

and the rest of Lyubeznik numbers depend on the connecting morphisms

$$\mathbb{K}^{\lambda_{p-1,n-(r+2)}(R/M+N)} = \mathbb{K}^{\lambda_{p-1,n-(r+1)}(R/L+K)} \xrightarrow{\partial_{p-1}^r} \mathbb{K}^{\lambda_{p,n-r}(R/L)}.$$

Assuming that the connecting morphisms are zero we get

$$\lambda_{p,n-r}(R/J(G)) = \lambda_{p,n-r}(R/L) + \lambda_{p,n-(r+1)}(R/L+K)$$

and the result follows since we have $L = J(G \setminus \{x_n\})$ and an isomorphism

$$\mathbb{K}[x_1, \dots, x_n]/M + N \cong \mathbb{K}[x_1, \dots, x_{n-3}]/J(H)$$

and thus $\Lambda(R/M + N) = \Lambda(\mathbb{K}[x_1, \dots, x_{n-3}]/J(H))$. □

We highlight the following particular case.

Corollary 3.8. *Under the assumptions of Proposition 3.7, if $\Lambda(R/J(G \setminus \{x_n\}))$ is trivial then the Lyubeznik table of $R/J(G)$ is*

$$\Lambda(R/J(G)) = \begin{pmatrix} 0 & 0 & \lambda'_{0,0} & \lambda'_{0,1} & \cdots & \lambda'_{0,d-3} & 0 \\ & 0 & 0 & \lambda'_{1,1} & \cdots & \lambda'_{1,d-3} & 0 \\ & & 0 & 0 & \cdots & \lambda'_{2,d-3} & 0 \\ & & & & \ddots & \vdots & \vdots \\ & & & & & \lambda'_{d-3,d-3} & 0 \\ & & & & & 0 & 0 \\ & & & & & 0 & 1 \\ & & & & & & 1 \end{pmatrix}$$

where

$$\Lambda(\mathbb{K}[x_1, \dots, x_{n-3}]/J(H)) = \begin{pmatrix} \lambda'_{0,0} & \cdots & \lambda'_{0,d-3} \\ & \ddots & \vdots \\ & & \lambda'_{d-3,d-3} \end{pmatrix}$$

3.1.3. *Splitting vertices of maximal degree.* We turn our attention to the case of a splitting vertex x_n of degree $n - 1$. That is, $\{x_i, x_n\}$ is an edge of G for $i = 1, \dots, n - 1$.

Proposition 3.9. *Let $J(G) \subseteq R$ be the cover ideal of a simple connected graph G . Let $x_n \in V_G$ be a vertex of degree $n - 1$. Then, the Lyubeznik table of $R/J(G)$ is trivial.*

Proof. We have a MV-splitting $J_G = L \cap K$ where $K = (x_1, x_n) \cap \cdots \cap (x_{n-2}, x_n)$ and, given the fact that all the vertices are in the neighbourhood of x_n , we have

$$L + K = \bigcap_{\substack{i,j \neq n \\ \{x_i, x_j\} \in E_G}} (x_i, x_j, x_n) = L + (x_n)$$

Recall that $\Lambda(R/K)$ is trivial so the long exact sequence associated to the MV-splitting reduces to

$$\cdots \longrightarrow H_m^p(H_{J(G)}^r(R)) \longrightarrow H_m^p(H_{L+(x_n)}^{r+1}(R)) \xrightarrow{\partial_p^r} H_m^{p+1}(H_L^r(R)) \longrightarrow \cdots$$

Using the interpretation of the connecting morphisms ∂_p^r 's in terms of the corresponding linear strands given at the end of Section 2 we observe that we are comparing the linear strands $\mathbb{F}_{\bullet}^{\langle r+1 \rangle}((L + (x_n))^\vee)^*$ and $\mathbb{F}_{\bullet}^{\langle r \rangle}(L^\vee)^*$ which are essentially the same (modulo a shifting), so the induced morphisms in homology are isomorphisms and the result follows. \square

3.2. **Examples.** The MV-splitting techniques developed in the previous section allow us to compute the Lyubeznik table of many families of graphs directly from the combinatorics of the graph without an explicit computation of the corresponding local cohomology modules. The idea is to choose a convenient splitting vertex and reduce the computation to the case of a graph in a smaller number of vertices.

An interpretation of Proposition 3.5 is that the Lyubeznik table remains invariant under the operation of removing whiskers. In this way we may simplify our original graph and, in the case of acyclic graphs, we can deduce the triviality of the Lyubeznik table by reducing the computation to the case of a single edge. In particular we get:

Corollary 3.10. *The Lyubeznik table of the cover ideal of a **path** is trivial.*

Corollary 3.11. *The Lyubeznik table of the cover ideal of a **tree** is trivial.*

Using Corollary 2.6 we can also consider the case of forests.

Corollary 3.12. *The Lyubeznik numbers of the cover ideal of a **forest** with c connected components are*


$$\lambda_{d-2k, d-k}(R/J(G)) = \binom{c}{k+1} \quad \text{for } k = 0, \dots, c-1$$

and the rest of Lyubeznik numbers are zero..


Another source of examples of trivial Lyubeznik tables is using Proposition 3.9. It says that the **cone** of any graph G , has a trivial Lyubeznik table. In particular:

Corollary 3.13. *The Lyubeznik table of the cover ideal of a **wheel** is trivial.*


Another way of simplifying our original graph is by means of Proposition 3.6, which says that we can remove what we call **handles** (or equivalently 3 and 4-cycles) having the following shape:



This gives as a very visual method to reduce the computation of Lyubeznik tables of graphs. For example, removing the yellow vertices indicated below do not modify the Lyubeznik table and thus, in the end, we see that the Lyubeznik table of the following graph is trivial.



Now we turn our attention to the case of cycles. Applying iteratively Corollary 3.8 we will obtain a closed formula for the Lyubeznik numbers. To illustrate our methods we present the case of a 6-cycle. The corresponding graphs L and H are represented as follows:



Notice that L is a path so it has trivial Lyubeznik table and thus we can use Corollary 3.8. Moreover we have that H is a 3-cycle.

Proposition 3.14. *The Lyubeznik numbers of the cover ideal of a n -cycle with $n = 3k + \ell$, $\ell \in \{-1, 0, 1\}$, are*

$$\lambda_{d-3i, d-i}(R/J(G)) = 1, \quad \lambda_{d-3i-1, d-i}(R/J(G)) = 1 \quad \text{for } i = 0, \dots, k-2.$$

$$\lambda_{d-3i, d-i}(R/J(G)) = 1 \quad \text{for } i = k-1.$$

and the rest of Lyubeznik numbers are zero.

Proof. Notice that, if we denote G the n -cycle, then $G \setminus \{x_n\}$ is a path so its Lyubeznik table is trivial. On the other hand, the graph H obtained from G is a $(n-3)$ -cycle so we can use induction on k and Corollary 3.8 to produce the formula for the Lyubeznik numbers. \square


Example 3.15. The Lyubeznik table of a cycle C_n in n vertices for $n = 5, \dots, 11$ are:

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ & 0 & 1 & 0 \\ & & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ & 0 & 0 & 0 & 0 \\ & & 0 & 1 & 0 \\ & & & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ & 0 & 0 & 1 & 0 & 0 \\ & & 0 & 0 & 0 & 0 \\ & & & 0 & 1 & 0 \\ & & & & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ & 0 & 0 & 0 & 0 & 1 & 0 \\ & & 0 & 0 & 0 & 0 & 0 \\ & & & 0 & 1 & 0 & 0 \\ & & & & 0 & 1 & 0 \\ & & & & & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ & & 0 & 0 & 0 & 1 & 0 & 0 \\ & & & 0 & 0 & 0 & 1 & 0 \\ & & & & 0 & 1 & 0 & 0 \\ & & & & & 0 & 1 & 0 \\ & & & & & & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ & & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ & & & 0 & 0 & 0 & 0 & 1 & 0 \\ & & & & 0 & 1 & 0 & 0 & 0 \\ & & & & & 0 & 1 & 0 & 0 \\ & & & & & & 0 & 1 & 0 \\ & & & & & & & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ & & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ & & & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ & & & & 0 & 1 & 0 & 0 & 0 & 0 \\ & & & & & 0 & 1 & 0 & 0 & 0 \\ & & & & & & 0 & 1 & 0 & 0 \\ & & & & & & & 0 & 1 & 0 \\ & & & & & & & & 0 & 1 \end{pmatrix}$$

Certainly a new source for finding non trivial Lyubeznik tables is to consider graphs obtained by joining cycles. After removing whiskers and handles we reduce to the case of cycles

joined by paths or sharing edges in such a way that we can still find degree two vertices that we can remove in order to simplify the graph. For example, the complement of a 6-cycle



can be interpreted as joining two 3-cycles and two 4-cycles. However all the vertices have degree three so we cannot apply our methods.

To start our study we will consider the following joining operation.

Definition 3.16. Let C_m and C_n be two cycles in m and n vertices respectively. We say that they are L -joined and we will denote by $C_m * C_n$ the corresponding graph if they share at most one edge or if they are joined by a path.

When we remove a vertex from $C_m * C_n$ we do not obtain a subgraph L with trivial Lyubeznik table. However, we still can use Proposition 3.7 to obtain the following formula for the Lyubeznik numbers.


Proposition 3.17. Let C_m and C_n be two cycles with $m \leq n$ and $m = 3k_1 + \ell_1$, $n = 3k_2 + \ell_2$ and $\ell_1, \ell_2 \in \{-1, 0, 1\}$. Then the Lyubeznik numbers of the cover ideal of $G = C_m * C_n$ are

$$\begin{aligned} \lambda_{d-3i, d-i}(R/J(G)) &= i + 1, & \lambda_{d-3i-1, d-i}(R/J(G)) &= i + 2 & \text{for } i &= 0, \dots, k_1 - 2, \\ \lambda_{d-3i, d-i}(R/J(G)) &= k_1, & \lambda_{d-3i-1, d-i}(R/J(G)) &= k_1 & \text{for } i &= k_1 - 1, \dots, k_2 - 2, \\ \lambda_{d-3i, d-i}(R/J(G)) &= k_1 + k_2 - i - 1, & \lambda_{d-3i-1, d-i}(R/J(G)) &= k_1 + k_2 - i - 2 \end{aligned}$$

for $i = k_2 - 1, \dots, k_1 + k_2 - 2$ and the rest of Lyubeznik numbers are zero. Here we follow the convention that in the case where $k_1 = k_2$ the set of indices $i = k_1 - 1, \dots, k_2 - 2$ is empty.

Proof. We are going to use Proposition 3.7 so we will follow the same terminology used in its proof. Pick a splitting vertex x_m of degree two in C_m . Notice that $G \setminus \{x_m\}$ has the same Lyubeznik table as C_n and the graph H associated to G is $C_{m-3} * C_n$ so we can proceed by induction on k_1 . The fact that C_m and C_n share at most one edge implies that the induction step will end up in a graph H having the Lyubeznik table of C_n . \square

We illustrate the case $G = C_6 * C_6$ as follows:



More generally we can consider the family of graphs obtained using the $*$ operation, which is a family that includes **cycles with chords** or **cactus graphs**. We can iterate the methods used in Proposition 3.17 in order to compute the Lyubeznik table of any L -joined graph of the form $C_{n_1} * \dots * C_{n_r}$. We are not going to give a closed formula for its Lyubeznik table since it depends on the indices $n_i = 3k_i + \ell_i$, with $\ell_i \in \{-1, 0, 1\}$, for $i = 1, \dots, r$ and one should distinguish too many cases which makes it very tedious and not very illustrative.

We will just point out that the non-vanishing Lyubeznik numbers are

$$\lambda_{d-3i, d-i}(R/J(G)), \quad \lambda_{d-3i-1, d-i}(R/J(G)) \quad \text{for } i = 0, \dots, k_1 + \dots + k_r - r$$

Moreover we have $\lambda_{d, d} = 1$, $\lambda_{d-1, d} = r$ and $\lambda_{d-3(k_1 + \dots + k_r - r), d - (k_1 + \dots + k_r - r)} = 1$.

In the case that we have two cycles C_m and C_n sharing more than one edge we can still use iteratively Proposition 3.7 to compute its Lyubeznik table. However, in this case we will not end up with a graph H having the Lyubeznik table of C_n as the following example shows.

Example 3.18. Let G be two 8-cycles sharing four consecutive vertices. The Lyubeznik table of G can be obtained, using Proposition 3.7, from the Lyubeznik tables of two graphs L_1 and H_1 . Notice that L_1 has the same Lyubeznik type as an 8-cycle C_8 , but in order to get the Lyubeznik table of H_1 we have to apply Proposition 3.7 once again. We illustrate the procedure as follows:

Remark 3.19. The number of connected graphs in a given number of vertices is very large so it seems infeasible to find all the possible configurations of Lyubeznik tables. If the number of vertices is small, we can use our methods to reduce the number of examples we have to consider and compute the Lyubeznik numbers of the remaining graphs using the algorithm presented in [6].

We made the computations in the case we have either 4, 5 or 6 vertices and the possible Lyubeznik types are reflected in the following table. Not surprisingly, having trivial Lyubeznik table is the most common situation.

n	trivial	cycle	complement cycle	total
4	6	–	–	6
5	20	1	–	21
6	106	5	1	112

TABLE 1. Number of graphs in n vertices with given Lyubeznik type.

In order to expand this list it would be desirable to develop techniques to deal with splitting vertices of degree bigger than two and identify graphs with non trivial Lyubeznik tables other than cycles or complement of cycles.

4. BASS NUMBERS OF LOCAL COHOMOLOGY MODULES OF COVER IDEALS OF GRAPHS

Let G be a simple graph and, given $\alpha \in \{0, 1\}^n$, we denote by G_α the subgraph of G obtained by removing the vertices x_i such that $\alpha_i = 0$. Indeed we will only consider those α 's for which G_α is not a set of isolated vertices, which means that $E_{G_\alpha} \neq \emptyset$. In the case that G_α has an isolated vertex, we can assume that we are working in a smaller set of vertices by removing the isolated one. Notice that the cover ideal of G_α is an ideal in the polynomial ring $R_{\mathfrak{p}_\alpha} = \mathbb{K}[x_i \mid \alpha_i = 1]$. As we mentioned in Remark 2.3, Bass numbers behave well with respect to restriction and thus, the Bass numbers of the local cohomology module $H_{J(G)}^{n-i}(R)$ with respect to the face ideal \mathfrak{p}_α are nothing but the Lyubeznik numbers corresponding to the subgraph G_α . More precisely,

$$\mu_p(\mathfrak{p}_\alpha, H_{J(G)}^{n-i}(R)) = \lambda_{p,i}(R_{\mathfrak{p}_\alpha}/J(G_\alpha))$$

Certainly G_α are not necessarily connected graphs and thus one needs to use Proposition 2.5 in order to compute these Lyubeznik numbers. In what follows we will denote c_α as the number of connected components of G_α and $c_{max} = \max\{c_\alpha \mid \alpha \in \{0, 1\}^n\}$.

Example 4.1. The maximal number of connected components of a path or a cycle is achieved when we remove every third vertex from the graph. For an n -path we have $c_{max} = \lceil \frac{n+1}{3} \rceil$. However for n -cycle we have to be a little more careful and we have $c_{max} = \lfloor \frac{n}{3} \rfloor$. In particular, for $n = 3k - 1$ we have that the n -path has $c_{max} = k$ and the n -cycle has $c_{max} = k - 1$. For the rest of cases they coincide.

4.1. Linear injective resolutions. Let G be a graph that $R/J(G)$ is Cohen-Macaulay. The Bass numbers of this class of graphs are completely determined as it has been shown in [8] using a simple spectral sequence argument. For completeness we include the result here but giving an equivalent description in terms of the connectivity of the subgraphs.

Proposition 4.2. *Let $J(G) \subseteq R$ be the cover ideal of a simple connected graph G . Then the following are equivalent*

- i) $R/J(G)$ is Cohen-Macaulay.
- ii) $\mu_p(\mathfrak{p}_\alpha, H_{J(G)}^2(R)) = \delta_{p,|\alpha|-2}$
- iii) G_α is connected and $\Lambda(R_{\mathfrak{p}_\alpha}/J(G_\alpha))$ is trivial for all $\alpha \in \{0, 1\}^n$.

In this case, the \mathbb{Z}^n -graded injective resolution $\mathbb{I}_\bullet(H_{J(G)}^2(R))$ has a very rigid structure which resembles the injective resolution of Gorenstein rings. Namely we have:

$$0 \longrightarrow H_{J(G)}^2(R) \longrightarrow \bigoplus_{|\alpha|=2} E_\alpha \longrightarrow \bigoplus_{|\alpha|=3} E_\alpha \longrightarrow \dots \longrightarrow \bigoplus_{|\alpha|=n-1} E_\alpha \longrightarrow E_1 \longrightarrow 0,$$

where, at each component of the resolution we are only considering those α 's such that $\mathfrak{p}_\alpha \in \text{Supp}R/J(G)$. Notice that this resolution is linear, it only has one linear strand.

We will show next that we may find non Cohen-Macaulay graphs still having a rigid injective resolution. To such purpose let's consider the following family.

Assumptions 4.3. Let $J(G) \subseteq R$ be the cover ideal of a simple (not necessarily connected) graph G such that, for any $\alpha \in \{0, 1\}^n$, all the connected components of G_α have trivial Lyubeznik table.

As we have seen in Section 3, the condition of having trivial Lyubeznik table is very common and is not difficult to find families of graphs satisfying Assumptions 4.3. An interesting example would be the case of **forests** but we may also include Cohen-Macaulay graphs. As a direct consequence of Corollary 2.6 we get:

Theorem 4.4. *Let $J(G) \subseteq R$ be an ideal satisfying Assumptions 4.3. Then, the Bass numbers of the corresponding local cohomology modules are*

$$\mu_p(\mathfrak{p}_\alpha, H_{J(G)}^{k+1}(R)) = \delta_{p,|\alpha|-2k} \cdot \binom{c_\alpha}{k} \quad \text{for } k = 1, \dots, c_\alpha,$$

where $\delta_{i,j}$ is Kronecker's delta function.

In this case we also have a rigid injective resolution in the sense that they only have one linear strand.

Example 4.5. Let G be a 5-path. We have that all the subgraphs G_α without isolated vertices are connected except for the case $\alpha = (1, 1, 0, 1, 1)$ in which G_α has two connected components. Applying the above result we get the following linear injective resolutions

$$\begin{aligned}
0 \longrightarrow H_{J(G)}^2(R) &\longrightarrow \bigoplus_{|\alpha|=2} E_\alpha \longrightarrow \bigoplus_{|\alpha|=3} E_\alpha \longrightarrow \left(\begin{array}{c} E_{(1,1,0,1,1)}^2 \\ \oplus \\ \bigoplus_{\substack{|\alpha|=4 \\ \alpha \neq (1,1,0,1,1)}} E_\alpha \end{array} \right) \longrightarrow E_1 \longrightarrow 0, \\
0 \longrightarrow H_{J(G)}^3(R) &\longrightarrow E_{(1,1,0,1,1)} \longrightarrow 0,
\end{aligned}$$

where we are only considering those α 's such that $\mathfrak{p}_\alpha \in \text{Supp}R/J(G)$. Notice that in this case we have two local cohomology modules different from zero.

Quite surprisingly we may provide a vanishing criterion for the local cohomology modules in terms of the connected components of the subgraphs. Recall that the **cohomological dimension** of an ideal J is the maximum r for which $H_J^r(R) \neq 0$.

Proposition 4.6. *Let $J(G) \subseteq R$ be an ideal satisfying Assumption 4.3. Then*

$$H_{J(G)}^{k+1}(R) \neq 0 \quad \text{for } k = 1, \dots, c_{max}.$$

In particular, the cohomological dimension of $J(G)$ is

$$\text{cd}(J(G), R) = c_{max} + 1.$$

Proof. A local cohomology module is different from zero if it has a non-vanishing Bass number. Then the result follows from Theorem 4.4. \square

Using the relation between the cohomological dimension and the projective dimension of the Alexander dual ideal given in [13] we deduce the following result which, in particular, gives a very simple description of the projective dimension of edge ideals of forests (compare with the results in [24]).

Proposition 4.7. *Let $J(G) \subseteq R$ be an ideal satisfying Assumption 4.3. Then, the projective dimension of the corresponding edge ideal $I(G)$ is*

$$\text{pd}(R/I(G)) = c_{max} + 1.$$

4.2. Non linear injective resolutions. As we have seen through the examples in Section 3, the easiest way to find non linear injective resolutions is to consider the case of cycles. Let's illustrate this fact with the following examples

Example 4.8. Let G be a 6-cycle. We have that all the subgraphs G_α are connected paths except for the cases $\alpha = (1, 1, 0, 1, 1, 0), (1, 0, 1, 1, 0, 1), (0, 1, 1, 0, 1, 1)$ in which G_α has two connected components. Therefore we have the injective resolutions

$$\begin{aligned}
0 \longrightarrow H_{J(G)}^2(R) &\longrightarrow \bigoplus_{|\alpha|=2} E_\alpha \longrightarrow \bigoplus_{|\alpha|=3} E_\alpha \longrightarrow \left(\begin{array}{c} E_{(1,1,0,1,1,0)}^2 \oplus E_{(1,0,1,1,0,1)}^2 \oplus E_{(0,1,1,0,1,1)}^2 \\ \oplus \\ \bigoplus_{\substack{|\alpha|=4 \\ \text{rest of } \alpha' \text{'s}}} E_\alpha \end{array} \right) \longrightarrow
\end{aligned}$$

$$\longrightarrow \begin{array}{c} \bigoplus_{|\alpha|=5} E_\alpha \\ \oplus \\ E_1 \end{array} \longrightarrow E_1 \longrightarrow 0 ,$$

$$0 \longrightarrow H_{J(G)}^3(R) \longrightarrow E_{(1,1,0,1,1,0)} \oplus E_{(1,0,1,1,0,1)} \oplus E_{(0,1,1,0,1,1)} \longrightarrow E_1 \longrightarrow 0 ,$$

where we are only considering those α 's such that $\mathfrak{p}_\alpha \in \text{Supp}R/J(G)$. In this case we have that both injective resolutions have two linear strands.

Example 4.9. Let G be a 6-wheel where x_6 is the dominating vertex. We have that all the subgraphs G_α are connected except for the case $\alpha = (1, 1, 1, 1, 1, 0)$ in which G_α is a 5-cycle. Therefore we have the injective resolutions

$$0 \longrightarrow H_{J(G)}^2(R) \longrightarrow \bigoplus_{|\alpha|=2} E_\alpha \longrightarrow \bigoplus_{|\alpha|=3} E_\alpha \longrightarrow \begin{array}{c} E_{(1,1,1,1,1,0)} \\ \oplus \\ \left(\bigoplus_{|\alpha|=4} E_\alpha \right) \end{array} \longrightarrow \begin{array}{c} E_{(1,1,1,1,1,0)} \\ \oplus \\ \left(\bigoplus_{\substack{|\alpha|=5 \\ a_6=1}} E_\alpha \right) \end{array} \longrightarrow E_1 \longrightarrow 0$$

$$0 \longrightarrow H_{J(G)}^3(R) \longrightarrow E_{(1,1,1,1,1,0)} \longrightarrow 0 ,$$

where we are only considering those α 's such that $\mathfrak{p}_\alpha \in \text{Supp}R/J(G)$. The injective resolution of $H_{J(G)}^2$ has two linear strands.

More generally we can consider the following family of examples.

Assumptions 4.10. Let $J(G) \subseteq R$ be the cover ideal of a simple (not necessarily connected) graph G that is obtained by joining cycles and paths in such a way that we can still find degree two vertices that we can remove in order to simplify the graph.

Even though it is not possible to give a closed formula for the Bass numbers $\mu_p(\mathfrak{p}_\alpha, H_{J(G)}^{k+1}(R))$ as the one given in Theorem 4.4, the methods developed in this work allow us to compute them. To do so we must performe the following steps:

- Describe the connected components of G_α .
- Compute the Lyubeznik numbers of each component as in Section 3.
- Apply Proposition 2.5.

We can also discuss the vanishing of local cohomology modules depending on the connected components of the corresponding subgraphs but the results are not going to be as clean as in Proposition 4.6. Indeed, using Proposition 3.14 and Example 4.1, we deduce the following formula for the case of cycles.

Proposition 4.11. *Let G be a cycle of the form C_{3k-1} . Then*

$$H_{J(G)}^{k+1}(R) \neq 0 \quad \text{for } k = 1, \dots, c_{max} + 1.$$

On the other hand, if G is a cycle of the form C_{3k} or C_{3k+1} we have

$$H_{J(G)}^{k+1}(R) \neq 0 \quad \text{for } k = 1, \dots, c_{max}.$$

Some partial results that we can provide are the following.

Proposition 4.12. *Let G be a graph in n vertices and assume that, given $\alpha \in \{0, 1\}^n$, the subgraph G_α has r connected components having the Lyubeznik type of cycles C_{n_1}, \dots, C_{n_r} and s connected components having trivial Lyubeznik table. Assume that $n_j = 3k_j + \ell_j$ with $\ell_j \in \{-1, 0, 1\}$. Then, if we denote $d_\alpha := |\alpha| - 2 = \dim(R_{\mathfrak{p}_\alpha}/J(G_\alpha))$, we have*

$$H_{J(G_\alpha)}^{d_\alpha - i}(R_{\mathfrak{p}_\alpha}) \neq 0 \quad \text{for } i = 0, \dots, (k_1 + \dots + k_r + s) - 1.$$

In particular, the cohomological dimension of $J(G_\alpha)$ is

$$\text{cd}(J(G_\alpha), R_{\mathfrak{p}_\alpha}) = |\alpha| - (k_1 + \dots + k_r + s) - 1.$$

Proof. Consider the decomposition $G_\alpha = G_1 \cup G_2$ where $\alpha = \alpha_1 + \alpha_2$ and G_1 is the subgraph in $|\alpha_1|$ vertices containing the components corresponding to cycles and G_2 is the subgraph in $|\alpha_2|$ vertices containing the rest. For G_2 , as a consequence of Corollary 2.6, we have

$$H_{J(G_2)}^{d_2 - i}(R_{\mathfrak{p}_{\alpha_2}}) \neq 0 \quad \text{for } i = 0, \dots, s - 1,$$

where $d_2 = |\alpha_2| - 2$.

On the other hand, using Corollary 3.14 and Corollary 2.7, we have that the smallest i for which the Lyubeznik number $\lambda_{p,i}$ corresponding to G_1 is non zero is

$$\begin{aligned} i &= (n_1 - 2) - (k_1 - 1) + \dots + (n_r - 2) - (k_r - 1) + (r - 1) \\ &= (n_1 + \dots + n_r) - 2 - 2(r - 1) - (k_1 + \dots + k_r) + r + (r - 1) \\ &= |\alpha_1| - 2 - (k_1 + \dots + k_r) + 1 \end{aligned}$$

Moreover, since the non-vanishing local cohomology modules of a cycle are consecutive, that is $H_{J(C_n)}^k(R) \neq 0$ where k runs from two to the cohomological dimension, it follows from Proposition 2.5 that the same consecutiveness property holds for G_1 . Namely we have

$$H_{J(G_1)}^{d_1 - i}(R_{\mathfrak{p}_{\alpha_1}}) \neq 0 \quad \text{for } i = 0, \dots, (k_1 + \dots + k_r) - 1,$$

where $d_1 = |\alpha_1| - 2$.

Finally, applying Corollary 2.7 for G_1 and G_2 , we have that the smallest i for which $\lambda_{p,i}(R_{\mathfrak{p}_\alpha}/J(G_\alpha)) \neq 0$ is

$$\begin{aligned} i &= (|\alpha_1| - 2 - (k_1 + \dots + k_r) + 1) + (|\alpha_2| - 2 - (s - 1)) + 1 \\ &= |\alpha_1| + |\alpha_2| - 2 - (k_1 + \dots + k_r + s) + 1 \\ &= |\alpha| - 2 - (k_1 + \dots + k_r + s) + 1 \\ &= d_\alpha - (k_1 + \dots + k_r + s) + 1 \end{aligned}$$

Once again Proposition 2.5 gives the consecutiveness of the non-vanishing local cohomology modules and the result follows. \square

More generally, and using the same type of arguments as above, we have.

Proposition 4.13. *Let G be a graph in n vertices and assume that, given $\alpha \in \{0, 1\}^n$, the subgraph G_α has r connected components having the Lyubeznik type of cycles $C_{n_{1,i}} * \dots * C_{n_{t_j,i}}$, for $i = 1, \dots, r$ and s connected components having trivial Lyubeznik table. Assume that*

$n_{j,i} = 3k_{j,i} + \ell_{j,i}$ with $\ell_{j,i} \in \{-1, 0, 1\}$. Then, if we denote $d_\alpha := |\alpha| - 2 = \dim(R_{\mathfrak{p}_\alpha}/J(G_\alpha))$, we have

$$H_{J(G_\alpha)}^{d_\alpha-i}(R_{\mathfrak{p}_\alpha}) \neq 0 \quad \text{for } i = 0, \dots, (k_{1,1} + \dots + k_{t_1,1} + \dots + k_{1,r} + \dots + k_{t_r,r} + s) - (t_1 + \dots + t_r) + (r-1).$$

REFERENCES

- [1] J. Álvarez Montaner, Characteristic cycles of local cohomology modules of monomial ideals, *J. Pure Appl. Algebra* **150** (2000), 1–25. [2](#), [10](#), [11](#)
- [2] J. Álvarez Montaner, Characteristic cycles of local cohomology modules of monomial ideals II, *J. Pure Appl. Algebra* **192** (2004), 1–20. [2](#)
- [3] J. Álvarez Montaner, *Some numerical invariants of local rings*, *Proc. Amer. Math. Soc.* **132** (2004), 981–986. [6](#)
- [4] J. Álvarez Montaner, *Operations with regular holonomic D-modules with support a normal crossing*, *J. Symb. Comp.* **40**, (2005), 999–1012. [6](#)
- [5] J. Álvarez Montaner, Local cohomology modules supported on monomial ideals, *Lecture Notes in Math.* **2083** (2013), 109–178. [4](#)
- [6] J. Álvarez Montaner and O. Fernández-Ramos, Local cohomology using Macaulay2, *Lecture Notes in Math.* **2083** (2013), 179–185. [2](#), [26](#)
- [7] J. Álvarez Montaner, R. García López and S. Zarzuela, *Local cohomology, arrangements of subspaces and monomial ideals*, *Adv. Math.* **174** (2003), 35–56. [5](#)
- [8] J. Álvarez Montaner and A. Vahidi, *Lyubeznik numbers of monomial ideals*, *Trans. Amer. Math. Soc.* **366** (2014), 1829–1855. [2](#), [6](#), [8](#), [10](#), [11](#), [27](#)
- [9] J. Álvarez Montaner and K. Yanagawa, *Lyubeznik numbers of local rings and linear strands of graded ideals*, *Nagoya Math. J.* **231** (2018), 23–54. [7](#), [8](#)
- [10] E. Batzies and V. Welker, *Discrete Morse theory for cellular resolutions*, *J. Reine Angew. Math.* **543** (2002), 147–168. [2](#)
- [11] W. Bruns and J. Herzog, *Cohen–Macaulay Rings*, revised edition, *Cambridge Studies in Advanced Mathematics*, vol. 39, Cambridge University Press, Cambridge, 1998. [1](#), [4](#)
- [12] A. Corso and U. Nagel, *Monomial and toric ideals associated to Ferrers graphs*, *Trans. Amer. Math. Soc.* **361** (2009), 1371–1395. [11](#)
- [13] J. A. Eagon and V. Reiner, *Resolutions of Stanley-Reisner rings and Alexander duality*, *J. Pure Appl. Algebra*, **130** (1998), 265–275. [11](#), [28](#)
- [14] S. Eliahou and M. Kervaire, *Minimal resolutions of some monomial ideals*, *J. Algebra* **129** (1990), 1–25. [2](#), [8](#)
- [15] C. Francisco, H. T. Hà and A. Van Tuyl, *Splittings of monomial ideals*, *Proc. Amer. Math. Soc.* **137** (2009), 3271–3282. [2](#), [8](#), [9](#)
- [16] R. Fröberg, *On Stanley-Reisner rings*. *Topics in algebra*, Part 2 (Warsaw, 1988), 57–70, Banach Center Publ., 26, Part 2, PWN, Warsaw, 1990. [11](#)
- [17] S. Goto and K. Watanabe, *On Graded Rings, II (\mathbb{Z}^n -graded rings)*, *Tokyo J. Math.* **1**(2) (1978), 237–261. [5](#)
- [18] D. Grayson and M. Stillman, *Macaulay 2*, <http://www.math.uiuc.edu/Macaulay2>. [2](#)
- [19] H. T. Hà and A. Van Tuyl, *Splittable ideals and the resolutions of monomial ideals*, *J. Algebra* **309** (2007), 405–425. [13](#)
- [20] M. Hochster, *Cohen-Macaulay rings, combinatorics, and simplicial complexes*, *Ring theory, II* (Proc. Second Conf., Univ. Oklahoma, Norman, Okla., 1975), pp. 171–223. *Lecture Notes in Pure and Appl. Math.*, Vol. 26, Dekker, New York, 1977. [2](#)
- [21] M. Hochster and C. Huneke, *Indecomposable canonical modules and connectedness*, *Commutative algebra: syzygies, multiplicities, and birational algebra* (South Hadley, MA, 1992), *Contemp. Math.* **159** (1994), 197–208, [6](#)
- [22] C. Huneke and R. Y. Sharp, *Bass numbers of local cohomology modules*, *Trans. Amer. Math. Soc.* **339** (1993), 765–779. [6](#)
- [23] S. Jacques, *Betti Numbers of Graph Ideals*, Ph.D. Thesis. University of Sheffield. 2004. Available at [arXiv:math/0410107](https://arxiv.org/abs/math/0410107). [11](#)

- [24] S. Jacques and M. Katzman, *The Betti numbers of forests*, Preprint available at arXiv:0501226. [28](#)
- [25] G. Lyubeznik, *The minimal non-Cohen-Macaulay monomial ideals*, J. Pure Appl. Algebra **51** (1988), 261–266. [11](#)
- [26] G. Lyubeznik, *On some local cohomology invariants of local rings*, Math. Z. **254** (2006), 627–640. [6](#)
- [27] G. Lyubeznik, *Finiteness properties of local cohomology modules (an application of D-modules to commutative algebra)*, Invent. Math., **113** (1993) 41–55. [2](#), [6](#)
- [28] E. Miller, *Resolutions and duality for monomial ideals*, Ph.D. Thesis. University of California, Berkeley. 2000. 136 pp. ISBN: 978-0599-86011-7 [2](#)
- [29] E. Miller, *The Alexander duality functors and local duality with monomial support*, J. Algebra **231** (2000), 180–234. [2](#), [4](#), [8](#)
- [30] E. Miller and B. Sturmfels, *Combinatorial commutative algebra*, Graduate Texts in Mathematics **227**. Springer-Verlag, New York, 2005. [4](#), [8](#)
- [31] M. Mustața, *Local Cohomology at Monomial Ideals*, J. Symbolic Comput. **29** (2000), 709–720. [2](#), [5](#)
- [32] R. P. Stanley, *Combinatorics and Commutative Algebra, Second Edition*, Progress in Mathematics **41**, Birkhäuser, 1996. [1](#)
- [33] N. Terai, *Local cohomology modules with respect to monomial ideals*, Preprint 1999 [5](#)
- [34] K. Yanagawa, *Bass numbers of local cohomology modules with supports in monomial ideals*, Math. Proc. Cambridge Philos. Soc. **131** (2001), 45–60. [2](#), [4](#), [5](#)
- [35] W. Zhang, *On the highest Lyubeznik number of a local ring*, Compos. Math. **143** (2007), 82–88. [6](#)

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT POLITÈCNICA DE CATALUNYA, AV. DIAGONAL 647,
BARCELONA 08028, SPAIN

E-mail address: Josep.Alvarez@upc.edu

FACULTY OF SCIENCES, DEPARTAMENT OF MATHEMATICS, UNIVERSITY OF MOHAGHEGH ARDABIL,
56199-11367 ARDABIL, IRAN

E-mail address: fateme_sohrabi@uma.ac.ir