

Metric dimension of maximal outerplanar graphs

M. Claverol · A. García · G.
Hernández · C. Hernando · M.
Maureso · M. Mora · J. Tejel

Abstract In this paper we study the metric dimension problem in maximal outerplanar graphs. Concretely, if $\beta(G)$ denotes the metric dimension of a maximal outerplanar graph G of order n , we prove that $2 \leq \beta(G) \leq \lceil \frac{2n}{5} \rceil$ and that the bounds are tight. We also provide linear algorithms to decide whether the metric dimension of G is 2 and to build a resolving set S of size $\lceil \frac{2n}{5} \rceil$ for G . Moreover, we characterize all maximal outerplanar graphs with metric dimension 2.

Keywords Metric dimension · Resolving set · Maximal outerplanar graph

Mathematics Subject Classification (2010) 05C12 · 05C10

M. Claverol
Departament de Matemàtiques, Universitat Politècnica de Catalunya, Barcelona, Spain
E-mail: merce.claverol@upc.edu, ORCID: 0000-0002-9138-8594

A. García
Departamento de Métodos Estadísticos, IUMA, Universidad de Zaragoza, Zaragoza, Spain
E-mail: olaverri@unizar.es, ORCID: 0000-0002-6519-1472

G. Hernández
Departamento de Matemática Aplicada a las TIC, Universidad Politécnica de Madrid,
Madrid, Spain
E-mail: gregorio@fi.upm.es, ORCID: 0000-0002-7178-0249

C. Hernando
Departament de Matemàtiques, Universitat Politècnica de Catalunya, Barcelona, Spain
E-mail: carmen.hernando@upc.edu, ORCID: 0000-0002-3864-6566

M. Maureso
Departament de Matemàtiques, Universitat Politècnica de Catalunya, Barcelona, Spain
E-mail: montserrat.maureso@upc.edu, ORCID: 0000-0001-6429-2776

M. Mora
Corresponding author
Departament de Matemàtiques, Universitat Politècnica de Catalunya, Barcelona, Spain
E-mail: merce.mora@upc.edu, ORCID: 0000-0001-6923-0320

J. Tejel
Departamento de Métodos Estadísticos, IUMA, Universidad de Zaragoza, Zaragoza, Spain
E-mail: jtejel@unizar.es, ORCID: 0000-0002-9543-7170

1 Introduction

Let $G = (V, E)$ be a finite connected simple graph. For two vertices $u, v \in V$, let $d(u, v)$ denote the length of a shortest path in G from u to v . If $S = \{x_1, \dots, x_k\}$ is a set of vertices of G , we denote by $r(u|S)$ the vector of distances from u to the vertices of S , that is, $r(u|S) = (d(u, x_1), \dots, d(u, x_k))$. We say that a vertex $x \in V$ *resolves* a pair of vertices $u, v \in V$ if $d(u, x) \neq d(v, x)$. A set of vertices $S \subseteq V$ is a *resolving set* of G if every pair of distinct vertices of G are resolved by some vertex in S . Therefore, S is a resolving set if and only if $r(u|S) \neq r(v|S)$ for every pair of distinct vertices $u, v \in V(G)$. The elements of $r(u|S)$ are the *metric coordinates* of u with respect to S . A resolving set S of G with minimum cardinality is a *metric basis* of G . The *metric dimension* of G , denoted by $\beta(G)$, is the cardinality of a metric basis. The *metric dimension problem* consists of finding a metric basis.

Resolving sets in general graphs were first studied by Slater [21] and Harary and Melter [16]. Since then, computing resolving sets and the metric dimension of a graph have been widely studied in the literature due to their applications in several areas, such as network discovery and verification [1], robot navigation [18], chemistry [5] or games [6]. The reader is referred to [3, 4, 10–15, 17, 19, 25] and the references therein for different results and variants of the metric dimension problem of graphs.

It is well-known that the metric dimension problem in general graphs is NP-hard [18]. The problem remains NP-hard even when restricting to some graph classes such as bounded-degree planar graphs [7]; split graphs, bipartite graphs and their complements, and line graphs of bipartite graphs [9]; interval graphs and permutation graphs of diameter 2 [11]. Polynomial algorithms are known for trees [18]; outerplanar graphs [7]; chain graphs [10]; k -edge-augmented trees, cographs and wheels [9]. A weighted variant of the metric dimension problem in several graphs, including paths, trees, and cographs, can be also solved in polynomial time [9].

While the algorithms to solve the metric dimension problem in trees, wheels or chain graphs are linear, the time complexity of the algorithm given in [7] for an outerplanar graph of order n is $O(n^{12})$. Thus, an interesting problem for such graphs is how to find more efficiently a not very large resolving set. Recall that a graph G is *outerplanar* if it can be drawn in the plane without crossings and with all the vertices belonging to the unbounded face.

In this paper, we focus on studying the metric dimension problem in *maximal outerplanar graphs*. A maximal outerplanar graph, *MOP graph* for short,

is an outerplanar graph such that the addition of an edge produces a non outerplanar graph. In particular, given a MOP graph G of order $n \geq 3$ we show that $2 \leq \beta(G) \leq \lceil \frac{2n}{5} \rceil$ and that the bounds are tight. The lower bound is shown to be tight in Section 2. Moreover, all MOP graphs with metric dimension 2 are characterized. We also provide in that section a linear algorithm to decide whether the metric dimension of a MOP graph is 2. The tightness of the upper bound is shown in Section 3.1 by exhibiting a family of MOP graphs attaining the given bound. Section 3.2 is devoted to show that the metric dimension of a MOP graph G is at most $\lceil \frac{2n}{5} \rceil$, by building in linear time a resolving set S for G such that $|S| = \lceil \frac{2n}{5} \rceil$. In [20], it is conjectured that $\beta(G) \leq \lceil \frac{2n}{5} \rceil$ for a maximal planar graph G , hence we are answering in the affirmative this conjecture for the particular case of MOP graphs. We conclude the paper with some open questions in Section 4. An extended abstract of this work has appeared at the 17th Spanish Meeting on Computational Geometry (EGC 2017).

To finish this section, we recall some well-known properties of MOP graphs. A MOP graph G of order at least 3 is biconnected, Hamiltonian and always admits a plane embedding such that all vertices belong to the unbounded face and every bounded face is a triangle. Unless otherwise stated, we assume throughout the paper that the MOP graph has order at least 3 and we are given this plane embedding of G . Thus, G can be seen as a triangulation of a convex polygon. Every edge on the boundary of the unbounded face belongs to only one triangle of G and any other edge (called diagonal) belongs to two triangles. The removal of the endvertices of a diagonal makes the graph to be disconnected. G always has at least 2 vertices of degree 2 and when removing any of them (if $|G| \geq 4$), the resulting graph is a MOP graph. From these properties, it is straightforward to see the following result:




Fig. 1 (a) Given a 2-tree G with metric dimension 2, a minimal induced 2-connected subgraph containing the basis $\{a_1, a_k\}$, as claimed in [2]. (b) A 2-tree with metric basis $\{a_1, a_k\}$ whose minimal induced 2-connected subgraph containing a_1 and a_k is different from the claimed subgraph in [2]

Remark 1 Let G be a MOP graph and let xy be a diagonal of G . If u and z are two vertices belonging to different components of $G \setminus \{x, y\}$, then $d(u, z) \geq \min\{d(u, x), d(u, y)\} + 1$.

2 MOP graphs with metric dimension two

Given a MOP graph G , its metric dimension must be greater than one, as paths are the only graphs with metric dimension one (see for example [5]). In this section, we characterize MOP graphs with metric dimension two.

There are several papers in the literature devoted to study properties of graphs with metric dimension two and to characterize such graphs for certain families of graphs. In [23], the authors give a general characterization for a graph G to have metric dimension two, based on the distance partition $\{U_1, U_2, \dots, U_k\}$ of the vertices of G , where vertices belonging to U_i are at distance i from a distinguished vertex v . They also give a $O(n^2 D^4)$ algorithm to check whether the metric dimension of a graph of order n is two, where D is the diameter of the graph. In [18], the authors show several properties that a graph with metric dimension two must satisfy.

Graphs with metric dimension two have been characterized for some families of graphs. In particular, unicyclic graphs [8] and Cayley graphs [24]. An incorrect characterization of the 2-trees with metric dimension 2 is given in [2]. Starting with a triangle, a 2-tree is formed by repeatedly adding vertices of degree 2 in such a way that each added vertex u is connected to two vertices v and w which are already adjacent. Thus, the family of 2-trees includes MOP graphs as a subfamily.

In [2], the authors define a family \mathcal{F} of 2-trees such that a 2-tree G belongs to \mathcal{F} if G satisfies a set of twelve conditions, and they claim that a 2-tree G has metric dimension 2 if and only if G belongs to \mathcal{F} . When proving that a 2-tree G with metric dimension two must belong to \mathcal{F} , the authors claim in one of the cases that the shape of the minimal induced 2-connected subgraph of G , containing the two vertices a_1 and a_k of the basis of G , is as shown in Figure 1(a): Two vertices of degree two (a_1 and a_k), two vertices of degree three (b_1 and b_k), a set of quadrilaterals with one of the two possible diagonals, and at most one vertex of degree five in the path a_1, a_2, \dots, a_k . But, part (b) of Figure 1 exhibits a 2-tree G (in fact a MOP graph) with metric dimension two, being $\{a_1, a_k\}$ the only basis of G , and the minimal induced 2-connected subgraph of G containing a_1 and a_k is precisely G , contradicting the shape claimed in [2]. As a consequence, their claimed characterization cannot be used to characterize MOP graphs with metric dimension 2.

We next give a characterization for MOP graphs with metric dimension two, based on embedding graphs with metric dimension 2 into the strong product of two paths. The strong product of two paths of order n , $P_n \boxtimes P_n$, has the cartesian product $[0, n-1] \times [0, n-1]$ as set of vertices and two different vertices (i, j) and (i', j') are adjacent if and only if $|i' - i| \leq 1$ and $|j' - j| \leq 1$. The distance between two vertices of this graph is $d((i, j), (i', j')) = \max\{|i' - i|, |j' - j|\}$. We will consider the representation of this graph in the plane identifying vertex (i, j) with the point with cartesian coordinates (i, j) . In this representation, a path of length k between two vertices (i, j) and (i', j') such that $d((i, j), (i', j')) = k$ is contained in the rectangle having (i, j) and (i', j') as opposite vertices and sides parallel to lines of slope 1 and -1




Fig. 2 Left: A graph G with metric basis $S = \{1, 3\}$. The metric coordinates of the vertices are: $r(1|S) = (0, 2)$, $r(2|S) = (1, 1)$, $r(3|S) = (2, 0)$, $r(4|S) = (3, 1)$, $r(5|S) = (2, 1)$, $r(6|S) = (3, 2)$, $r(7|S) = (3, 3)$, $r(8|S) = (2, 3)$, $r(9|S) = (1, 3)$, $r(10|S) = (1, 2)$ and $r(11|S) = (2, 2)$. Right: The representation G^* of G as a subgraph of $P_n \boxtimes P_n$ with respect to S . Vertex v in G is mapped to vertex v^* in G^* such that the cartesian coordinates of v^* are the metric coordinates of v

passing through these vertices. A set of four vertices of $P_n \boxtimes P_n$ of the form $\{(i, j), (i, j + 1), (i + 1, j + 1), (i + 1, j)\}$, for some $i, j \in [0, D]$, is a *unit square*. Three vertices of $P_n \boxtimes P_n$ are pairwise adjacent if and only if they all belong to a unit square and, in such a case, the edges joining them form a triangle with two consecutive sides one of a unit square and the diagonal joining them.

Let G be a graph with metric dimension 2 and let $S = \{u, v\}$ be a metric basis of G . It is straightforward to see that G is isomorphic to a subgraph of the strong product $P_n \boxtimes P_n$. See Figure 2 for an example. Indeed, we identify vertex $x \in V(G)$ with vertex $(x_1, x_2) \in V(P_n \boxtimes P_n)$, where $(x_1, x_2) = r(x|S) = (d(x, u), d(x, v))$. Recall that if two vertices w_1 and w_2 of G are adjacent and $d(w_0, w_1) = d$ for some vertex w_0 , then $d(w_0, w_2) \in \{d - 1, d, d + 1\}$. Thus, if x and y are adjacent vertices in G , then $|d(x, u) - d(y, u)| \leq 1$ and $|d(x, v) - d(y, v)| \leq 1$, hence $r(x|S)$ and $r(y|S)$ are adjacent in $P_n \boxtimes P_n$. We denote by G^* this representation of G , that is, $V(G^*) = \{r(x|S) : x \in V(G)\}$ and $r(x|S)r(y|S) \in E(G^*)$ if and only if $xy \in E(G)$. We say that G^* is the *representation of G as a subgraph of $P_n \boxtimes P_n$ with respect to S* , and vertex (i, j) is placed onto the point with cartesian coordinates (i, j) .

For every $d \geq 1$, consider the set $A_d = \{(i, j) \in [0, n - 1] \times [0, n - 1] : i + j \geq d, |j - i| \leq d\}$ (see Figure 3 left). The following properties can be easily derived.

Proposition 1 *Let G be a graph with metric dimension 2, and let $S = \{u, v\}$ be a metric basis of G such that $d(u, v) = d$. Consider the representation G^* of G as a subgraph of $P_n \boxtimes P_n$ with respect to S . The following properties hold:*

- (1) $S^* = \{(0, d), (d, 0)\}$ is a metric basis of G^* and all the vertices of G^* are in A_d .
- (2) There is only one shortest $((0, d), (d, 0))$ -path in G^* , and its vertices are the points (i, j) such that $i + j = d$.
- (3) Three vertices of G^* are pairwise adjacent if and only if they all belong to a unit square.




Fig. 3 Left: Illustrating Proposition 1. The shortest path from $(0, d)$ to $(d, 0)$ and the set A_d , which is in the shaded region. Right: Examples of horizontal and vertical MOP zigzags

Proof (1) If $d = d(u, v)$, then $r(u|S) = (0, d)$ and $r(v|S) = (d, 0)$. If $x \in V(G)$, then $r(x|S) = (x_1, x_2) = (d(x, u), d(x, v))$. On the one hand, $x_1 = d(x, u) \leq d(x, v) + d(u, v) = d(x, v) + d = x_2 + d$ and $x_2 = d(x, v) \leq d(x, u) + d(u, v) = d(x, u) + d = x_1 + d$, hence $|x_1 - x_2| \leq d$. On the other hand, $x_1 + x_2 = d(x, u) + d(x, v) \geq d(u, v) = d$.

(2) There is only one path of length d joining $(0, d)$ and $(d, 0)$ in $P_n \boxtimes P_n$, and its vertices are $\{(i, j) : i + j = d\}$. Thus, it is also the only shortest path between $(0, d)$ and $(d, 0)$ in G^* because we already know that $d_{G^*}((0, d), (d, 0)) = d$.

(3) It is also obvious, because three pairwise adjacent vertices of $P_n \boxtimes P_n$ belong to a unit square. \square

For $n \geq 5$, we say that a MOP graph G is a *MOP zigzag* if G has two vertices of degree 2, two vertices of degree 3, each one of them adjacent to a different vertex of degree 2, and the rest of the vertices have degree 4. See Figure 3 right for some examples of MOP zigzags. One can see a MOP zigzag as a MOP graph in which the diagonals form a zigzag path connecting the two vertices of degree 3. For $n = 3, 4$, we consider a triangle and a quadrilateral with a diagonal as MOP zigzags, respectively.

Given the representation G^* of a graph G , we say that an edge $e \in E(G^*)$ is *horizontal* if $e = (i, j)(i + 1, j)$, for some $i, j \geq 0$; *vertical* if $e = (i, j)(i, j + 1)$, for some $i, j \geq 0$; *1-slope diagonal* if $e = (i, j)(i + 1, j + 1)$, for some $i, j \geq 0$; and *(-1)-slope diagonal* if $e = (i, j + 1)(i + 1, j)$, for some $i, j \geq 0$. A *vertical MOP zigzag* with base line a vertical edge $(i, j)(i, j + 1)$ (see Figure 3 right) is a subgraph of the strong product induced by the set of vertices $\{(i + k, j + k) : 0 \leq k \leq r\} \cup \{(i + k, j + 1 + k) : 0 \leq k \leq s\}$, for some $r \geq 1$ and $s \in \{r - 1, r\}$, and a *horizontal MOP zigzag* with base line a horizontal edge $(i, j)(i + 1, j)$ is a subgraph of the strong product induced by the set of vertices $\{(i + k, j + k) : 0 \leq k \leq r\} \cup \{(i + 1 + k, j + k) : 0 \leq k \leq s\}$, for some $r \geq 1$ and $s \in \{r - 1, r\}$.

For any integer $k \geq 1$, let $V_k = \{(i, j) : i + j = k\}$. The following theorem characterizes the MOP graphs with metric dimension 2. Any of these MOP graphs consists of a base graph similar to the one shown in Figure 4(c) and several MOP zigzags joined to this base graph (see Figure 4(d)).

Theorem 1 *Let G be a MOP graph. Then, $\beta(G) = 2$ if and only if there is a representation G^* of G as a subgraph of the strong product of two paths such that for some $d \geq 1$,*

- (1) $V(G^*) \subseteq A_d$, $V_d \cap A_d \subseteq V(G^*)$, and $E(G^*)$ contains the edges of the shortest path joining $(0, d)$ and $(d, 0)$.
- (2) $V_{d+1} \cap A_d \subseteq V(G^*)$ and for each $(i, j) \in V_{d+1} \cap A_d$, $E(G^*)$ contains the edges $(i, j)(i-1, j)$ and $(i, j)(i, j-1)$.
- (3) For every pair of vertices $(i, j+1)$ and $(i+1, j)$ of V_{d+1} with $i, j \geq 1$, we have either $(i, j+1)(i+1, j) \in E(G^*)$ or $\{(i, j+1)(i+1, j+1), (i+1, j)(i+1, j+1), (i, j)(i+1, j+1)\} \subseteq E(G^*)$. Moreover, if $(i, j+1)(i+1, j) \in E(G^*)$ belongs to two triangles of G^* , then $(i+1, j+1) \in V(G^*)$ and $\{(i, j+1)(i+1, j+1), (i+1, j)(i+1, j+1)\} \subseteq E(G^*)$.
- (4) Any other vertex or edge of the graph belongs to a vertical or horizontal MOP zigzag with base line the edge $(0, d)(1, d)$, or the edge $(d, 0)(d, 1)$, or any other edge of G from those described in the preceding items with an endpoint in V_{d+1} and the other in V_{d+2} , with the additional condition that two distinct maximal vertical or horizontal MOP zigzags do not share any edge.

Proof Let us see first that if a MOP graph has metric dimension 2, then it satisfies items (1)-(4). Item (1) is a consequence of Proposition 1 (see Figure 4(a)).

Let us prove now (2). Recall that every edge of a MOP graph belongs to at least one triangle. Let $(i, j-1)(i-1, j)$ be an edge of the $((0, d), (d, 0))$ -path (and thus, $i+j-1 = d$). The only triangle of the strong product with vertices in A_d containing this edge is that with vertices $(i, j-1)$, $(i-1, j)$ and (i, j) . From here, the second item follows (see Figure 4(b)).




Fig. 4 An example of a MOP graph G with metric dimension 2. If the vertices of a basis are at distance d , then G can be represented as a subgraph G^* of the strong product $P_n \boxtimes P_n$ such that all vertices of G^* belong to the shaded region. Vertices described in Theorem 1 (1), (2), (3) and (4) are added in (a), (b), (c) and (d), respectively. Observe that all vertices of G^* belong to the unbounded face

To prove item (3), take $(i, j) \in V_d$. By item (2), we know that $(i, j)(i, j+1)$ and $(i, j)(i+1, j)$ are edges of G^* . Notice that the edges of the shortest

$((0, d), (d, 0))$ -path belong to exactly one triangle of G^* , thus, any other edge incident to $(i, j) \in V_d$ belongs to two triangles of G^* . Therefore, the edges $(i, j)(i, j+1)$ and $(i, j)(i+1, j)$ belong to two triangles of G^* , and there are only two possibilities, either $(i, j+1)(i+1, j) \in E(G^*)$ or $\{(i, j+1)(i+1, j+1), (i+1, j)(i+1, j+1), (i, j)(i+1, j+1)\} \subseteq E(G^*)$. In addition, if $(i, j+1)(i+1, j) \in E(G^*)$ belongs to two triangles, the only possibility is that $(i+1, j+1) \in V(G^*)$ and $\{(i, j+1)(i+1, j+1), (i+1, j)(i+1, j+1)\} \subseteq E(G^*)$ (see Figure 4(c)).

Finally, let us prove item (4). Let t be the number of triangles of the MOP graph G . The vertices and edges described in the preceding items (1), (2) and (3) induce a MOP graph, G_0^* , with t_0 triangles.

If $t = t_0$, then $G_0^* = G^*$ and we are done. Suppose now that $t > t_0$. In such a case, one of the edges of G_0^* limiting only one triangle in G_0^* must belong to two triangles in G^* . Let $e_0 = xy$ be one of these edges and let z be the third vertex of the triangle in G_0^* containing the endpoints of e_0 . By definition of G_0^* , e_0 must be a horizontal edge or a vertical edge. Besides, $(0, d)$, $(d, 0)$ and z belong to the same component in $G^* - \{x, y\}$. Assume that $e_0 = (i, j)(i+1, j)$ if e_0 is a horizontal edge, and $e_0 = (i, j)(i, j+1)$ if e_0 is a vertical edge, with $i, j \geq 0$. By Remark 1, we have that the third vertex of the other triangle of G^* limited by e_0 must be $(i+1, j+1)$.

Let G_1^* be the graph obtained by adding to the graph G_0^* the vertex $(i+1, j+1)$ and the edges joining $(i+1, j+1)$ with the endpoints of e_0 . Notice that one of the edges added to G_0^* is a 1-slope diagonal edge, and the other one is a horizontal edge if e_0 is vertical, or a vertical edge if e_0 is horizontal.

Now, if $G^* = G_1^*$, we are done. Otherwise, there is an edge e_1 belonging to exactly one triangle in G_1^* and to two triangles in G^* . By Remark 1, there is no 1-slope diagonal edge $(i, j)(i+1, j+1)$, with $i+j \geq d+1$, limiting two triangles in G^* . Hence, e_1 must be a horizontal edge or a vertical edge and we proceed as for e_0 . We repeat this procedure until we have added $t - t_0$ triangles to G_0^* . Observe that the new triangles added to G_0^* form a vertical or horizontal MOP zigzag with one of the considered base lines, since the triangles recursively added to G_0^* share vertical or horizontal edges.

Finally, it is not possible that two maximal vertical or horizontal MOP zigzags share an edge e . Indeed, in such a case, the edge e should be a 1-slope diagonal edge $e = (i, j)(i+1, j+1)$, with $i+j \geq d+1$, and $G^* - \{(i, j), (i+1, j+1)\}$ would be connected, a contradiction because e is not an edge of the unbounded face (see Figure 4(d)).

Now, we are going to prove that every graph satisfying (1) to (4) is a MOP graph with metric dimension 2. By construction, a graph satisfying conditions (1)-(4) is a biconnected plane graph with all vertices belonging to the unbounded face and any other face is a triangle. Therefore, G is a MOP graph. Moreover, $d((0, d), (i, j)) = i$ and $d((d, 0), (i, j)) = j$. Indeed, it is easy to give a path of length i from $(0, d)$ to (i, j) using some vertices of the shortest $(0, d) - (d, 0)$ path; all the vertices (i', j') such that $i' \leq i$, $j' \leq j$ and $i' - j' = i - j$; and vertex $((d+i-j)/2, 1+(d-(i-j))/2) \in V_{d+1}$, whenever d and $i-j$ have the same parity and with $i-j \neq -d$ (see Figure 5). In a similar




Fig. 5 A $(0, d) - (i, j)$ path of length i when $i - j$ and d have distinct parity (left); when $i - j$ and d have the same parity and $i - j \neq -d$ (center) and when $i - j = -d$ (right)

way, a path of length j from $(d, 0)$ to (i, j) can be given. Thus $\{(0, d), (d, 0)\}$ is a resolving set. Since G is not a path, we have $\beta(G) = \beta(G^*) = 2$. \square

If G is a MOP graph with metric dimension 2, we denote by G_0^* the graph induced by the vertices and edges described in items (1)-(3) of Theorem 1.

Deciding whether the metric dimension of a MOP graph is 2 can be done in linear time, as the following theorem shows.

Theorem 2 *Given a MOP graph G of order n we can decide in linear time and space whether the metric dimension of G is 2.*

Proof It is obvious for $n = 3$. From now on, suppose that $n \geq 4$.

If G is a MOP zigzag, then one can easily check that its metric dimension is 2, since two of the four vertices of degree 2 and 3 chosen in a suitable way form a resolving set. Thus, we may assume that G is not a MOP zigzag and we may also assume that the vertices of G are clockwise ordered along its boundary. From Theorem 1, the representation of a MOP graph G with metric dimension 2 consists of the graph G_0^* together with some vertical and horizontal MOP zigzags joined to G_0^* . Note that every vertical or horizontal MOP zigzag finishes in a vertex of degree 2 in G .

Given G , in the first step of the algorithm we calculate for every vertex v of degree 2 the *maximal MOP zigzag around v* , denoted by G_v . The set of vertices of G_v is the maximal set of consecutive vertices $S_v = \{u, \dots, v, \dots, w\}$ of G around v such that the subgraph induced by S_v is a MOP zigzag. By definition, uw is an edge of G_v , that will be denoted by e_v . This subgraph can be calculated by alternately exploring the vertices preceding and following v (see Figure 6).

Since calculating G_v only depends on its size, and two maximal MOP zigzags around two vertices v and v' of degree 2 are edge-disjoint, this first step only requires linear time and space. Notice that if G has metric dimension 2 and v is a vertex of degree 2, then the edge $e_v = uw$ of G_v must be a vertical or horizontal edge on the boundary of G_0^* , or a (-1) -slope diagonal edge of G_0^* . This last case (see for example the fourth MOP zigzag when moving along the border of G_0^* from $(0, d)$ in Figure 4(d)) only happens when the triangle defined by the vertices $(i, j + 1)$, $(i + 1, j)$ and $(i + 1, j + 1)$ belongs to G_0^* , with




Fig. 6 Examples of maximal MOP zigzags around v . On the left, vertices x and u are nonadjacent and, on the right, vertices x and w are nonadjacent

$(i, j + 1), (i + 1, j) \in V_{d+1}$, and only one of the edges $(i, j + 1)(i + 1, j + 1)$ and $(i + 1, j)(i + 1, j + 1)$ is the base line for a MOP zigzag that contains v .

Let $S = \{u_1, u_2, \dots, u_k\}$ be the set of vertices of degree 2 or 3 of G , clockwise ordered when moving along the boundary of G . From Theorem 1, we deduce that, if G has metric dimension 2, then a metric basis of G is formed by two consecutive vertices u_i and u_{i+1} in S (where $u_{k+1} = u_1$). Thus, in the second step of the algorithm, we check if the set $\{u_i, u_{i+1}\}$ is a metric basis of G , for every $i \in \{1, \dots, k\}$.

Given a pair (u_i, u_{i+1}) , this can be done as follows. Suppose that there are $d_i - 1$ vertices between u_i and u_{i+1} when traveling clockwise along the boundary of G . Note that using these vertices, checking (and building) if a graph G_0^* as described in items (1), (2) and (3) of Theorem 1 exists can be done in $O(d_i)$ time and space. If such a graph G_0^* exists, the rest of the vertices of G^* must belong to vertical and horizontal MOP zigzags joined to G_0^* . This can be again checked in $O(d_i)$ time by visiting clockwise the edges on the boundary of G_0^* . Indeed, an edge e_v associated with a vertex of degree 2 of those calculated in the first step must be a vertical, horizontal or (-1) -slope diagonal edge of G_0^* . Besides, it can be checked if all vertices of G appear in G_0^* or in a maximal MOP zigzag joined to G_0^* , since the number of vertices of G_0^* and of each maximal MOP zigzag is known. Therefore, as $\sum d_i = n$, this second step also requires linear time and space. \square

3 Upper bound on the metric dimension of MOP graphs

In this section, we show that $\beta(G) \leq \lceil \frac{2n}{5} \rceil$ for any MOP graph G of order n . We also show that, for some special MOP graphs of order n , their metric dimension is $\lceil \frac{2(n-2)}{5} \rceil$. Hence, the upper bound $\lceil \frac{2n}{5} \rceil$ is tight when n is a multiple of 5.

In the figures, we will assume that the vertices of a MOP graph G are placed on a circle labeled clockwise from 1 to n . The edges will be drawn on or inside the circle as segments or arcs.




Fig. 7 The fan $F_{1,6}$. Black vertices form a metric basis S . For a vertex not in S , its metric coordinates are given

3.1 Fan graphs

We first study the metric dimension of a special family of MOP graphs, the fan graphs. A *fan graph* of order n , denoted by $F_{1,n-1}$, is a MOP graph such that one of the vertices is connected to the $n - 1$ remaining vertices. For $n = 3, 4, 5, 6$, one can easily verify that $\beta(F_{1,n-1}) = 2$. For $n = 7$, we have $\beta(F_{1,6}) \geq 3$. This result follows from the fact that $n \leq \beta + D^\beta$ for a graph with metric dimension β and diameter D (see [18]). As $F_{1,6}$ has diameter 2, if it had metric dimension 2, then n would be at most 6. In addition, the three black vertices of Figure 7 form a metric basis for $F_{1,6}$, so $\beta(F_{1,6}) = 3$.

In the following theorem, we prove that $\beta(F_{1,n-1}) = \lceil \frac{2(n-2)}{5} \rceil$, for $n \geq 8$. The proof is based on locating-dominating sets. Given a graph $G = (V, E)$, let $N(u)$ be the set of neighbors of u in G , that is, $N(u) = \{v : uv \in E(G)\}$. A set $S \subseteq V$ is a *dominating set* if every vertex not in S is adjacent to some vertex in S . A set $S \subseteq V$ is a *locating-dominating set*, if S is a dominating set and $N(u) \cap S \neq N(v) \cap S$ for every two different vertices u and v not in S . The *location-domination number* of G , denoted by $\lambda(G)$, is the minimum cardinality of a locating-dominating set. It is easy to show that any locating-dominating set is a resolving set. Thus, $\beta(G) \leq \lambda(G)$.

Theorem 3 *Let $n \geq 8$. Then,*

$$\beta(F_{1,n-1}) = \left\lceil \frac{2(n-2)}{5} \right\rceil.$$

Proof Observe that $\beta(F_{1,n-1}) \geq 3$, because $F_{1,n-1}$ is not a path and graphs with metric dimension 2 and diameter 2 have order at most 6. Suppose that the vertices of $F_{1,n-1}$ are labeled so that n is the vertex of degree $n - 1$, and let P be the path of order $n - 1$ induced by vertices from 1 to $n - 1$.

We first prove that $\beta(F_{1,n-1}) \leq \lceil \frac{2(n-2)}{5} \rceil$. In [22], it is shown that a path of order $n - 2$ has a locating-dominating set of size $\lceil \frac{2(n-2)}{5} \rceil$ such that at least one endpoint of the path does not belong to it. Using this fact, we derive that the path of order $n - 2$ induced by the vertices from 2 to $n - 1$ has a locating-dominating set S of size $\lceil \frac{2(n-2)}{5} \rceil$ such that $2 \notin S$. We claim that S is a resolving set for $F_{1,n-1}$. On the one hand, as $n \geq 8$, then $|S| \geq 3$, so n is the only vertex at distance 1 from every vertex of S . On the other hand, 1 is the only vertex at distance 2 from every vertex of S , because of the choice of S .

Finally, every other vertex has different vector of distances to S because their neighborhoods in S are different, so that the 1's in the vectors of distances to S are located in different places. Consequently, S is a resolving set of $F_{1,n-1}$, and hence $\beta(F_{1,n-1}) \leq \lceil \frac{2(n-2)}{5} \rceil$.




Fig. 8 The set of black vertices is a metric basis of the fans of order 15, 16, 17, 18 and 19

We now prove that $\beta(F_{1,n-1}) \geq \lceil \frac{2(n-2)}{5} \rceil$. Let S be a metric basis of $F_{1,n-1}$. Since $d(i, n) = 1$ for $1 \leq i \leq n-1$, vertex n belongs to S only if it has the same coordinates as another vertex i with respect to the set $S \setminus \{n\}$. Then, $(S \setminus \{n\}) \cup \{i\}$ is also a metric basis of $F_{1,n-1}$. Hence, we may assume that $n \notin S$ and n is the only vertex with all metric coordinates 1, because $\beta(F_{1,n-1}) \geq 3$. Since $F_{1,n-1}$ has diameter 2, all metric coordinates of vertices not in S are 1 or 2. There is at most one vertex with all metric coordinates 2. If there is no vertex with all metric coordinates 2, then S is also a locating-dominating set of the path P of order $n-1$. Hence, $\beta(F_{1,n-1}) \geq \lambda(P_{n-1}) = \lceil \frac{2(n-1)}{5} \rceil \geq \lceil \frac{2(n-2)}{5} \rceil$. If there is one vertex i_0 with all metric coordinates 2, then S must be a locating-dominating set for $P - i_0$. If $i_0 \in \{1, n-1\}$, then $P - i_0$ is a path of order $n-2$ and $\beta(F_{1,n-1}) \geq \lambda(P_{n-2}) = \lceil \frac{2(n-2)}{5} \rceil$. If $i_0 \in \{3, \dots, n-3\}$, then $P - i_0$ has two connected components that are paths of order $r = i_0 - 1$ and $s = n - 1 - i_0$ respectively, with $r + s = n - 2$, and we have

$$\beta(F_{1,n-1}) \geq \lambda(P_r) + \lambda(P_s) = \left\lceil \frac{2r}{5} \right\rceil + \left\lceil \frac{2s}{5} \right\rceil \geq \left\lceil \frac{2(n-2)}{5} \right\rceil.$$

Finally, if $i_0 = 2$, then $1 \notin S$ and 1 would also be at distance 2 from every vertex in S , a contradiction. Therefore, $i_0 \neq 2$ and, analogously, $i_0 \neq n-2$. \square

It can be easily verified that if n is $5k$, $5k+1$ or $5k+2$ for some k , then $S = \{2 + 5r : 0 \leq r < \lfloor n/5 \rfloor\} \cup \{4 + 5r : 0 \leq r < \lfloor n/5 \rfloor\}$ is a metric basis of $F_{1,n-1}$, and if n is $5k+3$ or $5k+4$ for some k , then $S = \{2 + 5r : 0 \leq r < \lfloor n/5 \rfloor\} \cup \{4 + 5r : 0 \leq r < \lfloor n/5 \rfloor\} \cup \{n-1\}$ is a metric basis of $F_{1,n-1}$ (see Figure 8).

3.2 Upper bound

The main goal of this section is to show that every MOP graph $G = (V, E)$ of order n has a resolving set S of size $\lceil \frac{2n}{5} \rceil$ that can be built in linear time. For this purpose, we will begin with a certain set S of vertices of size $\lceil \frac{2n}{5} \rceil$. If S is a

resolving set, we are done. Otherwise, we will describe how S can be modified to obtain a resolving set of the same size. We will refer to the vertices belonging to S as black vertices, and vertices not in S as white vertices. Recall that the vertices of G are placed on a circle and labeled clockwise from 1 to n , so that all the edges are drawn inside the circle. A *run* will be a maximal set of consecutive vertices of the same color along the circle. We will denote by $[i, j]$ the set of vertices $\{i, i+1, \dots, j-1, j\}$, if $i < j$, and the set $\{i, i+1, \dots, n, 1, \dots, j-1, j\}$, if $i > j$.

We next prove some technical results.

Lemma 1 *Let G be a MOP graph of order n and $i, j \in [1, n]$. If $i, j, i-1$ and $j+1$ are four different vertices, then i and j are resolved by either $i-1$ or $j+1 \pmod{n}$.*

Proof Observe that G cannot contain at the same time the edges $(i, j+1)$ and $(j, i-1)$ because they cross, whenever $j+1 \not\equiv i-1 \pmod{n}$. Then, either $i-1$ or $j+1$ resolves i and j . See Figure 9. \square

We have seen in Section 3.1 that a resolving set of the fan can be obtained with alternating white runs of size 1 and 2 separated by black runs of size 1. Such a set is not a resolving set for a general MOP graph G , however, these kinds of sets will play an important role to construct a resolving set of G . This leads us to the following definition.




Fig. 9 Vertices i and j are resolved either by $i-1$ or $j+1$

We say that an interval $[i, j]$ is *(1,2)-alternating* if and only if all its white runs have size one or two, black runs have size one and there are no consecutive white runs of the same size.

Next lemma shows when two white vertices of a (1,2)-alternating interval are not resolved by any black vertex of the interval.

Lemma 2 *Let G be a MOP graph and let $[i_1, i_2]$ be a (1,2)-alternating interval such that the first and last vertices, i_1 and i_2 , are black. Let S' be the set of black vertices in the interval. The following properties hold.*

- (1) *Let $i \in [i_1, i_2]$ belong to a white run of size 1. Then, $r(i|S') = r(j|S')$ for some $j \in [i_1, i_2]$ if and only if j belongs to a white run of size 2 and one of the four cases (a), (b), (c) or (d) of Figure 10 holds.*
- (2) *Let $i, j \in [i_1, i_2]$ belong to white runs of size 2. Then, $r(i|S') = r(j|S')$ if and only if one of the four cases (e), (f), (g) or (h) of Figure 10 holds.*




Fig. 10 The eight cases in which two white vertices i and j , squared in the figure, are not resolved by black vertices

- (3) If $i \in [i_1, i_2]$ is a white vertex, then there is at most one white vertex $j \in [i_1, i_2]$ such that $r(i|S') = r(j|S')$.

Proof Let us prove item (1). By Lemma 1, two white vertices belonging to two different runs of size 1 are always resolved by vertices in S' . Suppose now that i belongs to a white run of size 1 and j belongs to a white run of size 2. If $r(i|S') = r(j|S')$, then, again by Lemma 1, either j is $i+2$ and j is connected to $i-1$, or j is $i-2$ and j is connected to $i+1$ (see Figure 10 top). Suppose first that $j = i+2$ and j is connected to $i-1$, so that $i-1$ and $i+4$ are black and $i+3$ is white. Since j is connected to $i-1$, we have that $2 \leq d(i, i+4) = d(j, i+4) \leq 2$. Hence, vertex $i-1$ is connected to $i+3$ and to $i+4$. Depending on which edge belongs to G , either $(i-1, i+1)$ or $(i, i+2)$, we have Cases (a)–(b) of Figure 10. Conversely, if Cases (a) or (b) hold, then for any black vertex $i' \in [i_1, i_2]$, the distances from i' to i and j are equal because the shortest path from i' to i or j goes through either $i-1$ or $i+4$, and in these cases the distances from i and j to $i-1$ (resp. to $i+4$) are the same. Thus, $r(i|S') = r(j|S')$. For the other case, that is, when $j = i-2$

and j is connected to $i + 1$, we have by symmetry that $r(i|S') = r(j|S')$ if and only if Cases (c)–(d) of Figure 10 hold. Therefore, we have proved (1).

We now prove item (2). By Lemma 1, it is clear that two white vertices i and $i + 1$ belonging to the same white run are resolved by either $i - 1$ or $i + 2$. Suppose now that i and j belong to different white runs of size 2 and $r(i|S') = r(j|S')$. In such a case, by Lemma 1, if $i - 1$ is a black vertex, then $j - 1$ is a black vertex, and there is an edge connecting i and $j - 1$ and another edge connecting j and $i - 1$. Also notice that, because of the assumption made in the hypothesis, $i + 2$ and $j + 2$ are black vertices. Since $2 \leq d(i, j + 2) = d(j, j + 2) \leq 2$, the only possibility for these two distances to be equal is that vertex $i - 1$ is connected to both $j + 1$ and $j + 2$. Analogously, taking into account that $2 \leq d(j, i + 2) = d(i, i + 2) \leq 2$, we derive that vertex $j - 1$ must be connected to $i + 1$ and $i + 2$. Depending on which edge belongs to G , either (i, j) or $(i - 1, j - 1)$, we have Cases (e)–(f) in Figure 10. Conversely, if Cases (e)–(f) hold, then one can easily check that $r(i|S') = r(j|S')$, as there are always shortest paths from i and j to any other black vertex passing through either $i - 1$ or $j - 1$. Cases (g)–(h) of Figure 10 appear by symmetry when $i + 1$ is black instead of $i - 1$. Therefore, (2) follows. Finally, item (3) is a direct consequence of items (1) and (2). \square

Before proving the main result of this section, we give some additional definitions. The vertex of degree two of Case (c) will be called a *special* vertex. Let $G = (V, E)$ be a graph. Given two subsets $S \subset V$ and $W \subset V$ we say that S *arranges* W if every pair of distinct vertices with at least one of them belonging to W is resolved by some vertex in S . If W consists of only one vertex u , we say that S *arranges* u . Observe that, by definition, if S_1 arranges W_1 and S_2 arranges W_2 , then $S_1 \cup S_2$ arranges $W_1 \cup W_2$. We next prove another technical lemma and the main result of this section, Theorem 4.

Lemma 3 *If $G = (V, E)$ is a MOP graph of order n , then there is a set $S_0 \subset V = [1, n]$ of black vertices such that $|S_0| = \lceil \frac{2n}{5} \rceil$, $\{2\}$ is a white run of size 1, the interval $[4, n]$ is (1, 2)-alternating and S_0 arranges the white run $\{2\}$.*

Proof Suppose that $n = 5k + t$, $t \in \{0, 1, 2, 3, 4\}$, for some $k \geq 1$. If $t \in \{0, 1, 2, 3\}$, we begin defining S_0 as the set of size $\lceil \frac{2n}{5} \rceil$ that consists of all the vertices of $[1, n]$ of the form $5j + 1$ and $5j + 3$. Vertices in S_0 are colored black and the rest, white. If $\{2\}$ is arranged by S_0 , we are done. Otherwise, by Lemmas 1 and 2, $\{2\}$ is the white run of size 1 of some of the subgraphs (a)–(d) of Figure 10. If we renumber all the vertices by rotating one place counterclockwise their labels, and update their colors according to the set S_0 , the new vertex with label 2 forms a white run of size 1 not belonging to any of the subgraphs (a)–(d) of Figure 10. Hence, S_0 arranges $\{2\}$. If $t = 4$, we define S_0 as the set formed by vertex 1 and all the vertices of $[1, n]$ of the form $5j + 3$ and $5j + 5$, with $j \geq 0$. Then, $\{n\}$, $\{2\}$ and $\{4\}$ are three consecutive white runs of size 1 separated by black vertices. By Lemma 1, S_0 arranges 2, and the interval $[4, n]$ is (1, 2)-alternating. \square

Theorem 4 *If $G = (V, E)$ is a MOP graph, then there exists a resolving set $S \subset V$ such that $|S| = \lceil \frac{2n}{5} \rceil$. Moreover, S can be computed in linear time.*

Proof The general procedure to obtain a resolving set for G is the following. We begin with the set $S := S_0$ defined in the proof of Lemma 3 that arranges the white run $\{2\}$ of size 1. If S_0 is a resolving set for G , we are done. Otherwise, we explore clockwise the white runs of S . Suppose that, after exploring the first h runs, S is a set consisting of $\lceil \frac{2n}{5} \rceil$ vertices which is a candidate to be a resolving set for G . If the next white run is not arranged by S because it belongs to some of the subgraphs shown in Figure 10, then we define a new set S' by removing some vertices in S and including new ones, such that $|S'| = |S|$ and all explored white runs are arranged by S' . Then, we update the set S , $S := S'$, and continue the exploration.

More precisely, suppose that, in a generic step of the exploration, the vertices in the interval $I = [1, i-1]$ have already been explored. Then, we denote by S the set of $\lceil \frac{2n}{5} \rceil$ black vertices of G and by W the set of white vertices of I , that is, $W = I \setminus S$.

We then prove that I and S satisfy the following invariant.

Invariant If $I = [1, i-1]$, $S \subseteq V$ and $W = I \setminus S$, then:

Property P1. $|S| = \lceil \frac{2n}{5} \rceil$ and S arranges W .

Property P2. Vertex i is white, vertices 1 and $i-1$ are black (that is, $\{1, i-1\} \subseteq S$ and $i \notin S$), and $[i, n]$ is an $(1, 2)$ -alternating interval.

Property P3. For every white vertex $w \in W \setminus \{i-2\}$ and every white special vertex $l \in V \setminus I$, there exists a black vertex $v \in S \cap (I \setminus \{i-1\})$ such that v resolves w and l .

Obviously, by Property P1, S will be a resolving set of size $\lceil \frac{2n}{5} \rceil$ for G after exploring all white runs. Properties P2 and P3 are technical facts that will be needed to proceed with the proof.

We begin verifying that Invariant holds for $S = S_0$ and $I = [1, 3]$. By Lemma 3, S_0 arranges 2, and interval $[4, n]$ is $(1, 2)$ -alternating. Thus, Property P2 obviously holds and Property P3 is true because in this case the set $W \setminus \{2\} = \emptyset$.

Assuming that Invariant is true for given sets $I = [1, i-1]$ and S , we next show that it holds for new sets I' and S' defined after exploring clockwise the next white run r not belonging to I . We will distinguish whether r is already arranged by S or not.

Suppose first that r is arranged by S . If r only consists of the white vertex i , then one can easily check that Invariant holds for $S' = S$ and $I' = I \cup [i, i+1]$. Indeed, as S arranges W and i , then S obviously arranges $W' = W \cup \{i\}$. Property P3 follows from the fact that a vertex in W satisfies Property P3, and a special vertex l cannot be connected to $i-1$, so $i-1$ resolves $i-2$ and l . Hence, I' satisfies Property P3. If r consists of two white vertices, $i, i+1$, consider $S' = S$ and $I' = I \cup [i, i+2]$. Sets S' and I' satisfy Property P1 of Invariant, because S arranges $W' = W \cup \{i, i+1\}$. To see that Property P3 is

true, notice that a special vertex $l \in V \setminus I'$ is not connected to $i - 1$. Hence, $i - 1$ resolves l and any of $i - 2$ and i .

Suppose now that r is not arranged by S . As S arranges W , the white vertices not resolved by S must belong to the interval $[i, n]$. Lemma 2 can be applied to the interval $[i - 1, i]$, if n is white, or to the interval $[i - 1, n]$, if n is black, since the interval $[i, n]$ is $(1, 2)$ -alternating by Property 2. Hence, r belongs to one of the subgraphs of Cases (a)-(h). Note that if r has size 1, then r consists of vertex i , and if r has size 2, then r consists of vertices i and $i + 1$.

In each one of these 8 cases, the general framework to construct new sets S' and I' satisfying Invariant is the following. The set I' is obtained by adding an interval $[i, i']$ to I , where i' is a black vertex and the vertices of the run r are in $[i, i']$. Then, we interchange the colors of some vertices from $[i - 1, i']$, so that the updated set S' of black vertices satisfies $|S'| = |S| = \lceil \frac{2n}{5} \rceil$, and Invariant holds for the new sets $I' = I \cup [i, i']$ and S' . To complete the validity of Property P1, it is needed to show that S' arranges W' after interchanging some colors in I' . This will be proved in two steps. First, we give a subset of S' that arranges the set of new white vertices, $X = W' \setminus W$. Secondly, we show that every pair of white vertices x and y , with $x \in W$ and $y \in V \setminus X$, that was resolved by a vertex from $S \setminus S'$, is now resolved by a vertex from $S' \setminus S$. Property P2 follows, because we have not changed the colors of the vertices from $V \setminus I'$. Finally, to prove that the sets S' and I' satisfy Property P3, it is enough to show that it holds for the white vertices in $(X \setminus \{i' - 1\}) \cup \{i - 2\}$. We next analyze the different cases.




Fig. 11 Squared vertices have the same coordinates. Case (a): interchanging the colors of i and $i - 1$, and $j = i + 2$ and $i + 1$. Case (b): interchanging the colors of $i + 1$ and $j = i + 2$

Case (a). The two vertices not resolved by S are i and j as shown in Figure 11 (a). In this case, we interchange the colors of vertices i and $i - 1$ and the colors of the vertices $j (= i + 2)$ and $i + 1$. We claim that Invariant holds for the new sets $I' = I \cup [i, i + 4]$ and $S' = (S \setminus \{i - 1, i + 1\}) \cup \{i, i + 2\}$.

Let us see that S' arranges $W' = W \cup \{i - 1, i + 1, i + 3\}$. On the one hand, the set $S^* = \{i, i + 2, i + 4\}$ arranges $\{i - 1, i + 1, i + 3\}$, because $r(i - 1 | S^*) = (1, 1, 1)$,

$r(i+1|S^*) = (1, 1, 2)$ and $r(i+3|S^*) = (2, 1, 1)$, respectively, and the only vertices adjacent to $i+2$ are precisely $i-1$, $i+1$ and $i+3$. On the other hand, the shortest path from $i' \in [i+5, i-2]$ to i , $i+1$ or $i+2$ necessarily goes through $i-1$, so $d(i', i) = d(i', i-1) + 1$ and $d(i', i+2) = d(i', i+1)$. This implies that, if $i-1$ (resp. $i+1$) resolves two vertices in $[i+5, i-2]$ then i (resp. $i+2$) also resolves them. In particular, S' arranges W' .

Let us see that Property P3 also holds. Take a special white vertex l in $V \setminus I'$. Property P3 clearly holds for the vertices of $W \setminus \{i-2\}$. Moreover, since $d(l, i+2) \geq 3$ and the distance from $i+2$ to any of $\{i-2, i-1, i+1\}$ is at most two, then we have that $i+2$ resolves l and any white vertex of $\{i-2, i-1, i+1\}$. Therefore, Property P3 is satisfied, and Invariant holds as claimed.

Case (b). The two vertices not resolved by S are i and j as shown in Figure 11 (b). In this case, we only need to interchange the colors of vertices $j (= i+2)$ and $i+1$. We claim that Invariant holds for the new sets $I' = [1, i+4]$ and $S' = (S \setminus \{i+1\}) \cup \{i+2\}$. Notice that $W' = W \cup \{i, i+1, i+3\}$.

The set $S^* = \{i+2, i+4\}$ arranges $\{i, i+1, i+3\}$ because $r(i|S^*) = (1, 2)$, $r(i+1|S^*) = (1, 3)$ and $r(i+3|S^*) = (1, 1)$, and the only white vertices adjacent to $i+2$ are precisely i , $i+1$ and $i+3$ (see Figure 11 (b)). Moreover, observe that for a vertex $i' \in [i+5, i-2]$, we have $d(i', i+2) = d(i', i+1) - 1$, so if $i+1$ resolves two vertices in $[i+5, i-2]$, then $i+2$ resolves them as well. As a consequence, S' arranges W .

Finally, to prove Property 3, note that the distance from $i+2$ to a special vertex $l \in V \setminus I'$ is at least 3. Thus, $i+2$ resolves the pairs formed by l and a vertex from $\{i-2, i, i+1\}$.




Fig. 12 Squared vertices have the same coordinates. Case (c): Interchanging the colors of $i+1$ and $i+2$, when $i+3$ is arranged by $(S \setminus \{i+2\}) \cup \{i+1\}$. Case (d): Interchanging the colors of $i+1$ and $i+2$

Case (c). In this case, the vertices not resolved by S are $i+1$ and $i+3$ (see Figure 12 (c)). We begin by interchanging the colors of the vertices $i+1$ and

$i + 2$, and distinguish two cases depending on whether $(S \setminus \{i + 2\}) \cup \{i + 1\}$ arranges $i + 3$ or not.

Suppose first that $i + 3$ is arranged by $(S \setminus \{i + 2\}) \cup \{i + 1\}$. Then, Invariant holds for the sets $S' = (S \setminus \{i + 2\}) \cup \{i + 1\}$ and $I' = I \cup [i, i + 4]$. Note that $W' = W \cup \{i, i + 2, i + 3\}$. Indeed, observe that $S^* = \{i - 1, i + 1\}$ arranges $\{i, i + 2\}$, because $r(i|S^*) = (1, 1)$, $r(i + 2|S^*) = (2, 1)$, and the only white vertices at distance 1 from $i + 1$ are i and $i + 2$. Besides, $d(i', i + 2) = d(i', i + 1)$ for every vertex $i' \in [i + 5, i - 2]$, implying that every pair of vertices belonging to $[i + 5, i - 2]$ that were resolved by $i + 2$ are now resolved by $i + 1$. In particular, S' arranges W . Therefore, S' arranges W' , and Property P1 holds. Since a special vertex l in $V \setminus I'$ is not connected to either $i - 1$ or $i + 1$, then l together with a vertex from $\{i - 2, i, i + 2\}$ are resolved by either $i - 1$ or $i + 1$. Then, Property P3 also holds.

Suppose now that $(S \setminus \{i + 2\}) \cup \{i + 1\}$ does not arrange $i + 3$. Let us see which vertex j has the same coordinates as $i + 3$ in relation to this set. Notice that by Property 3, since $i + 3$ is a special vertex, for any vertex $j \in W \setminus \{i - 2\}$ there is a vertex in $I \cap S$ resolving $i + 3$ and j . Besides, $i - 1$ resolves the pair $i + 3$ and $i - 2$, and $i + 1$ resolves $i + 3$ and any of i and $i + 2$. Hence, $j \notin [1, i + 2]$. By Property P2, a vertex $j \in [i + 5, n]$ is adjacent to a black vertex j' , but j' is not adjacent to $i + 3$ unless $j' = i + 4$ and $j = i + 5$. Then, $i + 6$ is white and $i + 7$ is black (see Figure 13). Since $2 \leq d(i + 3, i + 7)$ and $d(i + 5, i + 7) \leq 2$, we have that both distances are equal only when the edges $(i + 4, i + 7)$ and $(i + 4, i + 6)$ belong to G .

If this situation happens, then $i + 3$ and $i + 5$ have the same coordinates in relation to $(S \setminus \{i + 2\}) \cup \{i + 1\}$. We remark that Property P3 is important at this time to ensure that $i + 5$ is the only vertex with the same coordinates as $i + 3$. Otherwise, if Property P3 does not hold, then a vertex x in W could have the same coordinates as $i + 3$, because $i + 2$ could be the only vertex in S to resolve $i + 3$ and x .




Fig. 13 Case (c). Squared vertices have the same coordinates. When $i + 3$ is not arranged by $(S \setminus \{i + 2\}) \cup \{i + 1\}$, the colors of $i + 3$ and $i + 4$ are interchanged

We interchange the colors of vertices $i + 3$ and $i + 4$, as shown in Figure 13, and set $I' = [1, i + 7]$ and $S' = (S \setminus \{i + 2, i + 4\}) \cup \{i + 1, i + 3\}$. Thus,

$W' = W \cup \{i, i+2, i+4, i+5, i+6\}$. The argument to prove that Invariant holds for these new sets is similar to the previous ones, but a bit more elaborated.

Let us show that S' arranges W' . On the one hand, if $S^* = \{i-1, i+1, i+3\} \subseteq S'$, then $r(i|S^*) = (1, 1, 2)$, $r(i+2|S^*) = (2, 1, 1)$ and $r(i+4|S^*) = (1, 1, 1)$. In addition, the only vertices at distance 1 from $i+1$ are $i, i+2$ and $i+4$. Hence, S' arranges $\{i, i+2, i+4\}$. On the other hand, for every vertex $i' \in [i+5, i-2]$, we have $d(i', i+2) = d(i', i+1)$ and $d(i', i+3) = d(i', i+4)+1$. This implies that, any pair of vertices from $[i+5, i-2]$ resolved by $i+2$ or $i+4$ is also resolved by $i+1$ or $i+3$. Therefore, since $W \subseteq [i+5, i-2]$ and S arranges W , we derive that S' arranges W . It only remains to prove that S' arranges $\{i+5, i+6\}$. Notice that $i+7$ resolves the pair $i+5$ and $i+6$. Moreover, by Property P2, a white vertex j in $V \setminus I'$ is adjacent to a black vertex j' . Since $i+5$ and $i+6$ cannot be connected to j' , then j' resolves j and any vertex of $i+5$ and $i+6$. Note that if $j = i+8$, then $i+9$ is such a vertex j' . Finally, as S' arranges W and $\{i, i+2, i+4\}$, a vertex from $\{i+5, i+6\}$ and a vertex from $[1, i+4]$ are resolved by some vertex of S' . Hence, Property P1 is satisfied.

To show that Property P3 holds, we only need to prove this property for the vertices $i-2, i, i+2, i+4$ and $i+5$. For a special vertex l in $V \setminus I'$, its distance to $i+3$ is at least 3. Since the distance from $i+3$ to $i, i+2, i+4$, or $i+5$ is at most 2, vertex $i+3$ resolves l and any of these four vertices. The pair l and $i-2$ is resolved by $i-1$, because l is not adjacent to $i-1$.

Case (d). In this case, the vertices not resolved by S are $i+1$ and $i+3$ in the subgraph shown in Figure 12 (d). This case is symmetric to Case (b). Following the same kind of arguments used in that case, one can easily prove that Invariant holds for the sets $I' = [1, i+4]$ and $S' = (S \setminus \{i+2\}) \cup \{i+1\}$, defined after interchanging the colors of vertices $i+1$ and $i+2$ (see Figure 12 (d)).




Fig. 14 Cases (e) and (f): squared vertices have the same coordinates. Interchanging the colors of $i-1$ and i

Case (e). In this case, the vertices not resolved by S are i and j in the subgraph shown in Figure 14 (e). We interchange the colors of vertices $i - 1$ and i and we define $I' = I \cup [i, i + 2]$ and $S' = (S \setminus \{i - 1\}) \cup \{i\}$. Thus, $W' = W \cup \{i - 1, i + 1\}$ (see Figure 14 (e)).

Let us see first that S' arranges W' . On the one hand, the set $S^* = \{i, i + 2\}$ arranges $\{i - 1, i + 1\}$. Indeed, $r(i - 1|S^*) = (1, 3)$, $r(i + 1|S^*) = (1, 1)$, and the only white vertices belonging to V at distance 1 from i are $i - 1, i + 1$ and j , but $r(j|S^*) = (1, 2)$. On the other hand, $d(i', i) = d(i', i - 1) + 1$ for every vertex $i' \in [j + 1, i - 2]$. Hence, since $W \subseteq [j + 3, i - 2]$, every pair of vertices with at least one of them belonging to W and the other to $[j + 3, i - 2]$ that was resolved by $i - 1 \in S$ is now resolved by $i \in S'$. It only remains to prove that every pair formed by a vertex i' from W and a white vertex $j' \in [i + 3, j]$ is resolved by some vertex of S' . This is true because, by Property P2, j' is adjacent to a black vertex in $S' \cap [i + 3, j]$ that is not adjacent to i' . Hence, Property P1 holds. To prove property P3, it suffices to check that it holds for the vertices $i - 1$ and $i - 2$. If $l \in V \setminus I'$ is a special vertex different from $j - 2$, then $d(l, i) \geq 3$. Hence, the pairs formed by l and a vertex from $\{i - 1, i - 2\}$ are resolved by i , whenever $l \neq j - 2$. Suppose that $l = j - 2$ is a special vertex. If we take a black vertex $j'' \neq i - 1$ in I , then $d(j - 2, j'') = 3 + d(i - 1, j'')$ and $d(i - 2, j'') \leq d(i - 2, i - 1) + d(i - 1, j'') = 1 + d(i - 1, j'')$. Hence, $j'' \in I'$ resolves $j - 2$ and any of $i - 1$ and $i - 2$.

Case (f). In this case, the vertices not resolved by S are i and j in the subgraph shown in Figure 14 (f). This case is very similar to the previous one. By interchanging the colors of vertices $i - 1$ and i (see Figure 14 (f)), the proof that sets $I' = [1, i + 2]$, and $S' = (S \setminus \{i - 1\}) \cup \{i\}$ satisfy Invariant is essentially the same as the proof done in Case (e), with small differences due to the fact that the edge $(i - 1, j - 1)$ now belongs to G instead of edge (i, j) .

Case (g). In this case, the vertices not resolved by S are $i + 1$ and $j + 1$ in the subgraphs shown in Figure 15. We begin by interchanging the colors of vertices $i + 1$ and $i + 2$. We distinguish two cases depending on whether $(S \setminus \{i + 2\}) \cup \{i + 1\}$ arranges $i + 3$ or not.

Suppose first that $(S \setminus \{i + 2\}) \cup \{i + 1\}$ arranges $i + 3$ (see Figure 15, top). We claim that $S' = (S \setminus \{i + 2\}) \cup \{i + 1\}$ and $I' = I \cup [i, i + 4]$ satisfy Invariant. Note that $W' = W \cup \{i, i + 2, i + 3\}$.

On the one hand, the set $S^* = \{i - 1, i + 1\}$ arranges $\{i, i + 2, j + 1\}$. Indeed, $r(i|S^*) = (1, 1)$, $r(i + 2|S^*) = (3, 1)$, $r(j + 1|S^*) = (2, 1)$ and the only white vertices adjacent to $i + 1$ are $i, i + 2$ and $j + 1$. We include here vertex $j + 1$ to ensure that $j + 1$ and a vertex in W are resolved. On the other hand, a white vertex $i' \in [i + 4, j]$ is adjacent by Property P2 to a black vertex j' in this interval. Thus, j' resolves any pair formed by i' together with every white vertex of $W \subseteq [j + 3, i - 2]$ because the vertices of this last interval are not adjacent to j' . In addition, since $d(i', i + 1) = d(i', i + 2) - 1$ for every vertex $i' \in [j + 3, i - 1]$ and $W \subseteq [j + 3, i - 1]$, every pair of vertices in this interval,




Fig. 15 Case (g). Squared vertices have the same coordinates. Top: Interchanging the colors of $i + 1$ and $i + 2$. Middle and bottom: Interchanging the colors of $i + 1$ and $i + 2$ and the colors of $i + 4$ and $i + 5$

with one of them in W , that was resolved by $i + 2$ is now resolved by $i + 1$. Hence, S' arranges W . Besides, since a special white vertex l is not connected to either $i + 1$ or $i - 1$, and vertices $i - 2$, i and $i + 2$ are adjacent to at least one of them, Property P3 holds.

Suppose now that $(S \setminus \{i + 2\}) \cup \{i + 1\}$ does not arrange $i + 3$. Let us see which white vertex j' has the same coordinates as $i + 3$ with respect to this set. A vertex in the interval $[j, i + 1]$ is not adjacent $i + 4$, thus $j' \in [i + 2, j - 2]$. Moreover, $j' \neq i + 2$, because $i + 3$ is not adjacent to $i + 1$, and consequently, $j' \in [i + 5, j - 2]$. Observe now that if $d(i + 3, i + 1) = d(j', i + 1) = 2$ then $d(i + 3, i + 2) = d(j', i + 2) = 1$. Taking vertex $i + 2$ as a black vertex, we can apply Lemma 2 to the $(1, 2)$ -alternating interval $[i + 2, n]$ (or $[i + 2, 1]$), giving rise to the only two possibilities shown in Figure 15, middle and bottom, for a vertex $j' = i + 5$ to have the same coordinates as $i + 3$.

Consider the case shown in Figure 15, middle: vertex $i + 2$ connected to vertices $i + 3, i + 4, i + 5, i + 6$ and $i + 7$. In addition to the changes of color of $i + 1$ and $i + 2$, we interchange the colors of vertices $i + 4$ and $i + 5$. We

claim that the sets $S' = (S \setminus \{i+2, i+4\}) \cup \{i+1, i+5\}$ and $I' = I \cup [i, i+7]$ satisfy Invariant. Note that $W' = W \cup \{i, i+2, i+3, i+4, i+6\}$. The set $S^* = \{i-1, i+1, i+5, i+7\}$ arranges $\{i, i+2, i+4, i+6, j+1\}$, since $\{i, i+2, i+4, i+6, j+1\}$ are the only white vertices adjacent to $i+1$ or $i+5$, and $r(i|S^*) = (1, 1, 3, 3)$, $r(i+2|S^*) = (3, 1, 1, 1)$, $r(i+4|S^*) = (4, 2, 1, 2)$, $r(i+6|S^*) = (4, 2, 1, 1)$ and $r(j+1|S^*) = (2, 1, 2, 2)$. On the other hand, $i+3$ has no black neighbor. Hence, any other white vertex $i' \in V \setminus W'$ has at least a black neighbor that resolves $i+3$ and i' . If $i' \in W \subseteq [j+3, i-2]$, then $i+5$ resolves i' and $i+3$, because $d(i', i+5) \geq 4$, but $d(i+3, i+5) = 2$. Thus, S' arranges $i+3$. Finally, as S arranges W and $W \subseteq [j+3, i-2]$, then S' also arranges W taking into account that (1) $j+1$ is already resolved from a vertex in W , (2) for every vertex $i' \in [j+3, i-2]$, we have $d(i', i+5) = d(i', i+4)$ and $d(i', i+1) = d(i', i+2) - 1$ and that (3) a black vertex adjacent to a white vertex of $V \setminus I'$ in the interval $[i+8, j]$ cannot be connected to a vertex in W . Therefore, S' arranges W' and Property P1 holds.

To show that Property P3 holds we only need to resolve the pairs formed by a special vertex $l \in V \setminus I'$ and one of the vertices from $\{i-2, i, i+2, i+3, i+4\}$ using a vertex from $S' \cap [1, i+5]$. Vertex $i-1$ resolves l and any of $\{i-2, i\}$ because l is not adjacent to $i-1$, and $i+5$ resolves l and any of $\{i+2, i+3, i+4\}$ because $d(l, i+5) \geq 3$. Therefore, Property P3 also holds, and Invariant is satisfied, as claimed.

For the last case, the one shown in Figure 15, bottom, the analysis is very similar to the previous one. Following the same steps as described in the two previous paragraphs, one can prove that $S' = (S \setminus \{i+2, i+4\}) \cup \{i+1, i+5\}$ and $I' = I \cup [i, i+7]$ satisfy Invariant. The set W' is $W \cup \{i, i+2, i+3, i+4, i+6\}$. In this case, it can be shown that the set $S^* = \{i-1, i+1, i+5, i+7\}$ arranges $\{i, i+2, i+3, i+4, i+6, j+1\}$. Moreover, for every white vertex $i' \in [j+3, i-2]$, we have $d(i', i+1) = d(i', i+2) - 1$ and $d(i', i+5) = d(i', i+4) - 1$. Thus, every pair of white vertices from $[j+3, i-2]$ that was resolved by $i+2$ or $i+4$ is resolved now by $i+1$ or $i+5$. For every white vertex $i' \in [i+8, j]$, the black vertex adjacent to i' is not adjacent to a vertex in $W \subseteq [j+3, i-2]$. Hence, S' arranges W .

Finally, to show that Property P3 holds we only need to resolve the pairs formed by a special vertex $l \in V \setminus I'$ and one of the vertices from $\{i-2, i, i+2, i+3, i+4\}$ using a vertex from $S' \cap [1, i+5]$. All the vertices in $\{i-2, i, i+2, i+3, i+4\}$ are adjacent to either $i-1$ or $i+5$, but a special vertex $l \in V \setminus I'$ is not adjacent to either $i-1$ or $i+5$, so $i-1$ or $i+5$ resolves l and any of these five vertices. From this, Invariant holds as claimed.

Case (h). In this case, the vertices not resolved by S are $i+1$ and $j+1$ in the subgraphs shown in Figure 16. The analysis of Case (h) follows the same steps as Case (g), although there are small changes due to the fact that now the edge $(i+2, j+2)$ belongs to G instead of the edge $(i+1, j+1)$.

If $(S \setminus \{i+2\}) \cup \{i+1\}$ arranges $i+3$, it can be checked that $S' = (S \setminus \{i+2\}) \cup \{i+1\}$ and $I' = [1, i+4]$ satisfy Invariant (see Figure 16, top).




Fig. 16 Case (h). Squared vertices have the same coordinates. Top: Interchanging the colors of $i+1$ and $i+2$. Middle and bottom: Interchanging the colors of $i+1$ and $i+2$ and the colors of $i+4$ and $i+5$

If $(S \setminus \{i+2\}) \cup \{i+1\}$ does not arrange $i+3$, arguing exactly as in Case (g), we have that the vertex with the same coordinates as $i+3$ is $j' = i+5$, and one of the cases shown in the middle and bottom of Figure 16 holds. It can be checked in both cases that the sets $S' = (S \setminus \{i+2, i+4\}) \cup \{i+1, i+5\}$ and $I' = I \cup [i, i+7]$ satisfy Invariant (see Figure 16, middle and bottom).

To finish the proof of the theorem, let us see that S can be computed in linear time. Building $S = S_0$ obviously requires linear time. Besides, for every run r , we have to check if subgraphs (a)–(h) appear in G and, if it is the case, to update S accordingly. All of this can be done in constant time. Therefore, S can be computed in linear time. \square

4 Conclusions

In this paper, we have studied the metric dimension problem for maximal outerplanar graphs, and we have shown that $2 \leq \beta(G) \leq \lceil \frac{2n}{5} \rceil$ for any maximal outerplanar graph G . In relation to the lower bound, we have characterized all maximal outerplanar graphs with metric dimension two, based on embedding such graphs into the strong product of two paths. A first question is whether this technique can be applied to characterize graphs with metric dimension two in other families of graphs, as 2-trees or near-triangulations.

With respect to the upper bound, we have provided a linear algorithm to build a resolving set of size $\lceil \frac{2n}{5} \rceil$ for any maximal outerplanar graph. A second question is whether similar techniques as those described in the algorithm can be used to find efficiently resolving sets for other families of graphs, as Hamiltonian outerplanar graphs or near-triangulations. For near-triangulations, the conjecture is that there always exists a resolving set of size $\lceil \frac{2n}{5} \rceil$ for any near-triangulation.

Acknowledgements A. García, M. Mora and J. Tejel are supported by H2020-MSCA-RISE project 734922 - CONNECT; M. Claverol, A. García, G. Hernández, C. Hernando, M. Mora and J. Tejel are supported by project MTM2015-63791-R (MINECO/FEDER); M. Claverol is supported by project Gen. Cat. DGR 2017SGR1640; C. Hernando, M. Maureso and M. Mora are supported by project Gen. Cat. DGR 2017SGR1336; A. García and J. Tejel are supported by project Gobierno de Aragón E41-17R (FEDER).

References

1. Beerliova, Z., Eberhard, F., Erlebach T., Hall, A., Hoffman, M., Mihalák, M., Ram, L.S., Network discovery and verification, *IEEE J. Sel. Areas Commun.*, 24, 2168–2181 (2006)
2. Behtoei, A., Davoodi, A., Jannesari, M., Omoomi, B., A characterization of some graphs with metric dimension two, *Discrete Mathematics, Algorithms and Applications*, 9, 175027 (15 pages) (2017)
3. Cáceres, J., Hernando, C., Mora, M., Pelayo I.M., Puertas, M.L., On the metric dimension of infinite graphs, *Discrete Appl. Math.*, 160(18), 2618–2626 (2012)
4. Cáceres, J. , Hernando, C. , Mora, M., Pelayo, I.M., Puertas, M.L., Seara C. , Wood, D.R., On the Metric Dimension of Cartesian Products of Graphs, *SIAM Journal on Discrete Mathematics*, 21, 423–441 (2007)
5. Chartrand, G., Eroh, L., Johnson, M.A., Oellermann, O.R., Resolvability in graphs and the metric dimension of a graph, *Discrete Appl. Math.*, 105, 99–113 (2000)
6. Chvátal, V., Mastermind, *Combinatorica*, 3, 325–329 (1983)
7. Díaz, J., Potttonen, O. , Serna, M., van Leeuwen, E.J., Complexity of metric dimension on planar graphs, *Journal of Computer and System Sciences* 83, 132–158 (2017)
8. Dudenko, M., Oliynyk, B., On unicyclic graphs of metric dimension 2, *Algebra and Discrete Mathematics*, 23, 216–222 (2017)
9. Epstein, L., Levin, A., Woeginger, G.J., The (weighted) metric dimension of graphs: hard and easy cases, *Algorithmica*, 72, 1130–1171 (2015)
10. Fernau, H., Heggernes, P., van't Hof, P., Meister, D., Saei, R., Computing the metric dimension for chain graphs, *Information Processing Letters*, 115, 671–676 (2015)
11. Foucaud, F., Mertzios, G.B., Naserasr, R., Parreau, A., Valicov, P., Identification, location-domination and metric dimension on interval and permutation graphs. II. Complexity and algorithms, *Algorithmica*, 78, 914–944 (2017)

12. Foucaud, F., Mertzios, G.B., Naserasr, R., Parreau, A., Valicov, P., Identification, location-domination and metric dimension on interval and permutation graphs. I. Bounds, *Theoretical Computer Science*, 668, 43–58 (2017)
13. Garijo, D., González, A., Márquez, A., The difference between the metric dimension and the determining number of a graph, *Applied Mathematics and Computation*, 249, 487-501 (2014)
14. González, A., Hernando, C., Mora, M., Metric-locating-dominating sets of graphs for constructing related subsets of vertices, *Applied Mathematics and Computation*, 332, 449-456 (2018)
15. Grigoriou, C., Manuel, P., Miller, M., Rajan, B., Stephen, S., On the metric dimension of circulant and Harary graphs, *Applied Mathematics and Computation*, 248, 47-54 (2014)
16. Harary, F., Melter, R.A., On the metric dimension of a graph, *Ars Combinatoria*, 2, 191–195, (1976)
17. Hernando, C., Mora, M., Pelayo, I.M., Seara C., Wood, D.R., Extremal graph theory for metric dimension and diameter, *Electron. J. Combin.*, 17, R30 (2010)
18. Khuller, S., Raghavachari B., Rosenfeld, A., Landmarks in graphs, *Discrete Appl. Math.*, 70, 217–229 (1996)
19. Muhammad, H., Siddiqui A., Imran, M., Computing the metric dimension of wheel related graphs, *Applied Mathematics and Computation*, 242, 624-632 (2014)
20. Quines C.J., Sun, M., Bounds on metric dimension for families of planar graphs, [arXiv:1704.04066v1](https://arxiv.org/abs/1704.04066v1) (2017)
21. Slater, P.J., Leaves of trees, *Congressus Numerantium*, 14, 549–559 (1975)
22. Slater, P.J., Dominating and reference sets in a graph, *J. Math. Phys. Sci.*, 22, 445–455 (1988)
23. Sudhakara, G., Hemanth Kumar, A.R., Graphs with metric dimension two-a characterization, *International Journal of Mathematical and Computational Sciences*, 3, 1128–1133 (2009)
24. Vatandoost, E., Behtoei, A., Golkhandy Pour, Y., Cayley graphs with metric dimension two - A characterization, [arXiv: 1609.06565v1](https://arxiv.org/abs/1609.06565v1) (2016)
25. Yero, I. G., Estrada-Moreno, A., Rodríguez-Velázquez, J.A., Computing the k-metric dimension of graphs, *Applied Mathematics and Computation*, 300, 60-69 (2017)