

POINTWISE MONOTONICITY OF HEAT KERNELS

DIEGO ALONSO-ORÁN, FERNANDO CHAMIZO, ÁNGEL DAVID MARTÍNEZ,
AND ALBERT MAS

ABSTRACT. In this paper we present an elementary proof of a pointwise radial monotonicity property of heat kernels that is shared by the Euclidean spaces, spheres and hyperbolic spaces. The main result was discovered by Cheeger and Yau in 1981 and rediscovered in special cases during the last few years. It deals with the monotonicity of the heat kernel from special points on revolution hypersurfaces. Our proof hinges on a non straightforward but elementary application of the parabolic maximum principle. As a consequence of the monotonicity property, we derive new inequalities about classical special functions.

1. INTRODUCTION

The heat equation is one of the quintessential mathematical models for physical phenomena. Over the years, several properties of this equation had been studied from different points of view, including for instance: probabilistic, geometric and physical. In this paper we deal with the fundamental solution of the heat equation, namely, the heat kernel. In particular, we will focus on certain generalizations, due to Cheeger and Yau, of the following intuitive result

Theorem 1.1 ([4, 11, 13, 30]). *Let M be Euclidean space \mathbb{R}^n , the sphere \mathbb{S}^n or the hyperbolic space \mathbb{H}^n . Then, for any fixed $x \in M$ and time $t \in (0, \infty)$, the heat kernel $G(x, y, t)$ is a strictly decreasing function of the geodesic distance $d(x, y)$.*

First, note that by means of Fourier analysis one can provide an explicit expression of the heat kernel in the Euclidean space \mathbb{R}^n , namely

$$G(x, y, t) = \frac{1}{(4\pi t)^{n/2}} \exp\left(-\frac{|x - y|^2}{4t}\right),$$

where the assertion of Theorem 1.1 follows trivially. However, explicit expressions like the above are rare. The hyperbolic spaces are exceptions, e.g. for the hyperbolic plane one gets the following representation

$$G(x, y, t) = \frac{\sqrt{2}e^{-t/4}}{(4\pi t)^{3/2}} \int_{d(x,y)}^{\infty} \frac{\beta e^{-\beta^2/4t}}{\sqrt{\cosh \beta - \cosh(d(x, y))}} d\beta.$$

In general, distinguishing odd and even dimensional cases, rather involved formulas are available (cf. [11]). The case of the sphere does not share this good fortune, but one can use an identity¹ expressing the derivative of the kernel in terms of the heat kernel in

¹This formula also works in a broader generality for 3-dimensional space forms \mathbb{M}_k with sectional curvature k .

smaller dimensions to prove the monotonicity by an induction argument (the base case can be found in [11, §6.3]). Recently, Nowak, Sjögren and Szarek [30] provided sharp estimates for the heat kernel on the sphere \mathbb{S}^n that imply Theorem 1.1 in that specific case.

Here we propose a neat, direct and elementary proof of Theorem 1.1 which is built on a delicate application of the parabolic maximum principle. Interestingly enough, the same arguments also apply to more general situations described below, of which Theorem 1.1 is a beautiful particular case.

Theorem 1.2. *Let $M \subseteq \mathbb{R}^n$ be a smooth, compact and connected hypersurface of revolution around the x_n axis without boundary. If x is a point of intersection of M and the x_n axis, then the associated heat kernel $G(x, y, t)$ strictly decreases as a function of the geodesic distance $d(x, y)$ for any fixed $t > 0$.*

Remark 1.3. *The same proof covers the noncompact situation even in an intrinsic geometric setting beyond hypersurfaces of \mathbb{R}^n . In connection with this, recall that a celebrated theorem by Hilbert states that a complete regular surface of constant negative curvature, like \mathbb{H}^2 , cannot be isometrically immersed in \mathbb{R}^3 .*

Remark 1.4. *When we compare our approach to that of Cheeger and Yau [13], we strongly believe that ours is more PDE friendly and falls within a certain canon. Both proofs rely on the maximum principle, nevertheless. The proof in [13] is more technically demanding, namely it employs 1-forms and the small time asymptotics of the heat kernel. Furthermore, it relies on a certain contradiction (obtained using an integral quantity that is proved to be zero) while ours is more direct.*

Theorem 1.2 can be proven even in more general settings. Let us introduce some definitions before stating this. In a complete Riemannian manifold M of dimension n , a point p is called a pole if its cut locus is empty (see [12] for the definition of the cut locus). A manifold is said to be spherically symmetric (around p) if in terms of the geodesic polar coordinates $(\rho, \sigma) \in (0, \infty) \times \mathbb{S}^{n-1}$, its Riemannian metric ds^2 of M has the form $ds^2 = d\rho^2 + A^2(\rho)d\sigma^2$. In other words, the rotations around the origin in $T_p(M)$ become isometries of M under the exponential map². Finally, we need some control on the volume of the surface of balls in order to apply the maximum principle.

Theorem 1.5. *Let M be a complete spherically symmetric manifold with bounded curvature. Let S be the volume of the boundary of the ball of radius ρ in M centered on a pole p and suppose that $\frac{\partial^2}{\partial \rho^2} \log(S(\rho))$ is bounded from above. Then the heat kernel $G(p, y, t)$ based on the pole p , satisfies $\int_M G(p, y, t) dy = 1$ and it is a strictly decreasing function of the geodesic distance $d(p, y)$ for any fixed $t > 0$.*

Remark 1.6. *We have recently been aware of an extension of the results in [30] by the same authors where they provide sharp bounds for the heat kernels on all compact rank-one symmetric spaces, cf. [31].*

Our last result, also known to Cheeger and Yau, tackles the Dirichlet and Neumann heat kernels, G_D and G_N respectively, of a smooth hypersurface of revolution $M \subseteq \mathbb{R}^n$

²This kind of manifolds are called “model manifolds with $R_0 = \infty$ ” in [23] although we shall not employ this terminology here.

with boundary. Recall that G_D is the fundamental solution for the Dirichlet heat operator $\partial_t - \Delta_M$ (where we denoted by $-\Delta_M$ the positive Laplace-Beltrami operator on M) with the Dirichlet boundary condition, that is, for a fixed $x \in M$, $G_D(x, y, t)$ is the function in (y, t) satisfying

$$\begin{cases} (\partial_t - \Delta_M)G_D(x, y, t) = 0, & (y, t) \in M \times (0, +\infty), \\ G_D(x, y, t) = 0, & (y, t) \in \partial M \times (0, +\infty), \\ G_D(x, y, 0) = \delta_x(y), & y \in M, \end{cases}$$

where δ_x is the Dirac delta based on x . Similarly, for G_N with the Neumann boundary condition, that is, for a fixed $x \in M$, $G_N(x, y, t)$ is the function in (y, t) satisfying

$$\begin{cases} (\partial_t - \Delta_M)G_N(x, y, t) = 0, & (y, t) \in M \times (0, +\infty), \\ \frac{\partial}{\partial n} G_N(x, y, t) = 0, & (y, t) \in \partial M \times (0, +\infty), \\ G_N(x, y, 0) = \delta_x(y), & y \in M. \end{cases}$$

Theorem 1.7. *Let $M \subseteq \mathbb{R}^n$ be smooth and connected hypersurface of revolution around the x_n axis with boundary $\partial M \neq \emptyset$. If x is a point of intersection of the relative interior of M and the x_n axis, then the associated heat kernel with Dirichlet boundary condition (resp. Neumann boundary condition) $G_D(x, y, t)$ (resp. $G_N(x, y, t)$) strictly decreases as a function of the geodesic distance $d(x, y)$ for any fixed time $t > 0$.*

Remark 1.8. *An analogous result in the case of the Dirichlet heat kernel on a geodesic ball inside a n -dimensional simply connected space form of constant sectional curvature was already known (cf. [11, §8.3]). Theorem 1.7 has also been settled (cf. [13, §2]) for model manifolds. The interest of these results originates from the possibility of comparing the fundamental solutions to the heat equation on general manifolds with heat kernels of model manifolds.*

Plan of the paper: The paper is organized along the following lines: in the next section we present the proofs of the obtained results where we demonstrate Theorem 1.2, sketch the proof of Theorem 1.5 and infer as a consequence Theorem 1.1. Section 3 is devoted to show some consequences regarding inequalities for special functions and some observations of pointwise properties of the fractional Laplace-Beltrami operator. The article ends raising a natural question that bonds the decreasing properties of the heat kernel from a point with its cut locus.

2. PROOF OF THE MAIN RESULTS

Note that Theorem 1.1 is a direct corollary of Theorem 1.5. The main ideas in the proof of the latter are in that of Theorem 1.2. The whole argument relies on an application of the parabolic maximum principle.

Proof of Theorem 1.2. Thanks to the symmetry around the x_n axis, the heat kernel $G(x, y, t)$ is a function Γ only depending on the geodesic distance from x to y , $\rho = d(x, y)$, and the time variable t , that is, $G(x, y, t) =: \Gamma(\rho, t)$. This symmetry also justifies expressing the homogeneous heat equation on M for smooth³ functions $u : (0, L) \times (0, +\infty) \rightarrow \mathbb{R}$

³Recall that, for example, any function $u : (0, L) \times (0, +\infty) \rightarrow \mathbb{R}$ whose up to first order weak derivatives are locally square integrable and which solves (2.1) in the weak sense is indeed of class C^∞

which depend only on the geodesic distance and the time variable as [23, §3.2]

$$(2.1) \quad \partial_t u(\rho, t) - \partial_{\rho\rho} u(\rho, t) - \partial_\rho(\log(S(\rho))) \partial_\rho u(\rho, t) = 0$$

for all $\rho \in (0, L)$, $t > 0$, where L denotes the geodesic distance from x to its antipodal point, and $S(\rho)$ denotes the volume of the boundary of the ball of radius ρ in M centered at x . Differentiating (2.1) with respect to ρ we obtain

$$(2.2) \quad \partial_t(\partial_\rho u) - \partial_{\rho\rho}(\partial_\rho u) - \partial_{\rho\rho}(\log(S(\rho))) (\partial_\rho u) - \partial_\rho(\log(S(\rho))) \partial_\rho(\partial_\rho u) = 0.$$

Near x we see that $\partial_{\rho\rho}(\log(S(\rho)))$ behaves like in the euclidean case. This provides the approximation $-(n-2)\rho^{-2}$ for ρ near zero. By continuity and compactness we conclude that $\partial_{\rho\rho}(\log(S(\rho)))$ is bounded from above in $(0, L)$ if M is compact. With this at hand, we define

$$(2.3) \quad \begin{aligned} \lambda &:= \sup_{\rho \in (0, L)} \partial_{\rho\rho}(\log(S(\rho))), \\ c(\rho) &:= \lambda - \partial_{\rho\rho}(\log(S(\rho))), \\ v(\rho, t) &:= e^{-\lambda t}(\partial_\rho u)(\rho, t). \end{aligned}$$

By the previous considerations, we have that $\lambda \in \mathbb{R}$ and $c \geq 0$. Now, since $\partial_t(\partial_\rho u) = e^{\lambda t}(\lambda v + \partial_t v)$, from (2.2) we get

$$(2.4) \quad \begin{aligned} 0 &= \lambda v + \partial_t v - \partial_{\rho\rho} v - \partial_{\rho\rho}(\log(S(\rho))) v - \partial_\rho(\log(S(\rho))) \partial_\rho v \\ &= \partial_t v - \partial_{\rho\rho} v + b \partial_\rho v + cv = (\partial_t + \mathcal{L})v, \end{aligned}$$

where we have set $b(\rho) := -\partial_\rho(\log(S(\rho)))$ and $\mathcal{L} := -\partial_{\rho\rho} + b \partial_\rho + c$.

Our first target is to show that $\Gamma(\cdot, t)$ is a nonincreasing function on $[0, L]$ for all $t > 0$. In order to do so, let us regularize the heat kernel G employing a family of functions $\{\chi_\varepsilon(x, \cdot)\}_{\varepsilon > 0}$ that satisfy the following properties: they are smooth, radially symmetric (i.e. $\chi_\varepsilon(x, y) =: g_\varepsilon(\rho)$), and decreasing from x , their integral over M equals one, and they concentrate around the fixed point $x \in M$ as ε tends to zero. Notice that this family is, in particular, an approximation of the Dirac delta centered at x . Let us now introduce the aforementioned regularization of the heat kernel

$$G_\varepsilon(x, y, t) := \int_M \chi_\varepsilon(x, z) G(z, y, t) d\sigma(z),$$

where $d\sigma$ denotes the volume form on M . By construction, G_ε satisfies the heat equation in the variables (y, t) and the initial condition $\chi_\varepsilon(x, \cdot)$. Moreover, since G is smooth for $t > 0$ and χ_ε is also smooth, we deduce that G_ε is smooth up to $t = 0$. In addition, due to the radial symmetry of M with respect to x , and of χ_ε , $G_\varepsilon(x, y, t)$ only depends on the radial coordinate $\rho = d(x, y)$ emanating from x and on t , hence we can write $G_\varepsilon(x, y, t) =: u_\varepsilon(\rho, t)$. In particular $2u_\varepsilon(\rho, t) = G_\varepsilon(x, \exp_x(\rho \vec{e}_1), t) + G_\varepsilon(x, \exp_x(-\rho \vec{e}_1), t)$. Then $\partial_\rho u_\varepsilon(0, t) = 0$ for $t > 0$ and a similar argument at the antipodal point shows $\partial_\rho u_\varepsilon(L, t) = 0$. Therefore, if we set $v_\varepsilon := e^{-\lambda t} \partial_\rho u_\varepsilon$, taking $u = u_\varepsilon$ in (2.3) and $v = v_\varepsilon$ in (2.4) we deduce that, for every $T > 0$,

$$\begin{cases} \partial_t v_\varepsilon + \mathcal{L} v_\varepsilon = 0 & \text{in } (0, L) \times (0, T), \\ v_\varepsilon(0, t) = v_\varepsilon(L, t) = 0 & \text{for all } t \in (0, T), \\ v_\varepsilon(\cdot, 0) = \partial_\rho g_\varepsilon \leq 0 & \text{in } [0, L]. \end{cases}$$

in $(0, L) \times (0, +\infty)$; see for instance [21, Theorem 7 in page 390] and the comments below it. Hence, the smoothness assumption is not really necessary here, we introduce it only for the sake of simplicity in the exposition.

Recalling that $\mathcal{L} = -\partial_{\rho\rho} + b\partial_{\rho} + c$ with $c \geq 0$, by the weak maximum principle [21, Theorem 9 in page 392] we see that $v_{\varepsilon} \leq 0$ in $[0, L] \times [0, T]$. Taking now $T \rightarrow +\infty$ we get that $\partial_{\rho}u_{\varepsilon}(\cdot, t) = e^{\lambda t}v_{\varepsilon} \leq 0$ for all $t > 0$, which means that $u_{\varepsilon}(\cdot, t)$ is nonincreasing for all $t > 0$ and all $\varepsilon > 0$. Finally, since $G_{\varepsilon}(x, y, t) \rightarrow G(x, y, t)$ for $t > 0$ as $\varepsilon \rightarrow 0$, we have $\Gamma(\rho, t) = \lim_{\varepsilon \rightarrow 0} u_{\varepsilon}(\rho, t)$ and, thus, $\Gamma(\cdot, t)$ is a nonincreasing function on $[0, L]$ for all $t > 0$.

Our goal now is to show that $\Gamma(\cdot, t)$ is indeed a strictly decreasing function on $[0, L]$ for all $t > 0$. This will mean that, for every $t > 0$, the heat kernel $G(x, y, t)$ is strictly decreasing as a function of the geodesic distance $\rho = d(x, y)$ emanating from x , which is the claim of the theorem.

Take $u = \Gamma$ and $v = e^{-\lambda t}\partial_{\rho}u$ as in (2.3). Since u satisfies (2.1), we have $(\partial_t + \mathcal{L})v = 0$ by (2.4). Also, we already know that $\partial_{\rho}\Gamma(\cdot, t) \leq 0$ for all $t > 0$, which yields $v \leq 0$ in $[0, L] \times (0, +\infty)$. Therefore, given $t_0 > 0$, if we set $w(\rho, t) := v(\rho, t + t_0)$, w solves

$$\begin{cases} \partial_t w + \mathcal{L}w = 0 & \text{in } (0, L) \times (0, +\infty), \\ w \leq 0 & \text{in } [0, L] \times [0, +\infty). \end{cases}$$

We will show that if t_0 is small enough then $w < 0$ in $(0, L) \times (0, +\infty)$. Arguing by contradiction, assume that $w(r, \tau) = 0$ for some $(r, \tau) \in (0, L) \times (0, +\infty)$. Since $w \leq 0$ in $[0, L] \times [0, +\infty)$, we deduce that w attains a nonnegative maximum over $[0, L] \times [0, \tau]$ at the interior point (r, τ) . Then, since $c \geq 0$, the strong maximum principle [21, Theorem 12 in page 399] yields that $w = 0$ in $[0, L] \times [0, \tau]$, which leads to $\partial_{\rho}\Gamma = 0$ in $[0, L] \times [t_0, t_0 + \tau]$. In particular, $\Gamma(\cdot, t_0)$ is constant on $[0, L]$. By taking t_0 small enough in the definition of w , we get a contradiction with the fact that $\Gamma(\rho, t) = G(x, y, t)$ tends to the Dirac delta centered at $x \in M$ as $t \rightarrow 0$ (and hence $\Gamma(\cdot, t_0)$ is not constant for $t_0 > 0$ small enough).

Once we know that $w < 0$ in $(0, L) \times (0, +\infty)$, we deduce that $\partial_{\rho}\Gamma < 0$ in $(0, L) \times (t_0, +\infty)$ and, as t_0 can be chosen arbitrarily small, we conclude that $\partial_{\rho}\Gamma(\cdot, t) < 0$ for all $t > 0$. Therefore, $\Gamma(\cdot, t)$ is a strictly decreasing function on $[0, L]$ for all $t > 0$. The theorem is finally proven. \square

Remark 2.1. *The previous proof still applies dropping the compactness condition if we can assure that $\frac{\partial^2}{\partial \rho^2} \log(S(\rho))$ remains bounded from above and decays at infinity. The proof of the next result requires to justify the decay condition for the noncompact case.*

Next let us show the proof of Theorem 1.5:

Proof of Theorem 1.5. The proof follows the same lines as the previous one. Since the boundedness of $\frac{\partial^2}{\partial \rho^2} \log(S(\rho))$ is part of the hypothesis, we just need to provide an exponential decay of the radial derivative of $G(x, \cdot, t)$. Under the geometric hypothesis in the statement Cheng, Li, and Yau have shown already that, for $t \in [0, T]$,

$$|\nabla G(x, y, t)| \leq C_T t^{(n+1)/2} e^{-cd(x,y)^2/t},$$

for some constants $c, C_T > 0$ independent on the distance $d(x, y)$ (cf. Theorem 6, [14, §4] p. 1055). This is enough to ensure the decay at infinity to finish the argument. Observe that one may use the maximum principle on a space-time box $B(r) \times [0, T]$, then the maxima should be achieved at the boundary $t = 0$. Indeed, the heat equation prevents a maximum to be achieved in $t = T$; on the other hand one can extend the box to be a band

by letting r tend to infinity (i.e. $M \times [0, T]$) since the boundary terms tend to zero (due to the exponential decay at infinity model that the aforementioned bound provides). \square

Remark 2.2. *The proof of Theorem 1.7 is completely analogous to the one of Theorem 1.2. Indeed, for Dirichlet heat kernels one observes that $G_D(x, y, t) \geq 0$ in the interior and it vanishes at the boundary. This shows that its radial derivative is less than or equal to zero at the boundary, the rest of the proof is analogous. On the other hand, for the Neumann heat kernel the boundary replaces the antipodal point and the Neumann condition on the boundary replaces the radial derivative that vanishes at the antipodal point.*

3. CONSEQUENCES REGARDING SPECIAL FUNCTIONS AND THE FRACTIONAL LAPLACE-BELTRAMI OPERATOR

In this section, we will present certain consequences of the aforementioned theorems regarding inequalities about classical special functions. To conclude we will also show some observations about pointwise properties of the fractional Laplace-Beltrami operator.

To that purpose, let us first recall that a function $F : \mathbb{R}^+ \rightarrow \mathbb{R}$ is said to be *completely monotonic* if $(-1)^n F^{(n)} > 0$ for $n \in \mathbb{Z}_{\geq 0}$. The central result about them is Bernstein's theorem [38, §IV.12] stating that any completely monotonic function F can be written as $F(u) = \int_0^\infty e^{-tu} d\mu(t)$ for some nonnegative measure $d\mu$.

If M is as in Theorem 1.2, with the notation employed there, we have the spectral expansion (e.g. [19, Lemma 2.1])

$$G(x, y, t) = \sum_{k=1}^{\infty} e^{-\lambda_k t} \varphi_k(x) \varphi_k(y)$$

with φ_k normalized eigenfunctions of the Laplace-Beltrami operator $-\Delta_M$ corresponding to the eigenvalues λ_k . Combining the conclusion of Theorem 1.2 and Bernstein's theorem we deduce

Corollary 3.1. *Let M and x be as in Theorem 1.2, and F a completely monotonic function such that the following series converges absolutely*

$$\sum_{k=1}^{\infty} F(\lambda_k) \varphi_k(x) \varphi_k(y).$$

Then it defines a strictly decreasing function of the geodesic distance $d(x, y)$.

By definition, the kernel of the heat equation with the spectral fractional Laplace-Beltrami operator is (see [20] for the \mathbb{S}^2 case)

$$G(x, y, t) = \sum_{k=1}^{\infty} e^{-\lambda_k^\alpha t} \varphi_k(x) \varphi_k(y) \quad \text{with } 0 < \alpha \leq 1.$$

As the function $F(x) = \exp(-x^\alpha)$ is completely monotonic for $0 < \alpha \leq 1$, we conclude at once

Corollary 3.2. *For M and x as in Theorem 1.2, consider the fractional heat equation $u_t + (-\Delta_M)^\alpha u = 0$ with $0 < \alpha \leq 1$ and $(-\Delta_M)^\alpha$ the spectral fractional Laplace-Beltrami operator on M . Then, its fundamental solution decreases as a function of the geodesic distance $d(x, y)$.*

This generalizes to other operators if the symbol has a completely monotonic derivative because a simple calculation [27] shows that if $F'(x)$ is completely monotonic then $\exp(-tF(x))$ is also completely monotonic for all $t > 0$.

For some particular manifolds M , there exists a full description of the eigenvalues and eigenfunctions in terms of special functions. In these cases, taking derivatives, Corollary 3.1 can be read as the positivity of some weighted sums of special functions. We exploit this relation for a compact manifold, for a manifold with boundary and for a noncompact manifold to get some inequalities involving averages of some classic special functions which seem to be new.

We start with the Legendre polynomials P_n . Many authors have considered inequalities involving P_n and its derivatives (see [34]). Arguably the cleanest is Fejér's inequality $\sum_{n=0}^N P_n(x) > 0$ for $-1 < x < 1$ that we still find in recent research [3]. There are several classic results by Hilb, Stieltjes, and other authors [34] showing intricate oscillations of $P_n(x)$ and $P'_n(x)$ when n and x vary and it is a theoretical obstacle to get general simple inequalities. Here we use Corollary 3.1 and the link between the Legendre polynomials and spectral expansions on the sphere to get a neat result.

Corollary 3.3. *Let F be a completely monotonic function such that $F(x) = O(x^{-\sigma})$ for some $\sigma > 2$. Then,*

$$\sum_{n=0}^{\infty} (2n+1)P'_n(x)F(n(n+1)) \geq 0 \quad \text{for } -1 \leq x \leq 1.$$

Thanks to the identity $(1-x^2)(2n+1)P'_n(x) = n(n+1)(P_{n-1}(x) - P_{n+1}(x))$, see [22, 8.914.2], the previous corollary can also be rephrased in terms of a sum of P_n .

Proof. It is well known [15, §VII.5] that the eigenvalues of the Laplace-Beltrami operator on \mathbb{S}^2 are $\lambda_n = n(n+1)$ with $n \in \mathbb{Z}_{\geq 0}$ and λ_n has multiplicity $2n+1$ with corresponding eigenfunctions $P_n^m(\cos \theta) \cos(m\varphi)$ with $0 \leq m \leq n$ and $P_n^m(\cos \theta) \sin(m\varphi)$ with $0 < m \leq n$, where P_n^m are the associate Legendre polynomials and (θ, φ) are the usual spherical coordinates. Using the spherical harmonic addition theorem [22, 8.814] we have that the series in Corollary 3.1 is

$$\sum_{n=0}^{\infty} (2n+1)P_n(\cos \theta)F(n(n+1))$$

(see [15, §VII.5.4] for the details) where $0 \leq \theta \leq \pi$ gives the geodesic distance. From the bounds $|P_n(\cos \theta)| \leq 1$ and $|P'_n(\cos \theta)| \leq P'_n(1) = n(n+1)/2$ (see [34, (7.33.8)]) and our hypothesis $F(x) = O(x^{-\sigma})$, the series can be differentiated term by term. The result follows because Corollary 3.1 implies that it is decreasing in θ . The inequality at the end points follows by continuity. \square

Now we are going to deduce an inequality for a sum of Bessel functions. It is not strictly a consequence of Corollary 3.1 because we are going to impose Dirichlet conditions but the proof keep the same scheme appealing to Theorem 1.7 instead of Theorem 1.2.

Corollary 3.4. *Let F be a completely monotonic function such that $F(x) = O(x^{-\sigma})$ for some $\sigma > 5/4$. Then,*

$$\sum_{n=1}^{\infty} \frac{z_n J_1(z_n r)}{J_1^2(z_n)} F(z_n^2) \geq 0 \quad \text{for } 0 \leq r \leq 1,$$

where J_1 is the Bessel function of order 1 and $\{z_n\}_{n=1}^{\infty}$ is the sequence of positive zeros of J_0 , the Bessel function of order 0.

Proof. Consider the heat equation $u_t - \Delta u = 0$ in the unit disk $D \subset \mathbb{R}^2$ under the Dirichlet boundary condition $u(\partial D, t) = 0$. It is well known that the radial eigenfunctions of Δu satisfying the boundary condition are $J_0(z_n r)$. Then by separation of variables the radial solutions are of the form [8, §II.34]

$$u(r, t) = \sum_{n=1}^{\infty} a_n e^{-z_n^2 t} J_0(z_n r).$$

On the other hand, we have the Fourier-Bessel series expansion [8, §I.17]

$$f(x) = \sum_{n=1}^{\infty} a_n J_0(z_n x) \quad \text{with } a_n = \frac{2}{J_1^2(z_n)} \int_0^1 s f(s) J_0(z_n s) ds.$$

To compute the fundamental solution we can choose the following approximation of the identity: $f(r) = (\pi \varepsilon^2)^{-1}$ if $0 \leq r < \varepsilon$ and 0 otherwise. It is clear that formally we get the coefficients $a_n = \pi^{-1} J_1^{-2}(z_n)$ when $\varepsilon \rightarrow 0^+$, simply using $J_0(0) = 1$. Then the fundamental solution should be

$$u(r, t) = \sum_{n=1}^{\infty} \frac{J_0(z_n r)}{\pi J_1^2(z_n)} e^{-z_n^2 t}.$$

By Theorem 1.7, for any given t , it defines a strictly decreasing function of $r \in (0, 1)$. Using Bernstein's theorem, we can replace $e^{-z_n^2 t}$ by $F(z_n^2)$ and the result, under our hypothesis on F , is differentiable term by term because $\sqrt{t} J_0(t)$, n/z_n and $(\sqrt{z_n} J_1(z_n))^{-1}$ are uniformly bounded, indeed the last two sequences tend to $1/\pi$ and to $-\sqrt{\pi/2}$, respectively [22, 8.548, 8.541.1]. This can be used to show, by direct computation, that $u(r, t)$ is indeed the fundamental solution we were seeking. The derivative of the resulting expression must be nonpositive and using $J_0' = -J_1$, the proof is complete. \square

We finish with a result involving the classical Mehler (conical Legendre) function $P_{-1/2+iv}$. The Mehler-Fock transform involving $P_{-1/2+iv}$ is a kind of hyperbolic version of the Fourier transform, and then the following result has the same flavor as some old results by Bochner and other authors (e.g., [7] [25]) in the Euclidean setting.

Corollary 3.5. *Let F be a completely monotonic function such that $F(x) = O(x^{-\sigma})$ for some $\sigma > 5/4$. Then*

$$\int_0^{\infty} F\left(\frac{1}{4} + v^2\right) P'_{-1/2+iv}(\cosh r) v \tanh(\pi v) dv \leq 0 \quad \text{for } r > 0.$$

Proof. The heat kernel in the hyperbolic plane \mathbb{H}^2 is (see [36, (3.37)])

$$\frac{1}{2\pi} \int_0^\infty e^{-(\frac{1}{4}+v^2)t} P_{-1/2+iv}(\cosh r) v \tanh(\pi v) dv.$$

Again, Bernstein's theorem allows to replace the exponential by $F(\frac{1}{4} + v^2)$ keeping the monotonicity of the result, assured by Theorem 1.5 or Theorem 1.1, which must have a nonpositive derivative.

The convergence of the integral and the differentiation under the integral sign is justified thanks to [22, 8.723.1], with the trivial estimate $O(1)$ for the hypergeometric function, and the fact that $\Gamma(iv)/\Gamma(1/2 + iv) \sim v^{-1/2}$ leads to $P'_{-1/2+iv} = O(v^{1/2})$. \square

Let us now proceed to the observation regarding local representations for fractional operators. Recently, great attention has been paid on the study of problems involving fractional and non-local operators. In particular, the fractional Laplace-Beltrami operator on manifolds appears in the study of different type of models such as free-boundary problems [37], porous medium equations [24], the Yamabe problem [10] and game theory, as they are related to stochastic Lévy flights [9]. In the Euclidean space or tori, it is a well-known consequence of Fourier analysis that the fractional Laplacian can be written as a singular integral operator (cf. [16]) which combined with elementary algebra provides useful pointwise estimates (cf. [17]). In the case of compact manifolds, an explicit integral representation for the fractional Laplace-Beltrami operator was only recently shown in [1] (cf. [20, 28] for the sphere). This new representation turned out to be essential in the proof of global existence of solutions to the surface quasi-geostrophic equation on the sphere (cf. [2]). Unfortunately, compared with the Euclidean case, the explicit representation contains an error term that makes it less powerful when pointwise control is needed, cf. [1]. As a consequence, several properties that are *straightforward* to check using the integral representation in the Euclidean case, as the maximum principle property or the positivity of the kernel, require more involved proofs. However, in the case of the unit sphere \mathbb{S}^n , an application of the decreasing property of Theorem 1.1 provides the following remark

Corollary 3.6. *Let $\alpha \in (0, 1)$. There is a function $k_{n,\alpha} : [0, \pi] \rightarrow \mathbb{R}^+$ such that $k_{n,\alpha}(0) = 1$, $d(x, y)^{-n-\alpha} k_{n,\alpha}(d(x, y))$ is decreasing for $d(x, y) \in (0, \pi)$, and*

$$(3.1) \quad (-\Delta_{\mathbb{S}^n})^\alpha f(x) = c_{n,\alpha} P.V. \int_{\mathbb{S}^n} \frac{f(x) - f(y)}{d(x, y)^{n+2\alpha}} k_{n,\alpha}(d(x, y)) d\text{vol}(y).$$

The main improvement of this result in comparison with the one obtained in [1] for general compact manifolds, where the operator is expressed as a similar principal value up to an operator that gains derivatives, is the absence of the latter.

Proof. By the definition the heat kernel $G(x, y, t)$, we have that

$$(3.2) \quad e^{-t\Delta_{\mathbb{S}^n}} f(x) = \int_{\mathbb{S}^n} G(x, y, t) f(y) d\text{vol}(y), \text{ and } \int_{\mathbb{S}^n} G(x, y, t) d\text{vol}(y) = 1.$$

Using the semigroup-formula (cf. [20, §7.1])

$$(3.3) \quad (-\Delta_{\mathbb{S}^n})^\alpha f(x) = \frac{1}{\Gamma(-\alpha)} \int_0^\infty (f(x) - e^{-t\Delta_{\mathbb{S}^n}} f(x)) \frac{dt}{t^{1+\alpha}},$$

combined with (3.2) and changing the order of integration it yields

$$(3.4) \quad \lim_{\varepsilon \rightarrow 0} \int_{\mathbb{S}^n \setminus B_\varepsilon(x)} (f(x) - f(y)) \int_0^\infty G(x, y, t) \frac{dt}{t^{1+\alpha}} d\text{vol}(y).$$

Notice that in order to apply Fubini's theorem, we have subtract a small ball of radius ε with $\varepsilon \rightarrow 0^+$. This limit is the principal value from the statement. For large times, the heat kernel is comparable to a constant, this is $c \leq G(x, y, t) \leq C$ (cf. [30]). Alternatively, one can show the upper bound using Hörmander's L^∞ bound coupled with the exponential decay of the coefficients. To deal with small times, one can either invoke the sharper estimate for $G(x, y, t)$ in [30, §1] or use the heat kernel parametrix expansion (cf. [2, §2], [11, §6] or [29, §1]) to check that the singularity in (3.4) is of order $-n - 2\alpha$. Combining both ingredients one can readily obtain

$$\int_0^\infty G(x, y, t) \frac{dt}{t^{1+\alpha}} = \frac{k_{n,\alpha}(d(x, y))}{d(x, y)^{n+2\alpha}}$$

for some function of the distance k which only depends on the distance $d(x, y)$, n and α . Moreover, we can assure that $G(x, y, t)$ is positive and decreasing as a function of $d(x, y) \in (0, \pi)$ by Theorem 1.1 applied to the sphere. \square

Remark 3.7. *Following the ideas sketched in Corollary 3.6, an interesting question is whether this observation could be extended to cover the case of compact hypersurface of revolution (as an application of Theorem 1.2) or complete spherically symmetric manifold with bounded curvature (as an application of Theorem 1.5).*

4. FINAL REMARKS

If the manifold is not spherically symmetric, one cannot hope a monotone behavior on the geodesic distance. For instance, for the flat torus $\mathbb{R}^2/(\mathbb{Z} \times L\mathbb{Z})$ with $L > 1$, the heat kernel based on the origin is

$$\frac{1}{L} \sum_{n,m \in \mathbb{Z}} e^{-4\pi^2 t(n^2 + m^2/L^2) + 2\pi i(nx + my/L)} = \frac{1}{4\pi t} \sum_{n,m \in \mathbb{Z}} e^{-((n-x)^2 + (Lm-y)^2)/4t},$$

where the equality follows from the Poisson summation formula. It is apparent that it decays faster in the second coordinate. On the other hand, in this example we can separate the variables and for y fixed the kernel decreases when $0 < x < 1/2$ and, in the same way, it decreases when $0 < y < L/2$ for x fixed. Any radial derivative is a combination of these, which shows that the heat kernel on a torus is decreasing in any radial direction departing from the origin up to the boundary of the fundamental domain.

It seems natural to try to decide whether there is a region with this monotonicity property in an arbitrary manifold. Notice that, in such a case, the region would depend on the initial point due to the possible lack of symmetries. An inspection of the sphere and the standard torus seems to suggest that, for every point, the monotonicity should hold up to its cut locus. This might be a by-product of some wishful thinking. On the other hand the sphere is, in some sense, the worst case scenario because the south pole is heated through every geodesic coming from the north pole. Perhaps an interesting context to start the study of these difficult questions is that of the hyperbolic Riemann surfaces, defined by $\Gamma \backslash \mathbb{H}^2$ with Γ a Fuchsian group. The constant curvature and the connection with the arithmetic theory of automorphic forms could be of some help in

some compact and noncompact examples. As a matter of fact, the heat kernels for tori are theta functions, which are prominent examples of automorphic forms.

Acknowledgements. This paper emerged from conversations during the long coffee breaks at the Harmonic Analysis in Winter Workshop 2018, held in Madrid. The authors are grateful to its organizers. A. Mas is grateful to M. Cozzi for pointing out the version of the parabolic maximum principle we apply in the proof. We sincerely thank the anonymous referees for the comments and suggestions which have helped to improve the quality of the paper.

D. Alonso-Orán, F. Chamizo and Á. D. Martínez were partially supported by the PID2020-113350GB-I00 grant of the MCINN (Spain) and the “Severo Ochoa Programme for Centres of Excellence in R&D” (SEV-2015-0554). A. Mas is supported by MTM2017-84214 and MTM2017-83499 projects of the MCINN (Spain), 2017-SGR-358 project of the AGAUR (Catalunya) and ERC-2014-ADG project HADE Id. 669689

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

- [1] D. Alonso-Orán, A. Córdoba and A.D. Martínez, Integral representation for fractional Laplace-Beltrami operators, *Adv. in Math.* 328, 436–445, 2018.
- [2] D. Alonso-Orán, A. Córdoba and A.D. Martínez, Global well-posedness of critical surface quasi-geostrophic equation on the sphere, *Adv. in Math.* 328, 248–263, 2018.
- [3] H. Alzer and M. K. Kwong. On Fejér’s inequalities for the Legendre polynomials. *Math. Nachr.*, 290(17-18), 2740–2754, 2017.
- [4] D. Andersson. Estimates of the Spherical and Ultraspherical Heat Kernel. Master’s thesis, Chalmers University of Technology, Sweden, 2013.
- [5] R. Azencott. Behavior of diffusion semi-groups at infinity. *Bull. Soc. Math. France*, 102: 193–240, 1974.
- [6] V. Banica, M. González and M. Sáez Some constructions for the fractional Laplacian on noncompact manifolds *Rev. Mat. Iberoam.*, 31, no. 2, 2015.
- [7] S. Bochner. Stable laws of probability and completely monotone functions. *Duke Math. J.*, 3(4):726–728, 1937.
- [8] F. Bowman. *Introduction to Bessel functions*. Dover Publications Inc., New York, 1958.
- [9] L. Caffarelli Non-local diffusions, drifts and games. *Nonlinear Partial Differential Equations*, Abel Symposia, 7, 37–52, 2012.
- [10] S. Y. A. Chang, M. d. M. González. Fractional Laplacian in conformal geometry. *Adv. Math.*, 226, 1410–1432, 2010.
- [11] I. Chavel. *Eigenvalues in Riemannian geometry*, volume 115 of *Pure and Applied Mathematics*. Academic Press, Inc., Orlando, FL, 1984. Including a chapter by Burton Randol, With an appendix by Jozef Dodziuk.
- [12] J. Cheeger and D. G. Ebin. Comparison theorems in riemannian geometry *Amsterdam Publisher, North-Holland Pub. Co*, 1975.
- [13] J. Cheeger and S. T. Yau. A lower bound for heat kernel, *Comm. Pur. Appl. Math.* 34, pp. 465–480, 1981.
- [14] S. Y. Cheng, P. Li and S. T. Yau. *On the Upper Estimate of the Heat Kernel of a Complete Riemannian Manifold* American Journal of Mathematics, Vol. 103, No. 5, pp. 1021-1063, 1981.
- [15] R. Courant and D. Hilbert. *Methods of mathematical physics. Vol. I*. Interscience Publishers, Inc., New York, N.Y., 1953.

- [16] A. Córdoba and D. Córdoba. A Maximum Principle Applied to Quasi-Geostrophic Equations, *Commun. Math. Phys.* 249, pp. 511–528, 2004.
- [17] A. Córdoba and D. Córdoba. A pointwise estimate for fractionary derivatives with applications to partial differential equations, *Proc. Natl. Acad. Sci. USA*100 (26), pp. 15316–15317, 2003.
- [18] A. Córdoba and A. D. Martínez. A pointwise inequality for fractional laplacians, *Adv. of Math.*, Vol. 280, pp. 79–85, 2015.
- [19] E. B. Davies, and B. Simon. Ultracontractivity and the heat kernel for Schrödinger operators and Dirichlet Laplacians. 335–395, *J. Funct. Anal.* 59, 1984
- [20] P. L. de Nápoli and P. R. Stinga. Fractional Laplacians on the sphere, the Minakshisundaram zeta function and semigroups, *New developments in the analysis of nonlocal operators*, 167–189, *Contemp. Math.* 723, *Amer. Math. Soc.*, 2019.
- [21] L. C. Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2010.
- [22] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series, and products*. Elsevier/Academic Press, Amsterdam, seventh edition, 2007.
- [23] A. Grigor'yan. Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds. *Bull. Amer. Math. Soc. (N.S.)*, 36(2):135–249, 1999.
- [24] G. Grillo, M. Muratori, F. Punzo, The porous medium equation with large initial data on negatively curved Riemannian manifolds. *Journal de Mathématiques Pures et Appliquées*, Vol.113, 195–226, 2018.
- [25] C. S. Herz. A class of negative-definite functions. *Proc. Amer. Math. Soc.*, 14:670–676, 1963.
- [26] S. Kobayashi and K. Nomizu. *Foundations of differential geometry. Vol. II*. Interscience Tracts in Pure and Applied Mathematics, No. 15 Vol. II. Interscience Publishers John Wiley & Sons, Inc., New York-London-Sydney, 1969.
- [27] M. Merkle. Completely monotone functions: a digest. In *Analytic number theory, approximation theory, and special functions*, pages 347–364. Springer, New York, 2014.
- [28] S. Minakshisundaram. Zeta functions on the unitary sphere. *Canadian Journal of Mathematics*, 4, 26–30, 1952.
- [29] S. Minakshisundaram, A. Pleijel Some Properties of the Eigenfunctions of The Laplace-Operator on Riemannian Manifolds. *Canadian Journal of Mathematics*, 1(3), 242-256, 1949
- [30] A. Nowak, P. Sjögren, and T. Z. Szarek. Sharp estimates of the spherical heat kernel. *Journal de Mathématiques Pures et Appliquées*, Vol. 129, Pages 23-33, 2019.
- [31] A. Nowak, P. Sjögren, and T. Z. Szarek. Genuily sharp heat kernel estimates on compact rank-one symmetric spaces, for Jacobi expansions, on a ball and on a simplex. arXiv:1905.1058 [math.CA], 2019.
- [32] M. H. Protter and H. F. Weinberger. *Maximum principles in differential equations*. Springer-Verlag, New York, 1984. Corrected reprint of the 1967 original.
- [33] C. Sogge. Concerning the L^p norm of spectral clusters for second-order elliptic operators on compact manifolds *J. Funct. Anal.*, Vol. 77, Issue 1, pages 123-138, 1988.
- [34] G. Szegő. *Orthogonal polynomials*. American Mathematical Society, Providence, R.I., fourth edition, 1975. American Mathematical Society, Colloquium Publications, Vol. XXIII.
- [35] M. E. Taylor. *Partial differential equations*, volume 23 of *Texts in Applied Mathematics*. Springer-Verlag, New York, 1996. Basic theory.
- [36] A. Terras. *Harmonic analysis on symmetric spaces—Euclidean space, the sphere, and the Poincaré upper half-plane*. Springer, New York, second edition, 2013.
- [37] E. Valdinoci. From the long jump random walk to the fractional Laplacian. *Bol.Soc. Esp. Mat. Apl. SeMA*, (49):33–44, 2009.
- [38] D. V. Widder. *The Laplace Transform*. Princeton Mathematical Series, v. 6. Princeton University Press, Princeton, N. J., 1941.

D. ALONSO-ORÁN

INSTITUTE FÜR ANGEWANDTE MATHEMATIK, UNIVERSITÄT BONN, ENDENICHER ALLEE 60, BONN 53115, GERMANY

Email address: `diego.alonso@icmat.es`

F. CHAMIZO

INSTITUTO DE CIENCIAS MATEMÁTICAS (CSIC-UAM-UC3M-UCM) – DEPARTAMENTO DE MATEMÁTICAS (UNIVERSIDAD AUTÓNOMA DE MADRID), 28049 MADRID, SPAIN

Email address: `fernando.chamizo@uam.es`

A. D. MARTÍNEZ

INSTITUTO DE CIENCIAS MATEMÁTICAS (CSIC-UAM-UC3M-UCM) – DEPARTAMENTO DE MATEMÁTICAS (UNIVERSIDAD AUTÓNOMA DE MADRID), 28049 MADRID, SPAIN

Email address: `angel.martinez@icmat.es`

A. MAS

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT POLITÈCNICA DE CATALUNYA, CAMPUS DIAGONAL BESÒS, EDIFICI A (EEBE), AV. EDUARD MARISTANY 16, 08019 BARCELONA, SPAIN

Email address: `albert.mas.bleesa@upc.edu`