

The geometry of Hermitian self-orthogonal codes

Simeon Ball* and Ricard Vilar

Abstract

We prove that if $n > k^2$ then a k -dimensional linear code of length n over \mathbb{F}_{q^2} has a truncation which is linearly equivalent to a Hermitian self-orthogonal linear code. In the contrary case we prove that truncations of linear codes to codes equivalent to Hermitian self-orthogonal linear codes occur when the columns of a generator matrix of the code do not impose independent conditions on the space of Hermitian forms. In the case that there are more than n common zeros to the set of Hermitian forms which are zero on the columns of a generator matrix of the code, the additional zeros give the extension of the code to a code that has a truncation which is equivalent to a Hermitian self-orthogonal code.

1 Introduction

The main motivation to study Hermitian self-orthogonal codes is their application to quantum error-correcting codes. The most prevalent and applicative quantum codes are qubit codes, in which the quantum state is encoded on n quantum particles with two-states. In this case, the quantum code is a subspace of $(\mathbb{C}^2)^{\otimes n}$. More generally, a quantum code is a subspace of $(\mathbb{C}^q)^{\otimes n}$. The parameter q is called the *local dimension* and corresponds to the number of states each quantum particle of the system has. A qubit is then referred to as a quqit.

A quantum code with minimum distance d is able to detect errors, which act non-trivially on the code space, on up to $d - 1$ of the quqits and correct errors on up to $\frac{1}{2}(d - 1)$ of the quqits. If the code encodes k logical quqits onto n quqits then we say the quantum code is an $[[n, k, d]]_q$ code. It has dimension q^k .

Suppose that $q = p^h$ is a prime power and let \mathbb{F}_q denote the finite field with q elements. A linear code C of length n over \mathbb{F}_q is a subspace of \mathbb{F}_q^n . If the minimum weight of a non-zero element of C is d then the minimum (Hamming) distance between any two elements of C is d and we

*The first (corresponding) author acknowledges the support of the Spanish Ministry of Science and Innovation grant MTM2017-82166-P.

say that C is a $[n, k, d]_q$ code, where k is the dimension of the subspace C . If we do not wish to specify the minimum distance then we say that C is a $[n, k]_q$ code.

A canonical Hermitian form on $\mathbb{F}_{q^2}^n$ is given by

$$(u, v)_h = \sum_{i=1}^n u_i v_i^q.$$

If C is a linear code over \mathbb{F}_{q^2} then its *Hermitian dual* is defined as

$$C^{\perp_h} = \{v \in \mathbb{F}_{q^2}^n \mid (u, v)_h = 0, \text{ for all } u \in C\}.$$

The standard dual of C will be denoted by C^\perp . Observe that $v \in C^\perp$ if and only if $v^q \in C^{\perp_h}$, so both of the dual codes have the same weight distribution.

One very common construction of quantum stabiliser codes relies on the following theorem from Ketkar et al. [9, Corollary 19]. It is a generalisation from the qubit case of a construction introduced by Calderbank et al. [4, Theorem 2].

Theorem 1. *If there is a $[n, k]_{q^2}$ linear code C such that $C \leq C^{\perp_h}$ then there exists an $[[n, n - 2k, d]]_q$ quantum code, where d is the minimum weight of the elements of $C^{\perp_h} \setminus C$ if $k \neq \frac{1}{2}n$ and d is the minimum weight of the non-zero elements of $C^{\perp_h} = C$ if $k = \frac{1}{2}n$.*

If $C \leq C^{\perp_h}$ then we say the linear code C is *Hermitian self-orthogonal*. Theorem 1 is our motivation to study Hermitian self-orthogonal codes. We can scale the i -th coordinate of all the elements of C by a non-zero scalar v_i , without altering the parameters of the code. Such a scaling, together with a reordering of the coordinates, gives a code which is said to be *linearly equivalent* to C .

Thus, a linear code D is *linearly equivalent* to a linear code C over \mathbb{F}_q if, after a suitable re-ordering of the coordinates, there exist non-zero $\theta_i \in \mathbb{F}_q$ such that

$$D = \{(\theta_1 u_1, \dots, \theta_n u_n) \mid (u_1, \dots, u_n) \in C\}.$$

A *truncation* of a code is a code obtained from C by deletion of coordinates. Observe that a truncation can reduce the dimension of the code but the dual minimum distance can only increase.

We will be interested in the following question: Given a linear $[n, k, d]_q$ code C , what truncations does C have which are linearly equivalent to Hermitian self-orthogonal codes?

In the special case that C is a k -dimensional Reed-Solomon code, the above question was answered by the authors in [3]. The code C has a truncation of length $m \leq q^2$ which is linearly equivalent to a Hermitian self-orthogonal code if and only if there is a polynomial $g(X) \in \mathbb{F}_{q^2}[X]$ of degree at most $(q-k)q-1$ with the property that $g(x) + g(x)^q$, when evaluated at the elements $x \in \mathbb{F}_{q^2}$, has precisely $q^2 + 1 - m$ zeros.

2 Hermitian self-orthogonal codes

Let C be a linear code of length n over \mathbb{F}_{q^2} . We have that C is linearly equivalent to a Hermitian self-orthogonal code if and only if there are non-zero $\theta_i \in \mathbb{F}_{q^2}$ such that

$$\sum_{i=1}^n \theta_i^{q+1} u_i v_i^q = 0,$$

for all $u, v \in C$. Note that θ_i^{q+1} is a non-zero element of \mathbb{F}_q , so equivalently C is linearly equivalent to a Hermitian self-orthogonal code if and only if there are non-zero $\lambda_i \in \mathbb{F}_q$ such that

$$\sum_{i=1}^n \lambda_i u_i v_i^q = 0.$$

For any linear code C over \mathbb{F}_{q^2} of length n , Rains [10] defined the *puncture code* $P(C)$ to be

$$P(C) = \{\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{F}_q^n \mid \sum_{i=1}^n \lambda_i u_i v_i^q = 0, \text{ for all } u, v \in C\}.$$

Then, clearly we have the following theorem.

Theorem 2. *There is a truncation of a linear code C over \mathbb{F}_{q^2} of length n to a linear over \mathbb{F}_{q^2} of length $r \leq n$ which is linearly Hermitian self-orthogonal code if and only if there is an element of $P(C)$ of weight r .*

Thus, as emphasised in [8], the puncture code is an extremely useful tool in constructing Hermitian self-orthogonal codes. Observe that, the minimum distance of any quantum code, given by an element in the puncture code, will have minimum distance at least the minimum distance of C^\perp , since any element in the dual of the shortened code will be an element of C^\perp if we replace the deleted coordinates with zeros.

Given a linear code C over \mathbb{F}_{q^2} it is not obvious how one can efficiently compute the puncture code. Let $G = (g_{i\ell})$ be a generator matrix for C , i.e. a $k \times n$ matrix whose row-span is C . A straightforward approach would be to construct a $\binom{k+1}{2} \times n$ matrix $T(G) = (t_{ij,\ell})$ over \mathbb{F}_{q^2} , where for $\{i, j\} \subseteq \{1, \dots, k\}$ we define

$$t_{ij,\ell} = \begin{cases} g_{i\ell} g_{j\ell}^q & i < j, \\ g_{i\ell}^{q+1} & i = j. \end{cases} \quad (1)$$

Lemma 3. *The puncture code $P(C)$ is the intersection of the right-kernel of $T(G)$ with \mathbb{F}_q^n .*

Proof. For any $u, v \in C$,

$$u_\ell = \sum_{i=1}^k a_i g_{i\ell} \quad \text{and} \quad v_\ell = \sum_{j=1}^k b_j g_{j\ell}$$

for some $(a_1, \dots, a_k) \in \mathbb{F}_q^k$ and $(b_1, \dots, b_k) \in \mathbb{F}_q^k$.

Since

$$\sum_{\ell=1}^n \lambda_\ell u_\ell v_\ell^q = \sum_{i=1}^k \sum_{j=1}^k a_i b_j^q \sum_{\ell=1}^n \lambda_\ell g_{i\ell} g_{j\ell}^q,$$

we have that $\lambda = (\lambda_1, \dots, \lambda_n)$ is in the right-kernel of $T(G)$ if and only if

$$\sum_{\ell=1}^n \lambda_\ell u_\ell v_\ell^q = 0,$$

for all $u, v \in C$. □

Thus, the puncture code $P(C)$ can then be found by extracting the elements in the right-kernel of $T(G)$ all of whose coordinates are elements of \mathbb{F}_q . However, this quickly becomes unfeasible computationally for larger parameters.

Our first aim, which we will deal with now, is to construct a parity check matrix for $P(C)$, i.e. a matrix over \mathbb{F}_q whose right-kernel is $P(C)$. This allows one to determine, given a linear code C over \mathbb{F}_{q^2} , all truncations of C which are linearly equivalent to a Hermitian self-orthogonal code, provided that the dimension of $P(C)$ is not too large.

Let $e \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$.

Let $M(G) = (m_{ij,\ell})$ be a $k^2 \times n$ matrix where, for $i, j \in \{1, \dots, k\}$, we define

$$m_{ij,\ell} = \begin{cases} e g_{i\ell} g_{j\ell}^q + e^q g_{i\ell}^q g_{j\ell} & i < j \\ g_{i\ell} g_{j\ell}^q + g_{i\ell}^q g_{j\ell} & i > j \\ g_{i\ell}^{q+1} & i = j \end{cases}.$$

Theorem 4. *The matrix $M(G)$ is a parity check matrix for $P(C)$. i.e. $M(G)$ is defined over \mathbb{F}_q and its right-kernel is $P(C)$.*

Proof. Observe first that all the entries in the matrix $M(G)$ are in \mathbb{F}_q .

Suppose that $\lambda = (\lambda_1, \dots, \lambda_n)$ is in the right-kernel of $M(G)$. Hence, for all $i, j \in \{1, \dots, k\}$ with $i < j$,

$$\sum_{\ell=1}^n \lambda_\ell (e g_{i\ell} g_{j\ell}^q + e^q g_{i\ell}^q g_{j\ell}) = 0$$

and

$$\sum_{\ell=1}^n \lambda_{\ell} (g_{j\ell} g_{i\ell}^q + g_{j\ell}^q g_{i\ell}) = 0.$$

Multiplying the latter equation by e^q and subtracting the former implies

$$(e^q - e) \sum_{\ell=1}^n \lambda_{\ell} g_{i\ell} g_{j\ell}^q = 0.$$

Since $\lambda = (\lambda_1, \dots, \lambda_n)$ is in the right-kernel of $M(G)$ we also have that

$$\sum_{\ell=1}^n \lambda_{\ell} g_{i\ell}^{q+1} = 0.$$

Hence, λ is in the right-kernel of $T(G)$.

Since it is also in \mathbb{F}_q^n , by Lemma 3, $\lambda \in P(C)$.

Suppose that $\lambda = (\lambda_1, \dots, \lambda_n) \in P(C)$. Then, for all $i, j \in \{1, \dots, k\}$,

$$\sum_{\ell=1}^n \lambda_{\ell} g_{i\ell} g_{j\ell}^q = 0.$$

This implies that λ is in the right-kernel of $M(G)$. □

Example 5. *Theorem 4 can allow us to efficiently calculate the puncture code of a linear code. Then for each codeword of weight r in the puncture code, by Theorem 2, we can construct a quantum error correcting code of length r . For example, let C be the linear $[43, 7]_4$ code, which is dual to the cyclic linear $[43, 36, 5]_4$ code, constructed from the divisor of $x^{43} - 1$,*

$$x^7 + ex^5 + x^4 + x^3 + e^2x^2 + 1,$$

where e is a primitive element of \mathbb{F}_4 .

By Theorem 4, we can calculate the puncture code from the 49×43 matrix M over \mathbb{F}_2 , which turns out to have rank 29. The puncture code $P(C)$ has weights $14 + 2j$ for all $j \in \{0, 1, 2, 3, 4, 5, 6, 7, 8\}$.

The truncations to codes of length 14 give a $[14, 7, 6]_4$ code which is equal to its Hermitian dual. By Theorem 1, this implies the existence of a $[[14, 0, 6]]_2$ quantum code.

The truncations to codes of length $18 + 2j$ give a $[18 + 2j, 7]_4$ code with dual minimum distance 5, which by Theorem 1 implies the existence of a $[[18 + 2j, 4 + 2j, 5]]_2$ quantum code, for all $j \in \{0, 1, 2, 3, 4, 5, 6\}$.

These codes equal the best known qubit error-correcting codes, according to Grassl [7].

Example 6. Consider the dual C to the cyclic linear $[51, 42, 6]_4$ code, constructed from the divisor of $x^{51} - 1$,

$$x^9 + e^2x^8 + ex^6 + x^5 + e^2x^4 + e^2x^2 + e^2x + 1.$$

The dimension of the puncture code $P(C)$ is 10. The puncture code $P(C)$ has codewords of weight $18 + 2j$, for all $j \in \{0, 2, 3, 4, 6, 7, 8\}$, which implies that it truncates to codes equivalent to Hermitian self-orthogonal codes of length $18 + 2j$. One can check these are $[18 + 2j, 9]_4$ codes with dual minimum distance 6. By Theorem 1, this implies the existence of a $[[18 + 2j, 2j, 6]]_2$ quantum code, for all $j \in \{0, 2, 3, 4, 6, 7, 8\}$.

Example 7. Consider C the $[15, 5]_9$ code with generator matrix

$$G = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 0 & 0 & 0 \\ e^7 & e^6 & e^5 & e^4 & 1 & e & e^3 & e^5 & e^4 & 1 & 0 & 0 & 1 & 0 & 0 \\ e^3 & e & e^4 & e^5 & 1 & e^6 & e^7 & e^4 & e^5 & 1 & 0 & 0 & 0 & 1 & 0 \\ e^6 & e^7 & e^5 & e^2 & e^4 & e^2 & e^6 & e^7 & e^3 & 1 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

The dual code C^\perp is a linear $[15, 10, 5]_9$ code. The dimension of the puncture code $P(C)$ is 2 and has codewords of weight 9, 12 and 15. This implies that it truncates to codes equivalent to Hermitian self-orthogonal codes of length 9, 12 and 15 and one can check that these codes are a $[9, 4]_9$, a $[12, 5]_9$ and a $[15, 5]_9$ codes all with dual minimum distance 5. By Theorem 1, this implies the existence of a $[[9, 1, 5]]_3$, a $[[12, 2, 5]]_3$ and a $[[15, 5, 5]]_3$ code. The former of these attains the quantum Singleton bound, proved by Rains in [10], which states that

$$k \leq n - 2(d - 1).$$

It was proven in [3] that a $[9, 4, 6]_9$ MDS code does not come from a truncation of a generalised Reed-Solomon code. The only $[9, 4, 6]_9$ code which is not the truncation of a generalised Reed-Solomon code is the projection of Glynn's $[10, 5, 6]_9$ MDS code, see [6].

3 The geometry of Hermitian self-orthogonal codes

Let $\text{PG}(k - 1, q)$ denote the $(k - 1)$ -dimensional projective space over \mathbb{F}_q .

A Hermitian form is given by

$$H(X) = \sum_{1 \leq i < j \leq k} (h_{ij} X_i X_j^q + h_{ij}^q X_i^q X_j) + \sum_{i=1}^k h_{ii}^{q+1} X_i^{q+1}.$$

for some $h_{ij} \in \mathbb{F}_{q^2}$.

The set of Hermitian forms is a k^2 -dimensional vector space over \mathbb{F}_q .

Let $G = (g_{i\ell})$ be a $k \times n$ generator matrix for a linear code C whose dual minimum distance is at least three. Let \mathcal{X} be the set of columns of G considered as points of $\text{PG}(k-1, q)$. Observe that the condition that the dual code of C has minimum distance at least three ensures that \mathcal{X} is a set (and not a multi-set). Such a code is often called a *projective code*. Observe that the set \mathcal{X} is the same for all codes linearly equivalent to C . Let $\text{HF}(\mathcal{X})$ be the subspace of Hermitian forms that are zero on \mathcal{X} .

Lemma 8. *The dimension of the left kernel of the matrix $M(G)$ is equal to $\dim \text{HF}(\mathcal{X})$.*

Proof. Let $x \in \mathcal{X}$ and consider a vector v in the left kernel of $M(G)$.

Observe that the coordinates of v are indexed by $i, j \in \{1, \dots, k\}$.

Since x is a column of G ,

$$\sum_{i,j=0}^k v_{ij}(ex_i x_j^q + e^q x_i^q x_j) + v_{ji}(x_i x_j^q + x_i^q x_j) + \sum_{i=1}^k v_{ii} x_i^{q+1} = 0.$$

Thus, defining

$$h_{ij} = ev_{ij} + v_{ji} \text{ and } h_{ii}^{q+1} = v_{ii},$$

we have that

$$H(x) = 0.$$

Letting v run over a basis for the left kernel of $M(G)$, we obtain a set of linearly independent Hermitian forms. Indeed, let B be a basis for the left kernel of $M(G)$. Suppose there are $\lambda_v \in \mathbb{F}_q$, for $v \in B$, not all zero, such that, for all $i, j \in \{1, \dots, k\}$,

$$\sum_{v \in B} \lambda_v (ev_{ij} + v_{ji}) = 0, \quad \sum_{v \in B} \lambda_v v_{ii} = 0.$$

Since $e \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, this implies

$$\sum_{v \in B} \lambda_v v_{ij} = 0,$$

for all $i, j \in \{1, \dots, k\}$, contradicting the fact that B is a basis.

Vice-versa, if $H(x) = 0$ for some Hermitian form H , then we obtain v_{ij} by solving

$$h_{ij} = ev_{ij} + v_{ji} \text{ and } h_{ij}^q = e^q v_{ij} + v_{ji},$$

where $v_{ij}, v_{ji} \in \mathbb{F}_q$, and $v_{ii} = h_{ii}^{q+1}$. Letting H run over a basis for $\text{HF}(\mathcal{X})$, we obtain a set of linearly independent vectors in the left kernel of the matrix $M(G)$. \square

The previous lemma allows us to calculate the dimension of the puncture code in terms of the dimension of the space of Hermitian forms which are zero on \mathcal{X} . In the following \mathcal{X} is obtained, as before, as the set of columns of a generator matrix for C , viewed as points of $\text{PG}(k-1, q)$. Note, that the statement that \mathcal{X} imposes r conditions on the space of Hermitian forms is to say that the co-dimension of $\text{HF}(\mathcal{X})$ is r .

Theorem 9. *The set \mathcal{X} imposes $n - \dim P(C)$ conditions on the space of Hermitian forms and*

$$\dim P(C) = n - k^2 + \dim \text{HF}(\mathcal{X}).$$

Proof. By Lemma 8,

$$\dim \text{HF}(\mathcal{X}) = \dim \text{left kernel } M(G) = k^2 - \text{rank } M(G).$$

By Theorem 4,

$$n - \text{rank } M(G) = \dim P(C),$$

which proves the second statement. For the first statement, observe that $\dim \text{HF}(\mathcal{X}) = k^2 - r$, where r is the number of conditions imposed by \mathcal{X} on the space of Hermitian forms. \square

Note that in the following statements the truncation may be the code itself.

Theorem 10. *The set of points \mathcal{X} imposes $|\mathcal{X}|$ conditions on the space of Hermitian forms if and only if no truncation of C is equivalent to a Hermitian self-orthogonal code.*

Proof. Theorem 9 implies that the set of points \mathcal{X} imposes n conditions on the space of Hermitian forms if and only if $\dim P(C) = 0$ which, by Theorem 2, is if and only if no truncation of C is equivalent to a Hermitian self-orthogonal code. \square

Thus, from Theorem 10, we deduce that to find codes contained in their Hermitian dual it is necessary and sufficient to find a set of points \mathcal{X} which does not impose $|\mathcal{X}|$ conditions on the space of Hermitian forms.

Theorem 11. *The set of points \mathcal{X} imposes less than $|\mathcal{X}|$ conditions on the space of Hermitian forms if and only if some truncation of C is linearly equivalent to a Hermitian self-orthogonal code.*

Theorem 11 has some immediate consequences.

Theorem 12. *A linear $[n, k]_{q^2}$ code for which $n > k^2$ has a truncation which is linearly equivalent to Hermitian self-orthogonal code.*

Proof. Since n is larger than the dimension of the space of Hermitian forms, \mathcal{X} cannot impose n conditions on the space of Hermitian forms. Hence, Theorem 11 implies the statement. \square

Example 13. Let e be a primitive element of \mathbb{F}_9 , where $e^2 = e + 1$. Let D be the cyclic linear $[73, 66, 6]_9$ code, constructed from the divisor of $x^{73} - 1$,

$$x^7 + ex^6 + e^6x^5 + e^3x^4 + e^7x^3 + e^2x^2 + e^5x + 2.$$

Let C be the $[60, 7]$ code obtained from D^\perp by deleting coordinates 61 to 73. The dimension of the puncture code $P(C)$ is 11. The puncture code $P(C)$ has codewords of weight $\{26, 27, \dots, 55\}$ which implies the existence of a $[[n, n - 14, 6]]_3$ quantum codes, for all $n \in \{26, 27, \dots, 55\}$.

The previous theorem and following theorem are the main results of this paper.

Theorem 14. A linear $[n, k]_{q^2}$ code C of length n over \mathbb{F}_{q^2} which has no truncations which are linearly equivalent to a Hermitian self-orthogonal code can be extended to C' , a $[n + 1, k]_{q^2}$ code which does have a truncation to a code which is linearly equivalent to a Hermitian self-orthogonal code, if and only if \mathcal{X} imposes n conditions on the space of Hermitian forms and the set of common zeros of $\text{HF}(\mathcal{X})$ is larger than $|\mathcal{X}|$.

Proof. (\Rightarrow) Let \mathcal{X}' be the set of columns of a generator matrix for C' obtained by extending the matrix G . By Theorem 9, both \mathcal{X} and \mathcal{X}' impose n conditions on the space of Hermitian forms. Hence,

$$\text{HF}(\mathcal{X}) = \text{HF}(\mathcal{X}')$$

which implies that the set of common zeros of $\text{HF}(\mathcal{X})$ contains \mathcal{X}' .

(\Leftarrow) Let $\mathcal{X}' = \mathcal{X} \cup \{x\}$ be a subset of the set of common zeros of $\text{HF}(\mathcal{X})$. Let C' be the code with generator matrix whose columns are the elements of \mathcal{X}' . Then \mathcal{X}' imposes n conditions on the space of Hermitian forms, so Theorem 9 implies that $\dim P(C') = 1$. Thus, C' extends C to a $[n + 1, k]_{q^2}$ code which has a truncation to a code which is linearly equivalent to a Hermitian self-orthogonal code. \square

Theorem 14 indicates that to extend a linear code C to a Hermitian self-orthogonal code, we should calculate the set of common zeros of the Hermitian forms which are zero on the columns of a generator matrix for C .

Example 15. The $[13, 7]_4$ code generated by the matrix

$$G = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & e & 0 & e^2 & e & e \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & e & e & e & 0 & e^2 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & e & e^2 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & e & 1 & 0 & e & 0 & e^2 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & e^2 & e^2 & e & e & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & e^2 & e^2 & e & 1 & e^2 & e \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & e^2 & e & e^2 \end{pmatrix}$$

has dual minimum distance 6. As before, let \mathcal{X} be the 13 points which are the columns of the matrix G . The dimension of $\text{HF}(\mathcal{X})$ is 36, so \mathcal{X} imposes 13 conditions on the space of Hermitian forms. Theorem 9 implies that $\dim P(C) = 0$, so C has no truncations which are linearly equivalent to Hermitian self-orthogonal codes. However, there are 14 points which are common zeros of the zeros of $\text{HF}(\mathcal{X})$, the points of \mathcal{X} and the point

$$(0, e, 0, 1, e, 1, 1).$$

Thus, Theorem 14 implies that the $[14, 7]_4$ code, with generator matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & e & 0 & e^2 & e & e & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & e & e & e & 0 & e^2 & e \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & e & e^2 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & e & 1 & 0 & e & 0 & e^2 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & e^2 & e^2 & e & e & 0 & e \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & e^2 & e^2 & e & 1 & e^2 & e & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & e^2 & e & e^2 & 1 \end{pmatrix}$$

has a truncation which is Hermitian self-orthogonal. Indeed, one can check that the code itself is Hermitian self-orthogonal. Thus, from this code we can construct, by Theorem 1, a $[[14, 0, 6]]_2$ code.

4 Conclusions and further work

In conclusion, we give a summary of the main results.

Suppose that C^\perp is a $[n, n - k, d]_{q^2}$, where $d \geq 3$.

If $n > k^2$ then we have shown that there are truncations of C which are linearly equivalent to Hermitian self-orthogonal codes.

If $n \leq k^2$ and $\dim P(C) > 0$ then there are truncations of C which are linearly equivalent to Hermitian self-orthogonal codes.

If $n \leq k^2$ and $\dim P(C) = 0$ and there are points which are not in \mathcal{X} but are zeros of the forms in $\text{HF}(\mathcal{X})$ then we can extend C to a $[n + 1, k]_{q^2}$ which does have truncations which are linearly equivalent to Hermitian self-orthogonal codes.

Finally, if $n \leq k^2$ and $\dim P(C) = 0$ and there are no points which are zeros of the forms in $\text{HF}(\mathcal{X})$ but which are not in \mathcal{X} then C has no extension to a $[n + 1, k]_{q^2}$ which has truncations that are linearly equivalent to Hermitian self-orthogonal codes. In this case we can extend C trying to maintain the dual minimum distance. This will reduce the dimension of $\text{HF}(\mathcal{X})$ by one, which then creates the possibility that there are points which are not in \mathcal{X} but are zeros of the forms in $\text{HF}(\mathcal{X})$. Indeed we can try and find extensions of C so that this is the case.

In all of the above we can construct a $[[r, r - 2k', d]]_q$ code from a truncation of length r , for some $k' \leq k$.

It should be able to extend these methods to make use of the following recent result of Galindo and Hernando [5, Theorem 1.2], which is an extension of Theorem 1.

There is also the possibility to extend these methods to self-orthogonal codes, i.e. $C \leq C^\perp$. This will work well in the case that the characteristic is even, since λ^{q+1} is replaced by λ^2 and all elements in a field of even characteristic have a square root. The role of the Hermitian form is then replaced by a quadratic form.

References

- [1] S. Ball, *A Course in Algebraic Error-Correcting Codes*, Compact Textbooks in Mathematics, Birkhäuser, 2020.
- [2] S. Ball, Some constructions of quantum MDS codes, *Des. Codes Cryptogr.*, **89** (2021) 811–821.
- [3] S. Ball and R. Vilar, Determining when a truncated generalised Reed-Solomon code is Hermitian self-orthogonal, [arXiv:2106.10180](https://arxiv.org/abs/2106.10180).
- [4] A. R. Calderbank, E. M. Rains, P. W. Shor, and N. J. A. Sloane, Quantum error correction via codes over $GF(4)$, *IEEE Trans. Inform. Theory*, **44** (1998) 1369–1387.
- [5] C. Galindo and F. Hernando, On the generalization of the construction of quantum codes from Hermitian self-orthogonal codes, [arxiv:2012.11998](https://arxiv.org/abs/2012.11998).
- [6] D. G. Glynn, The non-classical 10-arc of $PG(4, 9)$, *Discrete Math.*, **59** (1986) 43–51.
- [7] M. Grassl, Bounds on the minimum distance of linear codes and quantum codes, (available online at <http://www.codetables.de>).
- [8] M. Grassl and M. Rötteler, Quantum MDS codes over small fields, in *Proc. Int. Symp. Inf. Theory (ISIT)*, 1104–1108 (2015).
- [9] A. Ketkar, A. Klappenecker, S. Kumar and P. K. Sarvepalli, Nonbinary stabilizer codes over finite fields, *IEEE Trans. Inform. Theory*, **52** (2006) 4892–4914. (available online at <https://arxiv.org/abs/quant-ph/0508070>)
- [10] E. M. Rains, Nonbinary quantum codes, *IEEE Transactions on Information Theory*, **45** (1999) 1827–1832.
- [11] J. H. van Lint, *Introduction to Coding Theory*, Graduate Texts in Mathematics, **86**, Springer, 1999.

Simeon Ball

Departament de Matemàtiques,
Universitat Politècnica de Catalunya,
Carrer Jordi Girona 1-3,
08034 Barcelona, Spain
simeon@ma4.upc.edu

Ricard Vilar

Departament de Matemàtiques,
Universitat Politècnica de Catalunya,
Carrer Jordi Girona 1-3,
08034 Barcelona, Spain
ricard.vilar@upc.edu