

## Empty triangles in good drawings of the complete graph

Oswin Aichholzer · Thomas Hackl ·  
Alexander Pilz · Pedro Ramos ·  
Vera Sacristán · Birgit Vogtenhuber

Received: date / Accepted: date

**Abstract** A good drawing of a simple graph is a drawing on the sphere or, equivalently, in the plane in which vertices are drawn as distinct points, edges are drawn as Jordan arcs connecting their end vertices, and any pair of edges intersects at most once. In any good drawing, the edges of three pairwise connected vertices form a Jordan curve which we call a triangle. We say that a triangle is empty if one of the two connected components it induces does not contain any of the remaining vertices of the drawing of the graph. We show that the number of empty triangles in any good drawing of the complete graph  $K_n$  with  $n$  vertices is at least  $n$ .

**Keywords** Good drawings · Empty triangles · Erdős-Szekeres type problems

### 1 Introduction

Consider a simple graph  $G = (V, E)$ . A good drawing  $D(G)$  of  $G$  on the sphere  $S^2$  or, equivalently, in the Euclidean plane  $E^2$  is a drawing with the following properties:

---

Research of OA and AP was partially supported by the ESF EUROCORES programme EuroGIGA—CRP ‘ComPoSe’, Austrian Science Fund (FWF): I648-N18. Research of TH was supported by the Austrian Science Fund (FWF): P23629-N18 ‘Combinatorial Problems on Geometric Graphs’. During parts of this work, AP was a recipient of a DOC-fellowship of the Austrian Academy of Sciences at the Institute for Software Technology, Graz University of Technology, Austria. Research of PR was partially supported by MICINN Project MTM2011-22792, and ESF EUROCORES programme EuroGIGA, CRP ComPoSe: MICINN Project EUI-EURC-2011-4306, for Spain. Research of VS was partially supported by projects MTM2012-30951, MTM2009-07242, Gen. Cat. DGR 2009SGR1040, and ESF EUROCORES programme EuroGIGA, CRP ComPoSe: MICINN Project EUI-EURC-2011-4306, for Spain.

Oswin Aichholzer, Thomas Hackl, Alexander Pilz, and Birgit Vogtenhuber  
Institute for Software Technology, University of Technology, Graz, Austria  
E-mail: [oaich|thackl|apilz|bvogt]@ist.tugraz.at

Pedro Ramos  
Departamento de Matemáticas, Universidad de Alcalá, Madrid, Spain  
E-mail: pedro.ramos@uah.es

Vera Sacristán  
Departament de Matemàtica Aplicada II, Universitat Politècnica de Catalunya, Barcelona, Spain  
E-mail: vera.sacristan@upc.edu

1. The vertices are drawn as distinct points on the sphere  $S^2$  (or in the Euclidean plane  $E^2$ ).
2. The edges are Jordan arcs (i.e., non-self-intersecting continuous curves containing their end points) that have the drawings of the vertices they connect as end points.
3. Edges do not pass through any drawn vertex except for their own end vertices.
4. Any pair of edges intersects in at most one point (either in the interior of both edges, forming a proper crossing; or at a common end point).

Besides being a reasonable restriction for a natural drawing of a graph, a main interest in good drawings comes from the fact that they are useful for minimizing the number of crossings: It is well-known that if in a drawing of a graph there are edges which have self-intersections or pairs of edges which cross more than once, then the graph can be redrawn with fewer crossings. Therefore, only good drawings need to be considered when the goal is to make drawings with few crossings or to minimize the number of crossings. See for example [13, 14] for results on this topic.

In a good drawing  $D(G)$  of a graph  $G$ , the edges of any three pairwise connected vertices in  $D(G)$  form a Jordan curve (i.e., continuous non-self-intersecting curves that are closed in the sense that the two “end points” are identical), which we call a *triangle*. This definition matches the usual definition for the special case of straight-line drawings of  $G$ , i.e., drawings in the plane where edges are straight-line segments. Any triangle, being a Jordan curve, partitions the sphere (or the plane) into two connected components. If, in  $D(G)$ , one of these components does not contain the drawing of any of the remaining vertices, then the triangle is called *empty*. Further, for the case of a good drawing  $D(G)$  in the plane, one of the connected components induced by a triangle is bounded while the other one is unbounded. We denote the former as *interior* and the latter as *exterior* of the triangle. If, in  $D(G)$ , no vertex of  $G$  is drawn in the interior of a triangle, then the triangle is *interior-empty*. Likewise, if, in  $D(G)$ , no vertex of  $G$  is drawn in the exterior of a triangle, then the triangle is *exterior-empty*.

In this work, we consider the number of empty triangles in good drawings  $D(K_n)$  of the complete graph  $K_n$ . The question of finding empty triangles in good drawings of the complete graph goes back to Erdős’ question [3] about the existence of convex  $k$ -holes (empty polygons spanned by  $k$  vertices and edges) in straight-line drawings of the complete graph  $K_n$  and the subsequently posed question about their number [9]. For the existence question, it is by now well-known that every sufficiently large point set contains empty convex triangles, quadrilaterals, pentagons [6], and also hexagons [12, 5], but that there exist arbitrarily large point sets without empty convex heptagons [8]. While the existence question is trivial for empty triangles, the question on the least number  $h_3(n)$  of empty triangles in straight-line drawings of  $K_n$  has attracted many researchers and has been the topic of a large number of publications. The currently best known bounds for  $h_3(n)$  are  $n^2 - \frac{32}{7}n + \frac{22}{7} \leq h_3(n) \leq 1.6196n^2 + o(n^2)$ , where the upper bound is due to Bárány and Valtr [2] and the lower bound can be found in [1]. Note that both the upper and the lower bound are quadratic in  $n$ . Especially, every edge in any straight-line drawing of  $K_n$  is incident to at least one empty triangle.

In contrast, for general good drawings, Harborth [7] showed in 1989 that it is possible to draw  $K_n$  such that it contains only  $2n - 4$  empty triangles. Note that this implies that most edges are not incident to any empty triangle, while in straight-line drawings, every edge is incident to at least one empty triangle. Harborth mentioned in the same work that for  $3 \leq n \leq 6$ , the number of empty triangles in any good drawing  $D(K_n)$  is at least  $2n - 4$ . For  $n \geq 7$ , the best general lower bound he could show was 2. However, Harborth conjectured that every vertex in any drawing  $D(K_n)$  is incident to at least two empty triangles. Recently,

Fulek and Ruiz-Vargas [4] proved Harborth's conjecture to be true, thus providing a lower bound of  $\frac{2n}{3}$  for the number of empty triangles in any good drawing  $D(K_n)$ . In this paper we improve that bound and show that the number of empty triangles in any such drawing is at least  $n$ . Further, for  $n \leq 8$ , we show that Harborth's upper bound of  $2n - 4$  is still tight and we conjecture this to be the case in general.

*Outline.* Before proving our main theorem in Section 3, we review Fulek's and Ruiz-Vargas' proof in Section 2 and show that it allows to obtain additional properties of the considered empty triangles. Further, we investigate the relation between rotation systems and the task of computing the minimum number of empty triangles in Section 4 and present results for graphs with few vertices. In Section 5, we conclude by giving a short account on our conjecture that every good drawing contains at least  $2n - 4$  empty triangles.


Note that for many purposes, including counting empty triangles, drawings on the sphere  $S^2$  are equivalent to drawings in the plane  $E^2$  by Riemann stereographic projection<sup>1</sup>: In any good drawing  $D(K_n)$  of the complete graph  $K_n$ , let a *cell* (of  $D(K_n)$ ) be an open region (of  $S^2$  or  $E^2$ , respectively) whose boundary is defined by (parts of) drawn edges of  $K_n$  and which does not contain any part of  $D(K_n)$  (i.e., no part of a drawn edge or vertex of  $K_n$ ). Then for drawings in  $E^2$  exactly one cell is unbounded, while for drawings on  $S^2$  all cells are bounded. Now consider a drawing  $D(K_n)$  on  $S^2$  and an arbitrary cell  $C$  of  $D(K_n)$ . Applying Riemann stereographic projection with the projection center in  $C$ , one obtains a drawing  $D'(K_n)$  in  $E^2$  where the unbounded cell is the projection of  $C$ . Note that for every triangle  $\Delta$  in  $D(K_n)$ ,  $C$  is completely contained in one of the two connected components of  $S^2$  induced by  $\Delta$ . Further, note that the projection does not change any crossing properties of the edges. Thus, all vertices of  $K_n$  which are drawn in the connected component of  $S^2$  induced by  $\Delta$  that contains  $C$  lie in the exterior of the projection  $\Delta'$  of  $\Delta$ , while all vertices of  $K_n$  which are drawn in the other connected component of  $S^2$  induced by  $\Delta$  lie in the interior of  $\Delta'$ . Particularly,  $\Delta$  is empty if and only if  $\Delta'$  is (interior- or exterior-)empty. While both models are equivalent in that sense, in some parts of our reasoning it will be more convenient to consider the drawings in the plane rather than on the sphere. Especially, all the drawings in all figures are assumed to be in the plane.

## 2 Empty star triangles


For any subgraph or element  $H$  of a graph  $G$ ,  $D(H)$  denotes the drawing of  $H$  induced by  $D(G)$ . Recall that in a good drawing  $D(G)$  of a graph  $G$ , the edges incident to a vertex  $v$  only intersect in  $D(v)$ . Thus, the (drawing of the) graph consisting of all vertices of  $G$  and all edges incident to a vertex  $v$  of  $G$  is always crossing-free. We denote this graph as the *induced star graph* (of  $v$  in  $D(G)$ ). Note that  $D(G)$  induces a circular order of the edges incident to  $v$ ; see Fig. 1.

If for a triangle  $\Delta = D(uvw)$ , the (drawing  $D(uw)$  of the) edge  $uw$  is not crossed by any (drawing of an) edge incident to  $v$  in  $D(G)$ , then we say that  $\Delta$  is a *star triangle* (at  $v$  in  $D(G)$ ). In the drawing in Fig. 1,  $D(v_1v_2v_4)$  is a star triangle at  $v_1$ . For comparison,  $D(v_1v_2v_3)$  is not a star triangle at  $v_1$ , as the edge  $v_2v_3$  crosses the edge  $v_1v_4$  in  $D(G)$ .  $D(v_1v_5v_6)$  and  $D(v_1v_6v_2)$  are other star triangles at  $v_1$ . As can be seen in Fig. 1, the induced star graph of a vertex in a general graph might have isolated vertices. In contrast, the star

<sup>1</sup> Riemann stereographic projection is a projection from the plane to a tangent sphere (or back) where the projection center lies on the sphere and opposite to the tangent point of the plane.



**Fig. 1** The induced star graph of vertex  $v_1$  (drawn bold) in a good drawing of a graph. The vertices that are incident to  $v_1$  are labeled with respect to their circular order around  $v_1$  in  $D(G)$ .



**Fig. 2** Examples of induced star graphs (drawn bold) and star triangles in a good drawing  $D(K_6)$  of the complete graph  $K_6$ . Vertices are labeled with respect to their circular order around  $v_1$  in  $D(G)$ .  $D(v_1v_2v_4)$  is a star triangle at  $v_1$ .  $D(v_1v_2v_3)$  is not a star triangle at  $v_1$  (as the edge  $v_2v_3$  crosses the edge  $v_1v_4$  in  $D(G)$ ), but it is a star triangle at  $v_3$ .  $D(v_1v_5v_6)$  is an interior-empty star triangle at  $v_1$ .  $D(v_1v_3v_5)$  is an exterior-empty triangle. Further, it is a star triangle at  $v_5$ , but not at  $v_1$  or  $v_3$ .

graph of a vertex in the complete graph  $K_n$  always contains  $n - 1$  edges and connects all vertices of  $K_n$ , see Fig. 2.

This property can be used to obtain the following proposition about star triangles in good drawings of complete graphs.


**Proposition 1** Consider a good drawing  $D(K_n)$  of the complete graph  $K_n$  with  $n \geq 3$  and let  $\Delta = D(uvw)$  be a star triangle at  $v$ . Then  $\Delta$  is empty if and only if  $u$  and  $w$  are adjacent in the circular order of the edges around  $v$  (in  $D(G)$ ).

*Proof* Let  $H \subseteq K_n$  be the induced star graph of  $v$  plus the edge  $uw$ . As  $\Delta$  is a star triangle at  $v$ ,  $D(H)$  is crossing-free. Further,  $uvw$  is the only simple cycle in  $H$ . Let  $V_1$  and  $V_2$  be the subsets of vertices of  $K_n$  which are drawn in the two connected components induced by  $\Delta$ , respectively. As  $\Delta$  is a star triangle at  $v$ ,  $uw$  is not crossed by any of the edges of the induced star graph of  $v$ , where the latter contains an edge between  $v$  and each other vertex of  $K_n$ . Thus, all edges from  $v$  to vertices of  $V_1$  are drawn completely in one connected component induced by  $\Delta$ , and all edges from  $v$  to vertices of  $V_2$  are drawn completely in the other connected component induced by  $\Delta$ , implying that the circular order of the vertices around  $v$  is  $u, V_1, w, V_2$ . Hence,  $u$  and  $w$  are adjacent in this order if and only if  $V_1 = \emptyset$  or  $V_2 = \emptyset$ , which is equivalent to  $\Delta$  being interior-empty or exterior-empty.  $\triangle$

In [4, Proposition 3.1], Fulek and Ruiz-Vargas show that in a good drawing in the plane, every vertex is incident to at least one interior-empty triangle. They do so by explicitly finding such a triangle  $\Delta$ . We reconsider the proof of this proposition, showing that  $\Delta$  is in fact a star triangle at  $v$ .

**Proposition 2** *For every good drawing  $D(K_n)$  of the complete graph  $K_n$  in the Euclidean plane with  $n \geq 4$  vertices and every vertex  $v$  of  $K_n$ , there exists at least one interior-empty star triangle at  $v$  in  $D(K_n)$ .*

*Proof* Let  $H_0$  be the star graph of  $v$ , and let  $u_0$  be a vertex of  $K_n \setminus \{v\}$ ; see Fig. 3 for an accompanying example. By [4, Corollary 2.3], there is an edge  $u_0w_0$ , with  $w_0 \in K_n \setminus \{u_0, v\}$ , such that  $D(H_0 \cup \{u_0w_0\})$  is still a crossing-free drawing. Consider the triangle  $\Delta_0 = D(vu_0w_0)$ . If  $\Delta_0$  is interior-empty then it only remains to show that  $\Delta_0$  is a star triangle at  $v$ ; see below. Otherwise, let  $H_1 = H_0 \cup \{u_0w_0\}$ , and let  $u_1$  be a vertex of  $K_n$  whose drawing lies in the interior of  $\Delta_0$ . Repeating the argument, there is a vertex  $w_1 \in K_n \setminus \{u_1, v\}$  such that  $D(H_1 \cup \{u_1w_1\})$  is still crossing-free (and thus,  $D(w_1)$  lies in the interior or on the boundary of  $\Delta_0$ ). As  $\Delta_1 = D(vu_1w_1)$  contains strictly fewer vertices of  $K_n$  than  $\Delta_0$ , repeating this process terminates with an interior-empty triangle  $\Delta_i = D(vu_iw_i)$  in a crossing-free drawing  $D(H_i \cup \{u_iw_i\}) \subset D(K_n)$ .



**Fig. 3** Finding interior-empty star-triangles at  $v$ :  $H_i$  is drawn bold, the edge  $u_iw_i$  is drawn dashed, and  $\Delta_i$  is drawn shaded. (a) First step:  $\Delta_0$  is not interior-empty. (b) Second step:  $\Delta_1$  is interior-empty, so this is also the last step in this example.


Finally consider the interior-empty triangle  $\Delta_i$ ,  $i \geq 0$ , that has been found by this procedure. As  $D(H_i \cup \{u_iw_i\})$  contains the star graph of  $v$  in  $D(K_n)$  and is crossing-free,  $\Delta_i$  is a star triangle at  $v$ .  $\triangle$

Consider a good drawing  $D(K_n)$  of the complete graph  $K_n$  in the Euclidean plane ( $n \geq 4$ ) and let  $v$  be a vertex of  $K_n$ . By Proposition 2, there exists at least one interior-empty star triangle  $\Delta$  at  $v$ . Let  $C$  be a cell of  $D(K_n)$  which lies completely in the interior of  $\Delta$ . From  $D(K_n)$ , we obtain a good drawing  $D'(K_n)$  in the plane where the projection of  $\Delta$  is an exterior-empty triangle by applying Riemann stereographic projection twice: First project  $D(K_n)$  to the sphere. Then project the result back to the plane with the (new) projection center inside the projection of  $C$ , i.e., in  $D'(K_n)$   $C$  is the unbounded cell. Repeating the above proof to the drawing  $D'(K_n)$ , we obtain an interior-empty triangle  $\Delta'$  which is a star triangle at  $v$  in  $D'(K_n)$ . As the projection does not change any crossing properties of the

edges, the inverse projection of  $\Delta'$  is a star-triangle of  $v$  in  $D(K_n)$  as well (either interior- or exterior-empty). Similarly, if we have a good drawing on the sphere, we can first project it to the plane (by this making an arbitrary cell unbounded) and then apply the same arguments as above. Thus, we altogether obtain the following corollary.

**Corollary 1** *For every good drawing  $D(K_n)$  of the complete graph  $K_n$  with  $n \geq 4$  vertices and every vertex  $v$  of  $K_n$ , there are at least two empty star triangles at  $v$  in  $D(K_n)$ .*

We remark that, by Proposition 1, the triangles from Corollary 1 are formed with vertices that are consecutive in the order around  $v$ . Further, the bounds from Proposition 2 and Corollary 1 are tight in the sense that there exist drawings of  $K_n$  in the plane where most vertices are incident to exactly one interior-empty and one exterior-empty triangle, or to exactly two interior-empty and no exterior-empty triangles. See for example Fig. 4(a) for the former and Fig. 4(b) or Harborth's upper bound drawing [7, Fig. 1] for the latter.




**Fig. 4** Good drawings of  $K_8$  with  $2n - 4 = 12$  empty triangles where (a) vertices  $v_3$ ,  $v_4$ , and  $v_5$  are incident to exactly one interior-empty and one exterior-empty triangle, and (b) vertices  $v_3$ ,  $v_4$ , and  $v_5$  are incident to exactly two interior-empty and no exterior-empty triangles.

### 3 Lonely and lucky vertices

Consider a good drawing  $D(K_n)$  of the complete graph  $K_n$  and a vertex  $v$  in this drawing. If there exists a triangle  $\Delta$  in  $D(K_n)$  for which  $v$  is the only vertex drawn in one of the two connected components induced by  $\Delta$ , then we say that  $v$  is *lonely* in  $\Delta$ . For example, in the drawing in Fig. 2, vertex  $v_3$  is lonely in  $D(v_1v_2v_4)$  as it is the only vertex drawn in the interior of  $D(v_1v_2v_4)$ . Likewise, vertex  $v_5$  is lonely in  $D(v_2v_3v_6)$  as it is the only vertex drawn in the exterior of  $D(v_2v_3v_6)$ .


**Proposition 3** *If a vertex  $v$  of  $K_n$ ,  $n \geq 4$ , is lonely in a good drawing  $D(K_n)$ , then  $v$  is incident to at least three empty triangles in  $D(K_n)$ .*

*Proof* Consider a triangle  $\Delta = D(v_1v_2v_3)$  in  $D(K_n)$  which witnesses the loneliness of  $v$ , i.e.,  $v$  is the only vertex in one of the two connected components induced by  $\Delta$ . Further, consider the edges  $e_1 = vv_1$ ,  $e_2 = vv_2$ , and  $e_3 = vv_3$  between  $v$  and the three vertices of  $\Delta$ . As  $D(K_n)$  is a good drawing, at most one of  $e_1$ ,  $e_2$ , and  $e_3$  can form a crossing with an edge of  $\Delta$ . We distinguish two cases.



**Fig. 5** None of the edges between  $v$  and  $\{v_1, v_2, v_3\}$  crosses an edge of  $\Delta = D(v_1 v_2 v_3)$ : (a)  $v$  in the interior of  $\Delta$ , and (b)  $v$  in the exterior of  $\Delta$ .

*Case 1: None of  $e_1, e_2,$  and  $e_3$  forms a crossing with an edge of  $\Delta$ .* Note that in this case,  $e_1, e_2, e_3,$  and  $v$  are all completely in the same connected component induced by  $\Delta$ . Moreover, as none of the other vertices is on this side of  $\Delta$ , each of the three triangles formed by  $v$  and two vertices of  $\Delta$  is empty; see Fig. 5.



**Fig. 6** One of the edges between  $v$  and  $\{v_1, v_2, v_3\}$ , w.l.o.g.  $vv_1$ , crosses an edge of  $\Delta = D(v_1 v_2 v_3)$ : (a)  $v$  in the interior of  $\Delta$ ; (b)  $v$  in the exterior of  $\Delta$ ,  $vv_3$  drawn such that  $D(vv_2 v_3)$  is interior-empty; and (c)  $v$  in the exterior of  $\Delta$ ,  $vv_3$  drawn such that  $D(vv_2 v_3)$  is exterior-empty.

*Case 2: One of the edges  $e_1, e_2,$  or  $e_3$  forms a crossing with an edge of  $\Delta$ .* W.l.o.g., let  $e_1 = vv_1$  be this edge. Then the crossed edge of  $\Delta$  is  $v_2 v_3$ , and  $\Delta' = D(vv_2 v_3)$  is an empty triangle; see Fig. 6. As the edge  $v_2 v_3$  is crossed by  $vv_1$ ,  $\Delta'$  is none of the empty star triangles at  $v$  which are encountered by the proof of Proposition 2 and by Corollary 1. Thus, together with these two empty star triangles,  $v$  is incident to at least three different empty triangles.  $\triangle$

In the following, let  $l(v)$  be the number of triangles in  $D(K_n)$  in which  $v$  is lonely, and  $t(v)$  be the number of empty triangles in  $D(K_n)$  incident to  $v$ . If  $t(v) - l(v) \geq 2$  then we say that  $v$  is *lucky*. Note that for  $n \geq 4$ , every vertex  $v$  that is not lonely is lucky, as we know by Corollary 1 that in this case  $t(v) \geq 2$ . Also, every vertex  $v$  with  $l(v) = 1$  is lucky, as in this case  $t(v) \geq 3$  by Proposition 3.

**Theorem 1** For  $n \geq 4$ , the number of empty triangles in any good drawing  $D(K_n)$  of the complete graph  $K_n$  with  $n$  vertices is at least  $n$ .

*Proof* We prove the bound by induction on the number  $n$  of vertices. For the induction base, it is straightforward that for  $n = 4$ , every good drawing contains exactly four empty triangles; see again Fig. 5 and 6. So assume that the statement is true for any good drawing  $D(K_{n'})$  with  $n' < n$ , and consider a good drawing of  $D(K_n)$ . We distinguish two cases.

*Case 1:  $D(K_n)$  contains a vertex  $v$  which is lucky.* As  $v$  is lucky, we know that  $t(v) - l(v) \geq 2$ . Removing  $v$  and all its incident edges results in a drawing  $D(K_{n-1})$ . By the induction hypothesis, this drawing contains at least  $n - 1$  empty triangles. When adding  $v$  and all its incident edges again, the number of empty triangles is increased by  $t(v)$  and decreased by  $l(v)$ . Thus,  $D(K_n)$  contains at least  $n - 1 + t(v) - l(v) \geq n + 1 > n$  empty triangles.

*Case 2: All vertices of  $K_n$  are lonely in  $D(K_n)$ .* By Proposition 3, every lonely vertex is incident to  $t(v) \geq 3$  empty triangles. Summing up the number of incident empty triangles per vertex over all vertices, every triangle is counted exactly thrice (once for each of its vertices). Thus,  $\frac{1}{3} \cdot \sum_{v \in K_n} t(v) \geq \frac{1}{3} \cdot \sum_{v \in K_n} 3 = n$  is a lower bound for the number of empty triangles in  $D(K_n)$ .  $\triangle$

#### 4 Rotation systems and graphs with few vertices

While the emptiness of a triangle clearly depends on the drawing, not all information of the good drawing is needed to decide whether a triangle  $v_1v_2v_3$  is empty. While for deciding interior- or exterior-emptiness we need to know which side of the triangle contains the unbounded face, we can decide whether a triangle is empty by only looking at the rotation system of the drawing. Given a drawing of a graph  $G$  on an oriented surface, the *rotation system* of the drawing of  $G$  gives the circular order of the edges around each vertex of  $G$ . Let  $v_1v_2v_3$  be a triangle in a good drawing. The rotation system of  $v_2$  is separated by the edges  $v_2v_1$  and  $v_2v_3$  into two disjoint (possibly empty) sequences. For any fixed direction of the circular order around  $v_2$ , let  $R_2$  be the sequence between the edges  $v_2v_1$  and  $v_2v_3$ , and let  $L_2$  be the sequence between the edges  $v_2v_3$  and  $v_2v_1$ . For  $v_1$  and  $v_3$ , we define  $R_1, L_1$  as well as  $R_3$  and  $L_3$  analogously. We call a sequence  $R_i$  a *right* sequence and a sequence  $L_i$  a *left* sequence, for  $1 \leq i \leq 3$ . In any rotation system (and any good drawing) of the complete graph, edges from  $v_1, v_2$ , and  $v_3$  to any vertex  $v$  are trivially contained either in at least two left sequences or at least two right sequences; we then say that  $v$  is *left* of  $v_1v_2v_3$  or *right* of  $v_1v_2v_3$ , respectively. In a good drawing, the triangle is empty if either all other vertices are left of  $v_1v_2v_3$  or all other vertices are right of  $v_1v_2v_3$ ; see again Fig. 5 and 6.


Exhaustively generating all possible rotation systems of  $K_n$  for small  $n$  and counting the number of empty triangles therein can therefore easily be done. It remains to verify whether a rotation system is actually realizable, i.e., whether it is the rotation system of at least one good drawing. Deciding realizability can be done in a combinatorial way by considering a drawing as a crossing-free graph where (i) each vertex is either a vertex of the original graph or a crossing of the original graph, (ii) each edge is a part of an edge of the original graph, and (iii) each face is a cell of the original graph. For small point sets, a simple backtracking procedure that subsequently adds edges of the original graph and checks whether the drawing is good is sufficient and can be implemented in a straightforward way. Note that Kynčl [10] gives a more sophisticated, polynomial-time algorithm to decide realizability of a given rotation system of the complete graph. For  $3 \leq n \leq 6$ , Harborth mentioned in [7] that the number of empty triangles in any good drawing  $D(K_n)$  is at least  $2n - 4$ . By extensive computer search (i.e., generating all realizable rotation systems, which are 11556 for  $n = 7$  and 5370725 for  $n = 8$ , and then counting the number of empty

triangles for each of them), we have been able to confirm this result and show the same to be true also for  $n = 7$  and  $n = 8$ . For  $4 \leq n \leq 8$  the interested reader can find the number of all realizable rotation systems (#RS) versus the number of realizable rotation systems with  $2n - 4$  empty triangles (#minRS) in the following table.

n	4	5	6	7	8
#RS	2	5	102	11556	5370725
#minRS	2	2	10	33	146

**Observation 2** For  $3 \leq n \leq 8$ , the number of empty triangles in a good drawing of  $K_n$  is at least  $2n - 4$ .

If every drawing with few empty triangles contained a lucky vertex, then, by the proof of Theorem 1, the number of empty triangles would always be at least  $2n - 4$ . This is the case for the upper bound example from Harborth [7, Fig. 1], as well as for the drawings shown in Fig. 4; there, none of the vertices  $v_3, \dots, v_{n-2}$  is lonely in any triangle, and thus all of them are lucky. Unfortunately, the drawing in Fig. 7 shows that, in general, an argumentation like this one is not possible.



v	triangles incident to v	triangles witnessing $\ell(v)$
v1	$v_1 v_2 v_5, \underline{v_1 v_2 v_7}, v_1 v_2 v_8, v_1 v_5 v_6$	$v_2 v_3 v_4, v_2 v_4 v_5, \underline{v_2 v_7 v_8}$
v2	$v_1 v_2 v_5, \underline{v_1 v_2 v_7}, v_1 v_2 v_8, \underline{v_2 v_4 v_7}, v_2 v_5 v_8$	$v_1 v_3 v_8, v_1 v_4 v_8, v_1 v_5 v_8, v_1 v_6 v_8, v_1 v_7 v_8$
v3	$v_3 v_4 v_7, v_3 v_5 v_6, v_3 v_6 v_7$	$v_2 v_6 v_7, v_5 v_6 v_7$
v4	$\underline{v_2 v_4 v_7}, v_3 v_4 v_7, v_4 v_5 v_6, v_4 v_6 v_7$	$v_1 v_6 v_7, \underline{v_2 v_3 v_7}, v_6 v_7 v_8$
v5	$v_1 v_2 v_5, v_1 v_5 v_6, v_2 v_5 v_8, v_3 v_5 v_6, v_4 v_5 v_6, v_5 v_6 v_8$	$v_1 v_3 v_6, v_1 v_4 v_6, v_2 v_6 v_8, v_3 v_6 v_8, v_4 v_6 v_8$
v6	$v_1 v_5 v_6, v_3 v_5 v_6, v_3 v_6 v_7, v_4 v_5 v_6, v_4 v_6 v_7, v_5 v_6 v_8$	$v_1 v_3 v_5, v_1 v_4 v_5, v_3 v_5 v_7, v_3 v_5 v_8, v_4 v_5 v_8$
v7	$\underline{v_1 v_2 v_7}, \underline{v_2 v_4 v_7}, v_3 v_4 v_7, v_3 v_6 v_7, v_4 v_6 v_7$	$v_1 v_3 v_4, v_2 v_3 v_4, v_3 v_4 v_5, v_3 v_4 v_6, v_3 v_4 v_8$
v8	$v_1 v_2 v_8, v_2 v_5 v_8, v_5 v_6 v_8$	$v_2 v_5 v_6, v_2 v_5 v_7$

**Fig. 7** Example of a drawing of the only rotation system for  $K_8$  where every vertex is lonely in at least two triangles and no vertex is lucky. In the table, underlined triangles are the ones that are exterior-empty in the drawing; the others are interior-empty in the drawing.

Still, also in this drawing the total number of empty triangles equals  $2n - 4$ . Moreover, for  $n = 8$ , the drawing represents the only realizable rotation system (out of 5370725 different ones) for which there is no lucky vertex. All other realizable rotation systems of

cardinality  $4 \leq n \leq 8$ , no matter whether or not they have few empty triangles, contain at least one lucky vertex.

## 5 Conclusion

In this paper we have shown that any good drawing of the complete graph  $K_n$  with  $n$  vertices contains at least  $n$  empty triangles, thus improving the best previous lower bound of  $\frac{2n}{3}$ . As mentioned in the last section, Harborth already stated that the number of empty triangles for the best known minimizing examples is  $2n - 4$ , and we have confirmed this for  $n$  up to 8. We thus state the following conjecture.

*Conjecture.* For  $n \geq 4$ , the number of empty triangles in any good drawing of the complete graph  $K_n$  is at least  $2n - 4$ .

A *triangulation* is a maximal, crossing-free drawing of a graph such that every face is an empty triangle. In the plane, the outer face can be an exception, i.e., it might be a larger face. It is interesting to observe that any triangulation of  $n$  points on the sphere has  $2n - 4$  triangular faces. Equivalently, any triangulation of a set of  $n$  points in the plane with triangular convex hull consists of  $2n - 5$  triangles plus the outer, triangular face.

A *geometric graph* consists of vertices which are embedded as points in the plane, and edges which are straight line segments connecting two such points. It is easy to see that any complete geometric graph contains a maximal crossing-free sub-graph, that is, a triangulation. In contrast, it is NP-complete to decide whether a general (non-complete) geometric graph contains a triangulation as a sub-graph [11].

Note that in the non-geometric case a good drawing might contain  $2n - 4$  empty triangles, but, as these triangles might overlap, no triangulation as a sub-drawing. See for example Fig. 6(a) where in any crossing-free sub-drawing one of the faces has to be at least a quadrilateral. Even if we allow the outer face of a triangulation in the plane to be larger, there exist good drawings which do not contain such a triangulation as a sub-drawing. Thus we raise the following question:

*Open Problem.* What is the complexity of deciding whether or not a good drawing  $D(K_n)$  contains a triangulation as a sub-drawing?

**Acknowledgements** This work was initiated during a research visit of Pedro Ramos and Vera Sacristán in May 2013 in Graz, Austria.

## References

1. Aichholzer, O., Fabila-Monroy, R., Hackl, T., Huemer, C., Pilz, A., Vogtenhuber, B.: Lower bounds for the number of small convex  $k$ -holes. In: Proc. 24th Canadian Conference on Computational Geometry CCCG'12, pp. 247–252. Charlottetown, Canada (2012)
2. Bárány, I., Valtr, P.: Planar point sets with a small number of empty convex polygons. *Studia Scientiarum Mathematicarum Hungarica* **41**(2), 243–266 (2004)
3. Erdős, P.: Some more problems on elementary geometry. *Australian Mathematical Society Gazette* **5**, 52–54 (1978)
4. Fulek, R., Ruiz-Vargas, A.J.: Topological graphs: empty triangles and disjoint matchings. In: Proc. 29th ACM Symposium on Computational Geometry (SoCG'13), pp. 259–265. Rio de Janeiro, Brazil (2013)

- 
5. Gerken, T.: Empty convex hexagons in planar point sets. *Discrete and Computational Geometry* **39**(1–3), 239–272 (2008)
  6. Harborth, H.: Konvexe Fünfecke in ebenen Punktmengen. *Elemente der Mathematik* **33**, 116–118 (1978). In German
  7. Harborth, H.: Empty triangles in drawings of the complete graph. *Discrete Mathematics* **191**, 109–111 (1998)
  8. Horton, J.: Sets with no empty convex 7-gons. *Canadian Mathematical Bulletin* **26**(4), 482–484 (1983)
  9. Katchalski, M., Meir, A.: On empty triangles determined by points in the plane. *Acta Mathematica Hungarica* **51**(3–4), 323–328 (1988)
  10. Kynčl, J.: Simple realizability of complete abstract topological graphs in  $\mathbb{P}$ . *Discrete Comput. Geom.* **45**(3), 383–399 (2011)
  11. Lloyd, E.L.: On triangulations of a set of points in the plane. In: 18th Annual Symposium on Foundations of Computer Science, pp. 228–240 (1977)
  12. Nicolás, C.: The empty hexagon theorem. *Discrete and Computational Geometry* **38**(2), 389–397 (2007)
  13. Pan, S., Richter, R.B.: The crossing number of  $K_{11}$  is 100. *Journal of Graph Theory* **56**(2), 128–134 (2007)
  14. Richter, R.B., Thomassen, C.: Relations between crossing numbers of complete and complete bipartite graphs. *American Mathematical Monthly* **104**(2), 131–137 (1997)