

Secular instability in the spatial three-body problem

J. Féjoz,^{*} M. Guardia[†]

May 21, 2015

Abstract. Consider the spatial three-body problem, in the regime where one body revolves far away around the other two, in space, the masses of the bodies being arbitrary but fixed; in this regime, there are no resonances in mean motions. The so-called secular dynamics governs the slow evolution of the Keplerian ellipses. We show that it contains a horseshoe and all the chaotic dynamics which goes along with it, corresponding to motions along which the eccentricity of the inner ellipse undergoes large, random excursions. The proof goes through the surprisingly explicit computation of the homoclinic solution of the first order secular system, its complex singularities and the Melnikov potential.

1 Introduction

The question of the stability of the Solar System is the oldest open problem in Dynamical Systems. A number of works have shown striking instability mechanisms in the three-body problems, e.g. oscillatory orbits and close to parabolic motion [Ale69, Sit60, LS80a, LS80b, GMS13, DKdIRS14], chaotic dynamics near double or triple collisions [Bol06, Moe89] (see also [Mos01]). But scarce mathematical mechanisms have been described regarding more astronomical regimes, which would be plausible for subsystems of solar or extra-solar systems. One of them [FGKR15] shows the existence of global instabilities along mean motion (i.e. Keplerian) resonances.

In this paper, we focus on secular resonances. Numerical evidence has long been suggesting that such resonances are a major source of chaos in the Solar system [LR93, Las08, FS89]. For example, astronomers have established that Mercury's eccentricity is chaotic and can increase so much that collisions with Venus or the Sun become possible, as a result from an intricate network of secular resonances [BLF12]. On the other hand, that Uranus's obliquity (97°) is essentially stable, is explained, to a large extent, by the absence of any low-order secular resonance [BL10, LR93]. It is the goal of this paper to provide a simple instability mechanism in the secular dynamics of the three-body problem.

^{*}Université Paris-Dauphine (Ceremade) & Observatoire de Paris (IMCCE), France, jacques.fejoz@dauphine.fr.

[†]Universitat Politècnica de Catalunya (Departament de Matemàtica Aplicada I), Spain, marcel.guardia@upc.edu.

2 The secular Hamiltonian in the lunar regime

Consider three point masses undergoing the Newtonian attraction in space. Call m_0 , m_1 and m_2 the masses and $(q_j, p_j)_{j=1,2,3} \in (\mathbb{R}^3 \times \mathbb{R}^3)^3$ the Jacobi coordinates. After having symplectically reduced the system by the symmetry of translations, the Hamiltonian depends only on $(q_j, p_j)_{j=1,2}$ and equals

$$F_{\text{Kep}} + F_{\text{per}}, \quad \begin{cases} F_{\text{Kep}} = \sum_{j=1,2} \left(\frac{p_j^2}{2\mu_j} - \frac{\mu_j M_j}{\|q_j\|} \right) \\ F_{\text{per}} = \left(\frac{\mu_2 M_2}{\|q_2\|} - \frac{m_0 m_2}{\|q_2 + \sigma_1 q_1\|} - \frac{m_1 m_2}{\|q_2 - \sigma_0 q_1\|} \right), \end{cases}$$

where the reduced masses are defined by

$$M_0 = m_0, \quad M_1 = m_0 + m_1, \quad \mu_1^{-1} = M_0^{-1} + m_1^{-1}, \quad \mu_2^{-1} = M_1^{-1} + m_2^{-1} \quad (1)$$

and coefficients σ_j are the barycentric weights of the inner bodies: $\sigma_j = m_j/M_1$ ($j = 0, 1$). F_{Kep} is the integrable Hamiltonian of two uncoupled Kepler problems and induces a ‘‘Keplerian’’ action of the 2-torus, while F_{per} is the so-called perturbing function.

In this paper, we consider the asymptotic regime (sometimes called *lunar* or *hierarchical* or *stellar*) where the masses are fixed, while body 2 revolves far away around the other two. In particular, each of the two terms of F_{Kep} is negative, the outer eccentricity is bounded away from 1, and the semi major axes a_1 of q_1 and a_2 of q_2 satisfy

$$a_1 = O(1) \ll a_2 \rightarrow \infty.$$

The expansion of F_{per} in the powers of $\|q_1\|/\|q_2\| \ll 1$ is

$$F_{\text{per}} = -\frac{\mu_1 m_2}{\|q_2\|} \sum_{n \geq 2} \sigma_n P_n(\cos \zeta) \left(\frac{\|q_1\|}{\|q_2\|} \right)^n, \quad (2)$$

where the P_n 's are the Legendre polynomials, and $\sigma_n = \sigma_0^{n-1} + (-1)^n \sigma_1^{n-1}$, $n \geq 2$. Note that the perturbing function is of the order of $1/a_2^3$.

Standard normal form theory¹ shows that, for every integer k , there is a local change of coordinates a_2^{-3} -close to the identity which transforms the Hamiltonian into

$$F = F_{\text{Kep}} + F_{\text{sec}} + O(a_2^{-k-1}), \quad (3)$$

where F_{sec} is the *secular Hamiltonian*² of order k

$$F_{\text{sec}} = \int_{\mathbb{T}^2} F_{\text{per}} d\ell_1 d\ell_2 + O\left(\frac{1}{a_2}\right), \quad (4)$$

obtained by averaging out the mean anomalies ℓ_j at successive orders; \mathbb{T}^2 is the orbit of the Keplerian action defined by F_{Kep} , parameterized by the mean anomalies ℓ_1 and ℓ_2 of the two planets. The secular Hamiltonian descends to the quotient by the Keplerian action of \mathbb{T}^2 , and induces a Hamiltonian system on the space of pairs of Keplerian ellipses with fixed semi major axes, describing the slow evolution of the Keplerian ellipses due to the perturbation.

¹See, in this context, [F ej02]: mean motions do not interfere since they are not of the same order.

²*Secular*, means century in Latin. This term governs the slow dynamics (as opposed to the fast, Keplerian dynamics), on a long time scale, symbolically one century.

In this paper, we primarily consider the principal part of the Hamiltonian F of equation (3), i.e. $F_{\text{Kep}} + F_{\text{sec}}$.

We establish the phase portrait of the secular system, in particular with a (well known) hyperbolic periodic orbit located at inclined and nearly circular ellipses. The main results of the paper are the following. We prove that this periodic orbit possesses transversal homoclinic points (Corollary 4.2) and that the secular system possesses a horseshoe (Theorem 4.3), and therefore chaotic motions. As a by-product, we prove that the secular system is non-integrable. The results we obtain are valid for any value of the masses of the three bodies.

To prove these results, we take advantage of the fact that the first order in $1/a_2$ of the secular Hamiltonian is integrable. We explicitly determine the homoclinic solution to the hyperbolic periodic orbit of this first order (Lemma 3.1). Then, we show that the Melnikov potential associated with this homoclinic orbit has non-degenerate critical points (Proposition 4.1), from what it follows that the secular homoclinic solution splits for the full secular system, as well as for the full initial system.

The result we present in this paper is a step towards proving the existence of unstable orbits in the three body problem in the lunar regime, for any value of the masses of the bodies, in the sense of Arnold diffusion. By “unstable orbits” here we mean orbits such that one of the three bodies undergoes a large change in the semi major axis of the associated osculating ellipse. This is explained more precisely in Section 4.1.

2.1 The quadrupolar and octupolar Hamiltonians

The first two terms of the expansion (2) of F_{per} , after averaging, yield the *quadrupolar* and *octupolar* Hamiltonians:³

$$F_{\text{sec}} = -\mu_1 m_2 (F_{\text{quad}} + (\sigma_0 - \sigma_1)F_{\text{oct}}) + O\left(\frac{a_1^4}{a_2^5}\right), \quad (5)$$

with

$$F_{\text{quad}} = \frac{a_1^2}{8 a_2^3 (1 - e_2^2)^{3/2}} \left((15 e_1^2 \cos^2 g_1 - 12 e_1^2 - 3) \sin^2 i + 3 e_1^2 + 2 \right) \quad (6)$$

and

$$F_{\text{oct}} = -\frac{15 a_1^3}{64 a_2^4 (1 - e_2^2)^{5/2}} \times \quad (7)$$

$$\left\{ \begin{array}{l} \cos g_1 \cos g_2 \left[\frac{G_1^2}{L_1^2} (5 \sin^2 i (-7 \cos^2 g_1 + 6) - 3) \right] \\ - \sin g_1 \sin g_2 \cos i \left[\frac{G_1^2}{L_1^2} (5 \sin^2 i (7 \cos^2 g_1 - 4) + 3) \right] \end{array} \right\}.$$

this is a standard computation (see [Féj02] for the reduction to integrating trigonometric polynomials), and we have used the following notations:

$$\begin{cases} e_j & = \text{eccentricities} \\ g_j & = \text{arguments of pericenters} \\ i & = \text{mutual inclination.} \end{cases}$$

³More generally, the n -th order term is called 2^n -polar, because, in electrostatic, this term is the dominating term of the potential of a system of 2^n well chosen charged particles.

The Hamiltonians F_{quad} and F_{oct} do not depend on the order k of averaging in the expression (3) (i.e., due to the special dependence of F_{Kep} and F_{per} in L_2 , these first two terms of the normal form are not modified by second and higher order averaging).

Recall that we want to analyze the secular Hamiltonian F_{sec} for any value of the masses, so that parameters m_i , μ_i and M_i are just given constants satisfying (1).

Let $(\ell_j, L_j, g_j, G_j, \theta_j, \Theta_j)_{j=1,2}$ denote the Delaunay variables. Jacobi's classical elimination of the node consists in considering a codimension-3 submanifold of fixed, vertical angular momentum, and quotienting by horizontal rotations. The reduced manifold has dimension 8, on which the Keplerian and eccentric variables $(\ell_j, L_j, g_j, G_j)_{j=1,2}$ induce symplectic coordinates. After averaging out the mean motions we are left with the symplectic coordinates $(g_j, G_j)_{j=1,2}$, and the variables L_j may be treated as parameters.

The variables L_i are given by $L_i = \mu_i \sqrt{M_i a_i}$. Therefore, in the lunar regime we have $L_1 \sim 1$ and $L_2 \gg 1$. Recall that the eccentricity of the ellipse is given in terms of the Delaunay coordinates by

$$e_i = \sqrt{1 - \frac{G_i^2}{L_i^2}}. \quad (8)$$

Since we want the outer body to describe non-degenerate ellipses, we even assume $G_2 \sim L_2$. Since we are doing a perturbative analysis in L_2^{-1} , we define the new variable

$$\Gamma = C - G_2, \text{ where } C = \delta L_2 \text{ and } \delta > 0 \text{ is a fixed constant.}$$

The coordinates

$$(g_1, G_1, \gamma, \Gamma) = (g_1, G_1, -g_2, C - G_2), \quad (9)$$

are symplectic; we also call them Delaunay coordinates (as opposed to some radically different coordinates used later). Now these variables are bounded as $a_2, L_2 \rightarrow \infty$ (recall that $C \sim G_2 \rightarrow \infty$). At the first order in L_2^{-1} , the mutual inclination i satisfies

$$\cos^2 i = \frac{(C^2 - G_1^2 - G_2^2)^2}{4G_1^2 G_2^2} = \frac{(C^2 - G_1^2 - (C - \Gamma)^2)^2}{4G_1^2 (C - \Gamma)^2} = \frac{\Gamma^2}{G_1^2} + O(L_2^{-1}). \quad (10)$$

Now we express the secular Hamiltonian in these new variables and expand it in inverse powers of L_2 . Recall that variables L_j are parameters, as well as the norm C of the angular momentum.

Lemma 2.1. *The secular Hamiltonian (5) has the form*

$$F_{\text{sec}} = \alpha_0 + \alpha_1 L_2^{-6} \left(H_0 + L_2^{-1} H_1 + (\sigma_0 - \sigma_1) L_2^{-2} H_2 + \mathcal{O}(L_2^{-3}) \right). \quad (11)$$

with

$$H_0 = \left(1 - \frac{G_1^2}{L_1^2} \right) \left[2 - 5 \left(1 - \frac{\Gamma^2}{G_1^2} \right) \sin^2 g_1 \right] - \frac{\Gamma^2}{L_1^2} \quad (12)$$

In the coordinates $(g_1, G_1, \gamma, \Gamma)$, F_{quad} and hence H_0 and H_1 do not depend on γ . We do not compute H_1 explicitly, here. The reason is that H_1 does not break the integrability of H_0 and therefore does not play any role in the Melnikov analysis. In contrast, the Hamiltonian H_2 does break integrability (as it will follow from our study), and is computed in Section 5.

Proof of Lemma 2.1. Formula (8) implies

$$\frac{1}{G_1^2} = \frac{e_1^2}{G_1^2} + \frac{1}{L_1^2}.$$

Hence, using (10),

$$\sin^2 i = 1 - \frac{\Gamma^2}{G_1^2} + O(L_2^{-1}) = 1 - \Gamma^2 \frac{e_1^2}{G_1^2} - \frac{\Gamma^2}{L_1^2} + O(L_2^{-1}). \quad (13)$$

Besides,

$$e_2 = \sqrt{1 - \frac{G_2^2}{L_2^2}} = \sqrt{1 - \frac{C^2}{L_1^2}} + O(L_2^{-1})$$

is a first integral of the first order approximation of F_{quad} . Hence, expanding in powers of L_2^{-1} , F_{quad} can be written as

$$F_{\text{quad}}(g_1, G_1, \Gamma) = \alpha_1 L_2^{-6} (H_0(g_1, G_1, \Gamma) + 2) + O(L_2^{-1})$$

with

$$\alpha_1 = -\frac{L_1^4 M_2 \mu^6 m_2}{8M_1^2 \mu_1^3 (1 - e_2^2)^{3/2}}$$

and

$$\begin{aligned} H_0 &= (5e_1^2 \cos^2 g_1 - 4e_1^2 - 1) \sin^2 i + e_1^2 \Big|_{L_2^{-1}=0} \quad (\text{using (6)}) \\ &= e_1^2 \left[(5 \cos^2 g_1 - 4) \left(1 - \frac{\Gamma^2}{G_1^2} \right) + \frac{\Gamma^2}{G_1^2} + 1 \right] - \frac{\Gamma^2}{L_1^2} \quad (\text{using (10) and (13)}) \\ &= e_1^2 \left[2 - 5 \left(1 - \frac{\Gamma^2}{G_1^2} \right) \sin^2 g_1 \right] - \frac{\Gamma^2}{L_1^2}. \end{aligned}$$

Factorization (12) follows. \square

3 The quadrupolar phase portrait

According to (6), the quadrupolar Hamiltonian F_{quad} and thus H_0 do not depend on γ . This happy coincidence (which does not repeat itself for the next order term F_{oct}) makes F_{quad} integrable. This has been extensively used (see [Zha14] and references therein). Here, we may thus think of Γ as a parameter.

Complex singularities of solutions of F_{quad} are hard to determine in general. In our regime, the first order of the quadrupolar Hamiltonian, H_0 , can be factorized as described in equation (12) (up to the additive constant $-\Gamma^2/L_1^2$), which dramatically simplifies our study.

The Hamiltonian H_0 is analytic on a neighborhood of the cylinder

$$(g_1, G_1) \in \mathbb{T}^1 \times]0, L_1[$$

in $\mathbb{T}^1 \times \mathbb{R}$. Since it is π -periodic with respect to g_1 , we may focus on the domain $0 \leq g_1 \leq \pi$, $0 \leq G_1 \leq L_1$, keeping in mind that the Delaunay coordinates blow up circular ellipses ($G_1 = L_1$).

Hamilton's vector field is

$$\begin{cases} \dot{g}_1 = \frac{2G_1}{L_1^2} \left[5 \left(1 - \frac{\Gamma^2}{G_1^2} \right) \sin^2 g_1 - 2 \right] - 10 \left(1 - \frac{G_1^2}{L_1^2} \right) \frac{\Gamma^2}{G_1^3} \sin^2 g_1 \\ \dot{G}_1 = 5 \left(1 - \frac{G_1^2}{L_1^2} \right) \left(1 - \frac{\Gamma^2}{G_1^2} \right) \sin 2g_1. \end{cases}$$

The second component \dot{G}_1 vanishes if and only if (assuming $G_1 > 0$)

$$G_1 \in \{|\Gamma|, L_1\} \quad \text{or} \quad g_1 = 0 \pmod{\pi/2}.$$

If $G_1 = |\Gamma|$, \dot{g}_1 cannot vanish. Moreover, $G_1 \in]0, L_1[$. Let

$$\tilde{\Gamma} = \sqrt{1 - \frac{\Gamma^2}{L_1^2}} \in]0, 1[. \quad (14)$$

The equilibrium points thus satisfy the following equations:

- If $G_1 = L_1$ (circular ellipse),

$$\sin^2 g_1 = \frac{2}{5\tilde{\Gamma}^2}. \quad (15)$$

Assuming that

$$|\Gamma| < L_1 \sqrt{\frac{3}{5}}, \quad (16)$$

whence $\tilde{\Gamma}^2 \geq \frac{2}{5}$ (this condition is satisfied if the inner ellipse has a large eccentricity or an inclination close to $\pi/2$), within the g_1 -interval $[0, \pi[$, there are two solutions $g_1^{\min} \in]0, \pi/2[$ and g_1^{\max} symmetric with respect to $\pi/2$, which are located on the energy level $H_0 = -\Gamma^2/L_1^2$.

In variables (x, y) such that

$$\sin^2 g_1 = \frac{2}{5\tilde{\Gamma}^2}(1+x) \quad \text{and} \quad G_1 = L_1(1-y)$$

(which are local coordinates in the neighborhood of either one of the two above singularities), we have

$$H_0 + \Gamma^2/L_1^2 = 4xy + O_3(x, y);$$

thus the two singularities are hyperbolic.

- If $g_1 = 0 \pmod{\pi}$, $G_1 = 0$ (collision ellipse) is an equilibrium point of the regularized Hamiltonian vector field $G_1^3 X_{F_{\text{quad}}^0}$.
- If $g_1 = \pi/2 \pmod{\pi}$,

$$3G_1^4 - 10\Gamma^2 G_1^2 + 5L_1^2 \Gamma^2 = 0$$

(two pairs of opposite real solutions).




Figure 1: Phase portrait of the reduced quadrupolar dynamics

We focus on the first two hyperbolic singularities, with $G_1 = L_1$, henceforth assuming that condition (16) is satisfied, with, say,

$$\Gamma > 0. \quad (17)$$

Lifted to the full secular phase space, these critical points become the periodic orbits

$$Z_{\min, \max}^0(t, \gamma_0) = \left(g_1^{\min, \max}, L_1, \gamma^0 + \gamma^1(t), \Gamma \right), \quad (18)$$

where γ^0 is just the initial condition for the variable γ and

$$\gamma^1(t) = -2 \frac{\Gamma}{L_1^2} t. \quad (19)$$

The whole circle $G_1 = L_1$ corresponds to circular motion. Along this circle the Delaunay variables are singular. Thus, in a neighborhood of circular motion it is more reliable to use the Poincaré variables

$$\begin{cases} \xi &= \sqrt{2(L_1 - G_1)} \cos g_1 \\ \eta &= -\sqrt{2(L_1 - G_1)} \sin g_1. \end{cases} \quad (20)$$

Those secular coordinates are symplectic (a well known fact). The Poincaré variables transform the Hamiltonian H_0 into

$$\tilde{H}_0(\xi, \eta) = -\frac{\Gamma^2}{L_1^2} + \frac{1}{L_1} \left[2\xi^2 - \left(3 - 5 \frac{\Gamma^2}{L_1^2} \right) \eta^2 \right] + \mathcal{O}_2(\xi^2 + \eta^2) \quad (21)$$

and the line segment $\{G_1 = L_1\}$ blows down to a single hyperbolic periodic orbit

$$(\xi, \eta, \gamma, \Gamma) = (0, 0, \gamma^0 + \gamma^1(t), \Gamma)$$

in the secular space.

As it has occurred above, in the (g_1, G_1) -plane there are two hyperbolic fixed points for $g_1 \in [0, \pi]$ and two more for $g_1 \in [\pi, 2\pi]$. On the line $G_1 = L_1$, there are heteroclinic connections between them. All these objects blow down to the critical point $(\xi, \eta) = (0, 0)$ in Poincaré variables. Moreover, there are two separatrix connections in the region $G_1 < L_1$, one between the critical points with $g_1 \in [0, \pi]$ and the other one between the critical points with $g_1 \in [\pi, 2\pi]$. Even if H_0 is well defined for $G_1 > L_1$, the corresponding solutions are spurious. In the secular space, we obtain a hyperbolic critical point at $(\xi, \eta) = (0, 0)$ with two homoclinic orbits forming a figure eight.

The main technical goal of this paper is to show that these separatrices split when one considers the complete secular Hamiltonian of arbitrary order k . We focus on the homoclinic on which $g_1 \in [0, \pi]$. We use both Delaunay and Poincaré variables, but need to keep in mind that the Delaunay variables are not defined on the secular space proper along circular inner ellipses ($G_1 = L_1$). Note the same orbit is homoclinic for the secular Hamiltonian (21) and heteroclinic after blow up (see (12)).

The expression of the energy (12) yields the following characterization of this orbit, where the interval $]g_1^{\min}, g_1^{\max}[\subset]0, \pi[$ is defined by the inequality (recall (14))

$$\sin^2 g_1 > \frac{2}{5\Gamma^2} \quad (22)$$

and

$$\chi = \sqrt{\frac{2|\Gamma|}{3L_1}} \frac{1}{\sqrt{1 - \frac{5\Gamma^2}{3L_1^2}}} > 0 \quad \text{and} \quad A_2 = 30 \sqrt{\frac{10|\Gamma|^3}{3L_1}} \sqrt{1 - \frac{5\Gamma^2}{3L_1^2}}. \quad (23)$$

Lemma 3.1. *Assume conditions (16)–(17). The Hamiltonian H_0 given in (12) has a heteroclinic solution which tends to the periodic orbits Z_{\min}^0 and Z_{\max}^0 in (18) in backward and forward times respectively. Its orbit is defined by the equation*

$$\left(1 - \frac{\Gamma^2}{G_1^2}\right) \sin^2 g_1 = \frac{2}{5}, \quad (24)$$

with range $g_1 \in]g_1^{\min}, g_1^{\max}[\subset]0, \pi[$. Its time parameterization is given by

$$Z^0(t, \gamma^0) = (g_1^h(t), G_1^h(t), \gamma^h(t), \Gamma) \quad (25)$$

with

$$\cos g_1^h(t) = -\sqrt{\frac{3}{5}} \frac{\sinh A_2 t}{\sqrt{\chi^2 + (1 + \chi^2) \sinh^2 A_2 t}} \quad (26)$$

$$G_1^h(t) = |\Gamma| \sqrt{\frac{5}{3}} \sqrt{1 + \frac{3}{5} \frac{L_1^2}{\Gamma^2} \sinh^2 A_2 t} (\cosh A_2 t)^{-1} \quad (27)$$

$$\gamma^h(t) = \gamma^0 + \gamma^1(t) + \gamma^2(t), \quad \gamma^2(t) = \arctan(\chi^{-1} \tanh A_2 t), \quad (28)$$

where $\gamma^0 \in \mathbb{T}$ is the arbitrary value of the γ coordinate at initial time, γ^1 has been introduced in (19).

In the γ -component of the separatrix, the angle $-\gamma^0$ is the (arbitrary) argument of the outer pericenter at the symmetry center of the separatrix $g_1 = \pi/2$ (recall the change of coordinates (9)). The term $\gamma^1(t)$ is the rotational part and the term $\gamma^2(t)$ is the transient part of $\gamma^h(t)$.

Remark 3.2. The assumed condition (16) implies that $\partial_\Gamma H_0 \neq 0$ in a neighborhood of the heteroclinic orbit. Therefore, one can rephrase Lemma 3.1 as the existence of a heteroclinic orbit in each energy level $H_0(g_1, G_1, \gamma, \Gamma) = h$ for any $h < 0$ (recall that by the expression of H_0 in (12) the heteroclinic orbits always have negative energy).

Proof. On the separatrix, we have

$$\dot{g}_1 = 10\Gamma^2 \frac{1 - G_1^2/L_1^2}{G_1^3} \sin g_1^2.$$

Using (24) and (22), one can eliminate G_1 by

$$G_1 = \frac{|\Gamma| \sqrt{5} \sin g_1}{\sqrt{3 - 5 \cos^2 g_1}}, \quad (29)$$

and get a closed differential equation

$$\dot{g}_1 = A_1 \frac{(1 - \frac{5}{3}(1 + \chi^2) \cos^2 g_1) \sqrt{1 - \frac{5}{3} \cos^2 g_1}}{\sin g_1} \quad (30)$$

where χ has been defined in (23) and

$$A_1 = 30\sqrt{3}\Gamma^2 \left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2}\right) > 0.$$

Note that, due to $\Gamma \neq 0$ and (22), $\dot{g}_1 \neq 0$ along the separatrix. So, one can separate variables and, choosing $g_1 = \pi/2$ at $t = 0$ yields

$$A_1 t = \int_{\pi/2}^{g_1} \frac{\sin g_1 dg_1}{\left(1 - \frac{5}{3}(1 + \chi^2) \cos^2 g_1\right) \sqrt{1 - \frac{5}{3} \cos^2 g_1}}.$$

The variable $\cos g_1 \in \left] -\frac{\sqrt{3}}{\sqrt{5(1+\chi^2)}}, \frac{\sqrt{3}}{\sqrt{5(1+\chi^2)}} \right[$ is a coordinate on the separatrix. Let x be defined by

$$\sqrt{\frac{5}{3}} \cos g_1 = \cos x \in \left] 0, \frac{1}{(1 + \chi^2)} \right[.$$

Then

$$\begin{aligned} A_1 t &= \sqrt{\frac{3}{5}} \int_{\pi/2}^{\arccos \sqrt{\frac{5}{3}} \cos g_1} \frac{dx}{1 - (1 + \chi^2) \cos^2 x} \\ &= \frac{1}{2\chi} \sqrt{\frac{3}{5}} \ln \frac{\tan x - \chi}{\tan x + \chi}, \quad \text{with } \cos x = \sqrt{\frac{5}{3}} \cos g_1. \end{aligned}$$

One can check that the constant A_2 defined in (23) satisfies $A_2 = \sqrt{\frac{5}{3}} A_1 \chi$. The equation can be solved for $\tan x$, which yields $\tan x = -\chi \coth(A_2 t)$. This equality, jointly with

$$\cos g_1 = \sqrt{\frac{3}{5}} \cos x = \sqrt{\frac{3}{5}} \frac{\pm 1}{\sqrt{1 + \tan^2 x}},$$

gives formula (26). Using (29), the G_1 -component of the separatrix given in (27) can be easily obtained. Finally, only the γ -component remains to be computed. But, along the separatrix solution, $\dot{\gamma} = \partial_\Gamma H_0$. Using (12), (26) and (29),

$$\begin{aligned} \dot{\gamma} &= 10 \frac{\Gamma}{G_1^2} \left(1 - \frac{G_1^2}{L_1^2}\right) \sin^2 g_1 - 2 \frac{\Gamma}{L_1^2} \\ &= \frac{4\Gamma}{L_1^2 \chi^2} \left(1 - \frac{5}{3}(1 + \chi^2) \cos^2 g_1\right) - 2 \frac{\Gamma}{L_1^2} \\ &= \frac{4\Gamma}{L_1^2} \frac{1}{((1 + \chi^2) \cosh^2 A_2 t - 1)} - 2 \frac{\Gamma}{L_1^2}. \end{aligned}$$

Since $\frac{4\Gamma}{L_1^2 \chi A_2} = \frac{4\Gamma \sqrt{3}}{L_1^2 \chi^2 A \sqrt{5}} = 1$, equation (28) follows. \square

4 Splitting of separatrices

Lemma 3.1 shows that the Hamiltonian H_0 in (6) has a heteroclinic connection (homoclinic in the secular space for the Hamiltonian (21)).

The next term in the asymptotic expansion of the secular Hamiltonian given in Lemma 2.1 does not depend on γ and therefore does not break integrability. For the amended Hamiltonian, according to classical perturbation theory, the hyperbolic periodic orbit and the homoclinic orbit persist. Moreover, the periodic orbit remains at $(\xi, \eta) = (0, 0)$. Thus, in Delaunay coordinates, there are two hyperbolic periodic orbits, which are of the form

$$\begin{cases} Z_{\min}^\varepsilon(t, \gamma^0) &= (g_{\min} + O(\varepsilon), L_1, \gamma^0 + \gamma^1(t) + O(\varepsilon), \Gamma) \\ Z_{\max}^\varepsilon(t, \gamma^0) &= (g_{\max} + O(\varepsilon), L_1, \gamma^0 + \gamma^1(t) + O(\varepsilon), \Gamma), \end{cases}$$

where $\varepsilon = L_2^{-1}$. We can choose the time origin so that

$$Z_{\min, \max}^\varepsilon(0, \gamma^0) = (g_{\min, \max} + O(\varepsilon), L_1, \gamma^0, \Gamma).$$

The second order H_2 in Lemma 2.1 depends on γ and therefore may (and will) break integrability. If we consider this Hamiltonian in Poincaré coordinates (20), it possesses a hyperbolic critical orbit $O(L_2^{-2})$ -close to $(\xi, \eta) = (0, 0)$.

To show that H_2 makes the stable and unstable invariant manifolds of this periodic orbit split, we use the classical Poincaré-Melnikov Theory [Mel63]. More precisely, here we define the Poincaré-Melnikov potential

$$\begin{aligned} \mathcal{L}^\varepsilon(\gamma^0) &= \int_0^{+\infty} H_2 \circ Z^\varepsilon(t, \gamma^0) - H_2 \circ Z_{\max}^\varepsilon(t, \gamma^0) dt \\ &\quad + \int_{-\infty}^0 H_2 \circ Z^\varepsilon(t, \gamma^0) - H_2 \circ Z_{\min}^\varepsilon(t, \gamma^0) dt, \end{aligned}$$

similarly to [DG00]), but taking into account the fact that the first order perturbation does not break integrability.

In other words, we consider as an integrable system $\mathcal{H}_0 = H_0 + \varepsilon H_1$ and as a perturbation $\mathcal{H}_1 = H_2$ (see Lemma 2.1 and recall that we have taken $\varepsilon = L_2^{-1}$). In Lemma 3.1 we have obtained the parameterization Z^0 of the separatrix (see (25)). For the Hamiltonian \mathcal{H}_0 , we have a parameterization of the separatrix $Z^\varepsilon(t, \gamma^0) = Z^0(t, \gamma^0) + O(\varepsilon)$. The time origin can be chosen so that

$$Z^\varepsilon(0, \gamma^0) = \left(\frac{\pi}{2}, \Gamma \sqrt{\frac{5}{3}} + O(\varepsilon), \gamma^0, \Gamma \right),$$

(recall that by Lemma 5.1, $Z^0(0, \gamma^0) = (\pi/2, \Gamma \sqrt{5/3}, \gamma^0, \Gamma)$).

In order to determine the splitting of separatrices, we consider the section $g_1 = \pi/2$ which is transversal to the flow locally in the neighborhood of the unperturbed separatrix. It is still transversal to the perturbed stable and unstable invariant manifolds. We measure the splitting in this section. The distance between of the invariant manifolds is parameterized by γ^0 , the value of the γ coordinate when $t = 0$.

Melnikov Theory ensures that the transversal homoclinic points in the section $g_1 = \pi/2$ are $\varepsilon^2 = L_2^{-2}$ -close to the non-degenerate critical points of the Poincaré-Melnikov potential. Expanding the potential in powers of ε , we get $\mathcal{L}^\varepsilon(\gamma^0) = \mathcal{L}(\gamma^0) + \mathcal{O}(\varepsilon)$ with

$$\begin{aligned} \mathcal{L}(\gamma^0) &= \int_0^{+\infty} H_2 \circ Z^0(t, \gamma^0) - H_2 \circ Z_{\max}^0(t, \gamma^0) dt \\ &\quad + \int_{-\infty}^0 H_2 \circ Z^0(t, \gamma^0) - H_2 \circ Z_{\min}^0(t, \gamma^0) dt. \end{aligned}$$

Moreover, since H_2 has a factor e_1 (see (7)) it vanishes over the periodic orbits $Z_{\min, \max}^0$. Thus, using (18) and (25), the potential reads

$$\begin{aligned} \mathcal{L}(\gamma^0) &= \int_{-\infty}^{+\infty} H_2(g_1^h(t), G_1^h(t), \gamma^h(t), \Gamma) dt \\ &= \int_{-\infty}^{+\infty} H_2(g_1^h(t), G_1^h(t), \gamma^0 + \gamma^1(t) + \gamma^2(t), \Gamma) dt. \end{aligned} \tag{31}$$

Melnikov theory then implies that, ε -close to non-degenerate critical points of \mathcal{L} , there exist transversal homoclinic points in the section $g_1 = \pi/2$. The potential \mathcal{L} naturally depends on parameters L_1 and Γ , which vary in the (non-empty) open set

$$O = \left\{ (L_1, \Gamma) : L_1 > 0, 0 < \Gamma < L_1 \sqrt{3/5} \right\} \subset \mathbb{R}^2. \quad (32)$$

Let us write

$$\mathcal{L}(\gamma^0) = \mathcal{L}_{L_1, \Gamma}(\gamma^0).$$

The next proposition shows that \mathcal{L} necessarily has non-degenerate critical points, except maybe for exceptional values of the parameters.

Proposition 4.1. *There exists a non constant real analytic function $\mathcal{L}^+ : O \rightarrow \mathbb{R}$ such that the potential (31) is of the form*

$$\mathcal{L}_{L_1, \Gamma}(\gamma^0) = \mathcal{L}_{L_1, \Gamma}^+ e^{i\gamma_0} + \overline{\mathcal{L}_{L_1, \Gamma}^+} e^{-i\gamma_0}.$$

In particular, outside an analytic subset of O of empty interior, \mathcal{L}^+ does not vanish and thus \mathcal{L} (as a function of γ^0) has non-degenerate critical points.

This proposition is proven in Section 5, where the function \mathcal{L}^+ is computed explicitly (see formula (38)). It is easier to describe the dynamics when the dimension is as low as possible, so let us carry out the symplectic reduction of the flow by time. Since $\dot{\gamma} \neq 0$, define the Poincaré section $\Sigma_{\gamma_0} = \{\gamma = \gamma_0\}$ within some fixed energy level, and the corresponding return map, induced by the Hamiltonian (11),

$$\mathcal{P}_{\gamma_0} : \Sigma_{\gamma_0} \longrightarrow \Sigma_{\gamma_0}$$

for which the Poincaré coordinates (ξ, η) may be used (see (20)). This map has a hyperbolic fixed point L_2^{-2} -close to the origin with one dimensional stable and unstable invariant manifolds. The classical Melnikov theory applied to the Melnikov potential obtained in Proposition 4.1 entails the following corollary and theorem. Note that the circle $g_1 = \pi/2$ locally corresponds to the line $\xi = 0$ (see (20)).

Corollary 4.2. *Fix $(L_1^0, \Gamma^0) \in O$ (defined in (32)) such that $\mathcal{L}_{L_1^0, \Gamma^0}^+ \neq 0$ and let γ_0^* be a non-degenerate critical point of $\mathcal{L}_{L_1^0, \Gamma^0}^+$. For L_2 large enough, there exists some $\tilde{\gamma}_0^* = \gamma_0^* + O(L_2^{-1})$ such that the Poincaré map $\mathcal{P}_{\tilde{\gamma}_0^*}$ has a transversal homoclinic point in the line $\{\xi = 0\} \subset \Sigma_{\tilde{\gamma}_0^*}$.*

Recall that all Poincaré maps are conjugate. Therefore, this corollary implies that there are transversal homoclinic points for \mathcal{P}_{γ_0} for all $\gamma_0 \in \mathbb{T}$.

The transversality of invariant manifolds given by the corollary a priori refers to the secular Hamiltonian obtained after one step of averaging. Nevertheless, the conclusion holds for the Poincaré maps of the secular Hamiltonian of any finite order. Indeed, higher order averaging only modifies higher orders of the asymptotic expansion of the splitting, while transversality at the first order was entailed by the first order of the expansion, as given by the Poincaré-Melnikov potential. Hence, by considering the analytic set defined by the condition $\mathcal{L}_{L_1^0, \Gamma^0}^+ \neq 0$ at all orders of averaging, one gets the main result, where we restrict to some fixed compact set

$$K \subset O = \left\{ (L_1, \Gamma) : L_1 > 0, 0 < \Gamma < L_1 \sqrt{3/5} \right\} \subset \mathbb{R}^2.$$

Theorem 4.3. *Fix any $\gamma_0 \in \mathbb{T}$. There is an analytic set K^o of full measure in K such that, for parameters (L_1^0, Γ^0) in K^o , for L_2 large enough, the Poincaré maps $\mathcal{P}_{\gamma_0} : \Sigma_{\gamma_0} \mapsto \Sigma_{\gamma_0}$ of the secular systems of all orders possess a horseshoe.*

4.1 Splitting of separatrices for the non-averaged Hamiltonian

As explained in Remark 3.2, for the secular Hamiltonian we have a hyperbolic periodic orbit at each energy level within a compact interval of such levels. These periodic orbits form a normally hyperbolic invariant cylinder. Corollary 4.2 implies that the invariant manifolds of this cylinder intersect transversally.

In Section 5 we consider L_1 and L_2 as constants. One can lift the dynamics to the extended phase space, with the additional Keplerian coordinates $(\ell_1, L_1, \ell_2, L_2)$. Then, the cylinder gets enlarged by four extra dimensions: (L_1, L_2) , which are just constants of motion, and (ℓ_1, ℓ_2) , which are performing a rigid rotation.

This extended system, after rescaling, is just the first order of (3), that is

$$F_0 = F_{\text{Kep}} + F_{\text{sec}}.$$

Call Λ the cylinder of this Hamiltonian. Now one would like to analyze Hamiltonian (3), that is, the full (non averaged) three-body problem. The first step is to prove the persistence of the invariant cylinder when L_2 is large enough, using that the reminder is $O(L_2^{-k})$ for some $k \in \mathbb{N}$. Since we are in a singular perturbation regime, the classical theory of persistence of normally hyperbolic invariant manifolds [Fen72] cannot be applied directly. However, now there are results which deal with singularly perturbed problems and which can be applied to the present setting [Yan09, GdlLT15]. Note that we can do global averaging up to high order. Therefore, one does not need to face the problems in [BKZ11] and the obtained cylinder can be as smooth as needed. Call $\tilde{\Lambda}$ the perturbed cylinder.

In this setting, one can expect Arnold diffusion i.e., orbits whose action components drift by a large amount, uniformly with respect to large L_2 's. Since

- G_1 is prescribed by $\tilde{\Lambda}$ and the homoclinic channel,
- G_2 (resp. L_1) is constrained by the conservation of the angular momentum (resp. the energy),

one would expect to obtain orbits with drift in L_2 , that is in the semi major axis of the outer body. This is a remarkable feature, since semi major axes are known to be very stable when two of the three masses are very small [Nie96].

To obtain unstable orbits one usually combines two types of dynamics. The “inner dynamics” is the dynamics of Hamiltonian (3) in restriction to the cylinder $\tilde{\Lambda}$. The “outer dynamics” and is the so-called scattering map [DdlLS08], obtained as the following limit. Assume that the invariant manifolds of $\tilde{\Lambda}$ split transversally along a homoclinic channel. Consider a homoclinic orbit in the channel, which is asymptotic to the trajectory of some point $x_- \in \tilde{\Lambda}$ as $t \rightarrow -\infty$ and to the trajectory of some other point $x_+ \in \tilde{\Lambda}$ as $t \rightarrow +\infty$. Then, we say that the scattering map \mathcal{S} maps x_- to x_+ . hat \mathcal{S} be a map (as opposed to a more general correspondance) is proved in [DdlLS08] under general hypotheses which are satisfied here. Note that this map depends on the chosen homoclinic channel and therefore it may not be defined globally –usually it is multivaluated. Provided that these two maps do not have common invariant circles, by iterating them in a random order, one gets pseudo-orbits which have the wanted unstable behavior. A shadowing argument then permits to approximately realize these pseudo-orbits as real orbits of the system.

Understanding both the inner and outer dynamics is certainly not easy for Hamiltonian (3). Concerning the inner dynamics, Jefferys and Moser [JM66] have used KAM theory to show that this cylinder contains quasiperiodic hyperbolic tori which form a

positive measure set. There should be a very rich dynamics in the complement of these tori in the cylinder. In particular, since the Hamiltonian restricted to the cylinder has three degrees freedom, there may exist Arnold diffusion in the cylinder itself.

Concerning the outer dynamics, one needs first to prove that the invariant manifolds of the cylinder $\tilde{\Lambda}$ split transversally and then derive some precise dynamical behavior for the scattering map, as in [DdlLS08]. The analysis in the present paper allows us to perform only the first step.

Theorem 4.4. *Fix $L_1^+ > L_1^- > 0$. Consider the Hamiltonian (3) with $L_1 \in [L_1^-, L_1^+]$ and $L_2 \geq L_2^0$. If L_2^0 is large enough,*

1. *there exists a normally hyperbolic invariant cylinder $\tilde{\Lambda}$,*
2. *the invariant manifolds of the cylinder $\tilde{\Lambda}$ intersect transversally along a homoclinic channel, which is diffeomorphic to $\tilde{\Lambda}$,*
3. *and there exists a scattering map $\mathcal{S} : \tilde{\Lambda} \rightarrow \tilde{\Lambda}$ associated to this homoclinic channel.*

Note that, in the above statement, the cylinder is invariant in the sense that the vector field is tangent to the cylinder, but orbits may escape from its boundary (sometimes one rather refers to these manifolds as weakly invariant).

Proof. As explained above, the persistence of the invariant cylinder can be obtained by the available results of persistence of normally hyperbolic invariant cylinders in the singular perturbation setting [Yan09, GdlLT15].

For the other statements, we may use the results in Proposition 4.1 and [DdlLS08]. Using the expression of F_{sec} in (5), one can split the Hamiltonian (3) as $F = H_0 + H_1$ with $H_0 = F_{\text{Kep}} - \mu_1 m_2 F_{\text{quad}}$ and $H_1 = F - H_0$. Therefore,

$$H_1 = -\mu_1 m_2 (\sigma_0 - \sigma_1) F_{\text{oct}} + O\left(\frac{a_1^4}{a_2^5}\right) - \mu_1 m_2 (\sigma_0 - \sigma_1) F_{\text{oct}} + O(L_2^{-5}).$$

and therefore satisfies $H_1 = O(L_2^{-2})$.

Let us abuse notation and assume that H_0 and H_1 are expressed in the variables given in (9) and that time has been suitably scaled. As we have explained the Hamiltonian H_0 has as normally hyperbolic invariant cylinder. Recall that L_1 , L_2 and Γ are first integrals of H_0 . Thus, the dynamics in this cylinder is foliated by three dimensional invariant tori. Let us define $Y_0 \equiv Y_0(t, \ell_1^0, L_1^0, \ell_2^0, L_2^0, \gamma^0, \Gamma^0)$ the trajectory on the invariant tori $L = L_i^0$, and $\Gamma = \Gamma^0$ in the cylinder with initial condition in the (ℓ_1, ℓ_2, γ) variables given by $(\ell_1^0, \ell_2^0, \gamma^0)$.

Since H_0 is integrable, the stable and unstable invariant manifolds of the cylinder agree, and form a homoclinic manifold. The latter is given (at the first order) by the homoclinic of the quadrupolar Hamiltonian obtained in Lemma 3.1 in the space $(g_1, G_1, \gamma, \Gamma)$. Recall that for H_0 the variables L_1 and L_2 are first integrals. The dynamics of the variables ℓ_1 and ℓ_2 is close to a rigid rotation and can be easily deduced. These homoclinic manifold can be parameterized by time t and the coordinates of the cylinder $Y \equiv Y(t, \ell_1^0, L_1^0, \ell_2^0, L_2^0, \gamma^0, \Gamma^0)$. Note that Y can be asymptotic to different points in the cylinder as $t \rightarrow \pm\infty$. There exist smooth functions $(\ell_1^\pm, \ell_2^\pm, \gamma^\pm) = (\ell_1^\pm, \ell_2^\pm, \gamma^\pm)(\ell_1^0, L_1^0, \ell_2^0, L_2^0, \gamma^0, \Gamma^0)$ such that

$$Y(t, \ell_1^0, L_1^0, \ell_2^0, L_2^0, \gamma^0, \Gamma^0) - Y_0(t, \ell_1^\pm, L_1^0, \ell_2^\pm, L_2^0, \gamma^\pm, \Gamma^0) \rightarrow 0$$

as $t \rightarrow \pm\infty$.

Then, to prove that the invariant manifolds split and that one can define the scattering map of the full Hamiltonian $H_0 + H_1$, one may apply the results in [DdlLS08] (see also [DdlLS06]). Define the Poincaré-Melnikov potential

$$\tilde{\mathcal{L}}(\ell_1^0, L_1^0, \ell_2^0, L_2^0, \gamma^0, \Gamma^0) = \int_{-\infty}^0 H_1 \circ Y - H_1 \circ Y_0^- dt + \int_0^{+\infty} H_1 \circ Y - H_1 \circ Y_0^+ dt,$$

where Y stands for $Y(t, \ell_1^0, L_1^0, \ell_2^0, L_2^0, \gamma^0, \Gamma^0)$ and Y_0^\pm for $Y_0(t, \ell_1^\pm, L_1^0, \ell_2^\pm, L_2^0, \gamma^\pm, \Gamma^0)$.

Consider the function

$$\tau \mapsto \tilde{\mathcal{L}}(\ell_1^0 + \omega_1\tau, L_1^0, \ell_2^0 + \omega_2\tau, L_2^0, \gamma^0 + \omega_3\tau, \Gamma^0) \quad (33)$$

where $\omega = (\omega_1, \omega_2, \omega_3)$ is the frequency vector associated with the torus $L = L_i^0$, and $\Gamma = \Gamma^0$. Results in [DdlLS08] imply that each non-degenerate critical point of this function gives rise to a transversal intersection of the invariant manifolds. The non-degeneracy of the critical point allows us to define a local scattering map.

Using the formula of H_1 , we see that

$$\tilde{\mathcal{L}} = \mathcal{L} + O(L_2^{-2}).$$

where \mathcal{L} is the Melnikov potential introduced in (31). Proposition 4.1 implies that, as long as $\mathcal{L}_{L_1^0, \Gamma^0}^+ \neq 0$, the function $\mathcal{L}(\gamma^0 - \omega_1\tau)$ has non-degenerate critical points with respect to τ . Thanks to the non-degeneracy, the function (33) has critical points, which are $O(L_2^{-2})$ -close to those of $\mathcal{L}(\gamma^0 - \omega_1\tau)$. Each critical point of (33) gives rise to a transversal intersections and to an associated scattering map. \square

Theorem 4.4 implies that Hamiltonian (3) fits in the classical framework of Arnold diffusion along single resonances: a normally hyperbolic invariant cylinder whose invariant manifolds intersect transversally [DdlLS00, DdlLS06, BKZ11]. Nevertheless, the results obtained in this paper are not enough to obtain unstable orbits drifting along the cylinder. Indeed, we cannot derive formulas for the scattering map and therefore we have no information about the outer dynamics beyond the fact that it is well defined.

The reason is the following. To prove that the invariant manifolds of the cylinder split, it is enough to project them into a certain plane and see that they intersect transversally in this plane. But, in order to build unstable orbits, one needs more precise information on the scattering map. Namely, to derive a first order approximation of the scattering map, one needs to analyze how the invariant manifolds of the cylinder split in every direction (see [DdlLS08]), in particular in the L_1 and L_2 directions. This would give the size of the possible jumps that the scattering maps makes in these directions. That is, how far can be two points in the cylinder which are connected by a heteroclinic orbit. Since the mean anomalies ℓ_1 and ℓ_2 are rapidly oscillating in our regime, the transversality in the conjugate directions L_1 and L_2 is exponentially small and therefore very difficult to detect.

An intermediate step would be to tackle the 1-averaged Hamiltonian, where only the fastest mean anomaly ℓ_1 has been averaged out. Then, one has a three degree of freedom Hamiltonian system with only one fast frequency. As a starter, we provide the expression of the analogue to the quadrupolar term:

$$-\frac{\mu_1 m_2}{\|q_2\|} \int_{\mathbb{T}} \sigma_2 P_2(\cos \zeta) \frac{\|q_1\|^2}{\|q_2\|^3} d\ell_1 =$$

$$-\frac{3\mu_1 m_2}{4a_1^2 r_2^3} \begin{pmatrix} ((4e_1^2 + 1) \sin^2 g_1 + (1 - e_1^2) \cos^2 g_1) \cos^2 i \sin^2 (v_2 + g_2) + \\ 10e_1^2 \cos g_1 \sin g_1 \cos i \cos (v_2 + g_2) \sin (v_2 + g_2) + \\ ((1 - e_1^2) \sin^2 g_1 + (4e_1^2 + 1) \cos^2 g_1) \cos^2 (v_2 + g_2) \\ -e_1^2 - \frac{2}{3} \end{pmatrix}.$$

where $v_2 \equiv v_2(G_2, L_2, \ell_2)$ is the eccentric anomaly.

Analyzing how the invariant manifolds split either for the full three-body problem or for the 1-averaged Hamiltonian, and deriving formulas for the corresponding scattering maps, would be the major step towards proving Arnold diffusion in the three body problem in the lunar regime.

5 Proof of Proposition 4.1

In Lemma 3.1, we have called $(g_1^h, G_1^h, \gamma^h = \gamma^0 + \gamma^1 + \gamma^2, \Gamma^h \equiv \Gamma)$ the separatrix solution.⁴ We compute the potential defined in (31). To this end, we expand it in Fourier series in γ^0 . Since H_2 is a trigonometric polynomial of degree 1 in γ (or equivalently in g_2 , see equation (7)), it can be written as

$$H_2(g_1, G_1, \gamma, \Gamma) = H_2^+(g_1, G_1, \Gamma)e^{i\gamma} + H_2^-(g_1, G_1, \Gamma)e^{-i\gamma}.$$

Because $\gamma^h = \gamma^0 + \gamma^1(t) + \gamma^2(t)$ depends linearly on γ^0 , $\mathcal{L}(\gamma^0)$ too has only two harmonics:

$$\mathcal{L}(\gamma_0) = \mathcal{L}^+ e^{i\gamma^0} + \mathcal{L}^- e^{-i\gamma^0}, \quad (34)$$

where

$$\mathcal{L}^\pm = \int_{-\infty}^{\infty} H_2^\pm(g_1^h(t), G_1^h(t), \Gamma) e^{\pm i(\gamma^1(t) + \gamma^2(t))} dt.$$

Since \mathcal{L} is a real function, it is determined by, say, the positive harmonic $\mathcal{L}^+ = \bar{\mathcal{L}}^-$.

As a first step, we parameterize the separatrix by g_1 instead of t , using the following lemma (where we omit the upper index h).

Lemma 5.1. *The following identities are satisfied on the separatrix given by Lemma 3.1:*

$$\begin{cases} G_1 &= |\Gamma| \sqrt{\frac{5}{3}} \frac{\sin g_1}{\sqrt{1 - \frac{5}{3} \cos^2 g_1}} \\ e_1 &= \sqrt{\frac{2}{3}} \frac{\Gamma}{L_1 \chi} \sqrt{\frac{1 - \frac{5}{3}(1 + \chi^2) \cos^2 g_1}{1 - \frac{5}{3} \cos^2 g_1}} \\ \cos i &= \sqrt{\frac{3}{5}} \frac{\sqrt{1 - \frac{5}{3} \cos^2 g_1}}{\sin g_1} \\ \sin i &= \sqrt{\frac{2}{5}} \frac{1}{\sin g_1} \\ \cos \gamma^2 &= \sqrt{1 - \frac{5}{3} \cos^2 g_1} \\ \sin \gamma^2 &= -\sqrt{\frac{5}{3}} \cos g_1. \end{cases}$$

The proof of this lemma is a direct consequence of Lemma 3.1 and formulas (8), (10), (13). We can use this lemma to express the function $\mathcal{F}^+ = H_2^+ e^{i\gamma^2}$ on the separatrix as a function of g_1 .

⁴The variable t used in section 3 is a rescaled time, since in Section 3 we have simplified the quadrupolar Hamiltonian by some multiplicative constant, when replacing F_{quad} by H_0 .

Lemma 5.2. *The function \mathcal{F}^+ can be written, on the separatrix, as a function of g_1 as $\mathcal{F}^+ = \frac{1}{2}(\mathcal{F}_1 + i\mathcal{F}_2)$ with*

$$\begin{cases} \mathcal{F}_1 &= C_1 \frac{\sqrt{1 - \frac{5}{3}(1 + \chi^2) \cos^2 g_1}}{1 - \frac{5}{3} \cos^2 g_1} \cos g_1 \left(1 - \frac{5}{3} \frac{1 - \frac{11}{7} \frac{\Gamma^2}{L_1^2}}{1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2}} \cos^2 g_1 \right) \\ \mathcal{F}_2 &= C_2 \frac{\sqrt{1 - \frac{5}{3}(1 + \chi^2) \cos^2 g_1}}{\sqrt{1 - \frac{5}{3} \cos^2 g_1}} \left(\frac{5}{3} \frac{\Gamma^2}{L_1^2} \frac{\cos^2 g_1 (9 - 11 \cos^2 g_1)}{1 - \frac{5}{3} \cos^2 g_1} + \right. \\ &\quad \left. \frac{21}{5} - 5 \frac{\Gamma^2}{L_1^2} - \left(14 - 11 \frac{\Gamma^2}{L_1^2} \right) \cos^2 g_1 \right) \end{cases}$$

and

$$C_1 = \frac{105 a_1^3}{32 a_2^4} \frac{e_2}{(1 - e_2^2)^{5/2}} \left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2} \right)^{3/2} \quad \text{and} \quad C_2 = \frac{15}{64} \sqrt{\frac{5}{3}} \frac{a_1^3}{a_2^4} \frac{e_2}{(1 - e_2^2)^{5/2}} \left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2} \right)^{1/2}.$$

Proof. We have

$$H_2 = A_{\text{oct}} e_1 (\cos g_1 \cos \gamma A + \sin g_1 \sin \gamma \cos i B)$$

where

$$\begin{cases} A = \frac{G_1^2}{L_1^2} (5 \sin^2 i (-7 \cos^2 g_1 + 6) - 3) - 35 \sin^2 g_1 \sin^2 i + 7 \\ B = \frac{G_1^2}{L_1^2} (5 \sin^2 i (7 \cos^2 g_1 - 4) + 3) + 35 \sin^2 g_1 \sin^2 i - 7 \end{cases}$$

and

$$A_{\text{oct}} = -\frac{15 a_1^3}{64 a_2^4} \frac{e_2}{(1 - e_2^2)^{5/2}}.$$

Thus,

$$\begin{aligned} \mathcal{F}^+ &= \frac{A_{\text{oct}} e_1}{2} \left[(A \cos g_1 \cos \gamma_2 + B \sin g_1 \sin \gamma_2 \cos i) \right. \\ &\quad \left. + i (A \cos g_1 \sin \gamma_2 - B \sin g_1 \cos \gamma_2 \cos i) \right], \end{aligned}$$

which we want to express in terms of g_1 . Using Lemma 5.1, the functions A and B can be written as

$$\begin{cases} A = \frac{5}{3} \frac{\Gamma^2}{L_1^2} \frac{1}{1 - \frac{5}{3} \cos^2 g_1} (9 - 11 \cos^2 g_1) - 7 \\ B = -\frac{5}{3} \frac{\Gamma^2}{L_1^2} \frac{1}{1 - \frac{5}{3} \cos^2 g_1} (5 - 11 \cos^2 g_1) + 7. \end{cases}$$

Let

$$\begin{cases} S_1 = e_1 (\cos g_1 \cos \gamma_2 A + \sin g_1 \sin \gamma_2 \cos i B) \\ S_2 = e_1 (\cos g_1 \sin \gamma_2 A - \sin g_1 \cos \gamma_2 \cos i B). \end{cases}$$

Then,

$$S_1 = -\frac{14\Gamma}{L_1 \chi} \sqrt{\frac{2}{3}} \left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2} \right) \frac{\sqrt{1 - \frac{5}{3}(1 + \chi^2) \cos^2 g_1}}{1 - \frac{5}{3} \cos^2 g_1} \cos g_1 \left(1 - \frac{5}{3} \frac{1 - \frac{11}{7} \frac{\Gamma^2}{L_1^2}}{1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2}} \cos^2 g_1 \right)$$

and

$$\begin{aligned} S_2 &= -\frac{\sqrt{10}}{3} \frac{\Gamma}{L_1 \chi} \frac{\sqrt{1 - \frac{5}{3}(1 + \chi^2) \cos^2 g_1}}{\sqrt{1 - \frac{5}{3} \cos^2 g_1}} \times \\ &\quad \left(\frac{5}{3} \frac{\Gamma^2}{L_1^2} \frac{\cos^2 g_1 (9 - 11 \cos^2 g_1)}{1 - \frac{5}{3} \cos^2 g_1} + \frac{21}{5} - 5 \frac{\Gamma^2}{L_1^2} - \left(14 - 11 \frac{\Gamma^2}{L_1^2} \right) \cos^2 g_1 \right). \end{aligned}$$

□

Now we compute \mathcal{F}^+ as a function of t . Recall that the constant A_2 was defined in (23).

Lemma 5.3. *The functions \mathcal{F}_1 and \mathcal{F}_2 on the heteroclinic (cf. Lemma 5.2), can be written as functions of $\tau = A_2 t$ as*

$$\begin{cases} \mathcal{F}_1 = \tilde{C}_1 \frac{\sinh \tau}{1 + \sinh^2 \tau} \cdot \frac{7 + 6 \sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau} \\ \mathcal{F}_2 = C_2 \frac{1}{\cosh \tau} \left[\frac{21}{5} - 5 \frac{\Gamma^2}{L_1^2} + \frac{\sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau} \left(\frac{\Gamma^2}{L_1^2 \chi^2} \frac{9\chi^2 + (\frac{12}{5} + 9\chi^2) \sinh^2 \tau}{\cosh^2 \tau} - \frac{3}{5} \left(14 - 11 \frac{\Gamma^2}{L_1^2} \right) \right) \right], \end{cases}$$

with

$$\tilde{C}_1 = -\frac{15}{16\sqrt{10}} \frac{a_1^3}{a_2^4} \frac{e_2}{(1 - e_2^2)^{5/2}} \frac{\Gamma}{L_1} \left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2} \right).$$

Proof. We compute each term as a function of τ . By (26), we have that

$$\sqrt{1 - \frac{5}{3}(1 + \chi^2) \cos g_1^0(t)} = \frac{\chi}{\sqrt{\chi^2 + (1 + \chi^2) \sinh^2 \tau}}$$

and

$$1 - \frac{5}{3} \cos g_1^0(t) = \chi^2 \frac{1 + \sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau}.$$

So,

$$\frac{\sqrt{1 - \frac{5}{3}(1 + \chi^2) \cos g_1^0(t)} \cos g_1^0(t)}{1 - \frac{5}{3} \cos g_1^0(t)} = \frac{1}{\chi} \sqrt{\frac{3}{5}} \frac{\sinh \tau}{1 + \sinh^2 \tau}.$$

For the last term, we have

$$\left(1 - \frac{5}{3} \frac{1 - \frac{11}{7} \frac{\Gamma^2}{L_1^2}}{1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2}} \cos^2 g_1^0(t) \right) = \frac{\left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2} \right) \chi^2 + M \sinh^2 \tau}{\left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2} \right) (\chi^2 + (1 + \chi^2) \sinh^2 \tau)}$$

where

$$M = \left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2} \right) (1 + \chi^2) - \left(1 - \frac{11}{7} \frac{\Gamma^2}{L_1^2} \right).$$

Using the definition of χ in (23), we have that

$$\begin{aligned} \left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2} \right) \chi^2 &= \frac{2}{3} \frac{\Gamma^2}{L_1^2} \\ M &= \frac{4}{7} \frac{\Gamma^2}{L_1^2}. \end{aligned}$$

Therefore,

$$\left(1 - \frac{5}{3} \frac{1 - \frac{11}{7} \frac{\Gamma^2}{L_1^2}}{1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2}} \cos^2 g_1^0(t) \right) = \frac{2\Gamma^2}{21L_1^2 \left(1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2} \right)} \frac{7 + 6 \sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau}.$$

Putting all these formulas together and defining

$$\tilde{C}_1 = \frac{2}{21\chi} C_1 \sqrt{\frac{3}{5}} \frac{\Gamma^2}{L_1^2} \frac{1}{1 - \frac{5}{3} \frac{\Gamma^2}{L_1^2}}.$$

we complete the proof. \square

The formulas of Lemma 5.3 allow us to compute the Poincaré - Melnikov potential (34). We split it as $\mathcal{L}^+ = \mathcal{L}_1 + i\mathcal{L}_2$ with

$$\mathcal{L}_j = \frac{1}{2} \int_{\mathbb{R}} \mathcal{F}_j(t) e^{i\gamma^1(t)} dt, \quad j = 1, 2. \quad (35)$$

We first compute

$$\begin{aligned} \mathcal{L}_1 &= \tilde{C}_1 \int_{-\infty}^{\infty} \frac{\sinh(A_2 t)}{1 + \sinh^2(A_2 t)} \cdot \frac{7 + 6 \sinh^2(A_2 t)}{\chi^2 + (1 + \chi^2) \sinh^2(A_2 t)} e^{-i \frac{2\Gamma}{L_1^2} t} dt \\ &= \frac{\tilde{C}_1}{A_2} \int_{-\infty}^{\infty} \frac{\sinh \tau}{1 + \sinh^2 \tau} \cdot \frac{7 + 6 \sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau} e^{-i \frac{2\Gamma}{A_2 L_1^2} \tau} d\tau. \end{aligned}$$

using the residue theorem. First note that if we look at this integral along the path $\tau = -i\pi + \sigma$, $\sigma \in \mathbb{R}$, instead of the real line, we get

$$\begin{aligned} &\int_{-i\pi + \mathbb{R}} \frac{\sinh \tau}{1 + \sinh^2 \tau} \cdot \frac{7 + 6 \sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau} e^{-i \frac{2\Gamma}{A_2 L_1^2} \tau} d\tau \\ &= -e^{-\frac{2\Gamma\pi}{A_2 L_1^2}} \int_{\mathbb{R}} \frac{\sinh \sigma}{1 + \sinh^2 \sigma} \cdot \frac{7 + 6 \sinh^2 \sigma}{\chi^2 + (1 + \chi^2) \sinh^2 \sigma} e^{-i \frac{2\Gamma}{A_2 L_1^2} \sigma} d\sigma. \end{aligned}$$

So, if we define the function

$$f_1(\tau) = \frac{\sinh \tau}{1 + \sinh^2 \tau} \cdot \frac{7 + 6 \sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau} e^{-i \frac{2\Gamma}{A_2 L_1^2} \tau}, \quad (36)$$

Cauchy's integral formula shows

$$\mathcal{L}_1 = -\frac{2\pi i}{1 + e^{-\frac{2\Gamma\pi}{A_2 L_1^2}}} \sum \text{Residues} \quad (37)$$

where the summation stands over the residues of the function f_1 in $-\pi < \text{Im } \tau < 0$; the negative sign comes from the index of the curve.

Lemma 5.4. *The function f_1 defined in (36) has three singularities in the region $-\pi < \text{Im } \tau < 0$ given by*

$$\begin{cases} a_0^- = -i \frac{\pi}{2} \\ a_1^- = -i \arcsin \frac{\chi}{\sqrt{1+\chi^2}} \\ a_2^- = -i \left(\pi + \arcsin \frac{\chi}{\sqrt{1+\chi^2}} \right) \end{cases}$$

and the associated residues are

$$\begin{cases} \text{Res}(f, a_0^-) = -\frac{2\Gamma}{A_2^2} e^{-\frac{\pi\Gamma}{A_2 L_1^2}} \\ \text{Res}(f, a_1^-) = \frac{7+\chi^2}{2\sqrt{1+\chi^2}} e^{-\frac{2\Gamma}{A_2 L_1^2} \arcsin \frac{\chi}{\sqrt{1+\chi^2}}} \\ \text{Res}(f, a_2^-) = -\frac{7+\chi^2}{2\sqrt{1+\chi^2}} e^{-\frac{2\Gamma}{A_2 L_1^2} \left(\pi - \arcsin \frac{\chi}{\sqrt{1+\chi^2}} \right)}. \end{cases}$$

Before proving the lemma, we proceed to deduce the value of the real part of the Melnikov integral. The lemma shows that the sum of formula (37) is

$$\sum \text{Residues} = -\frac{2\Gamma}{A_2 L_1^2} e^{-\frac{\pi\Gamma}{A_2 L_1^2}} + \frac{7 + \chi^2}{2\sqrt{1 + \chi^2}} e^{-\frac{2\Gamma}{A_2 L_1^2} \arcsin \frac{\chi}{\sqrt{1+\chi^2}}} \left(1 - e^{-\frac{2\Gamma}{A_2 L_1^2} \pi} \right),$$

hence the function \mathcal{L}_1 introduced in (35) satisfies

$$\mathcal{L}_1 = -\frac{2\pi i}{1 + e^{-\frac{2\Gamma\pi}{A_2 L_1^2}}} \frac{\tilde{C}_1}{A_2} \left(\begin{array}{l} -\frac{2\Gamma}{A_2 L_1^2} e^{-\frac{\pi\Gamma}{A_2 L_1^2}} + \\ + \frac{7+\chi^2}{2\sqrt{1+\chi^2}} e^{-\frac{2\Gamma}{A_2 L_1^2} \arcsin \frac{\chi}{\sqrt{1+\chi^2}}} \left(1 - e^{-\frac{2\Gamma}{A_2 L_1^2} \pi} \right) \end{array} \right).$$

Proof of Lemma 5.4. The singularities are the solutions of

$$1 + \sinh^2 \tau = 0 \text{ or either } \chi^2 + (1 + \chi^2) \sinh^2 \tau = 0.$$

The first equation is just $1 + \sinh^2 \tau = \cosh^2 \tau = 0$ and the only possible solution in $-\pi < \text{Im } \tau < 0$ is $a_0^- = -i\pi/2$. For the second equation, one can take $\tau = i\sigma$. Then, it is equivalent to

$$\sin \sigma = \pm \frac{\chi}{\sqrt{1 + \chi^2}},$$

which gives the other two singularities.

Now we compute the residues. We start by a_0^- . We compute the Laurent series of each term in f . From the fact that

$$\sinh \tau = -i(1 + \mathcal{O}_2(\tau + i\pi/2))$$

one can easily deduce the following,

$$\begin{aligned} \frac{1}{\cosh^2 \tau} &= -\frac{1}{(\tau + i\pi/2)^2} (1 + \mathcal{O}_2(\tau + i\pi/2)) \\ 7 + 6 \sinh^2 \tau &= 1 + \mathcal{O}_2(\tau + i\pi/2) \\ \chi^2 + (1 + \chi^2) \sinh^2 \tau &= -1 + \mathcal{O}_2(\tau + i\pi/2). \end{aligned}$$

Therefore

$$\frac{\sinh \tau}{1 + \sinh^2 \tau} \cdot \frac{7 + 6 \sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau} = -\frac{i}{(\tau + i\pi/2)^2} (1 + \mathcal{O}_2(\tau + i\pi/2)).$$

For the exponential, we know have that

$$e^{-i\frac{2\Gamma}{A_2 L_1^2} \tau} = e^{-\frac{\pi\Gamma}{A_2 L_1^2}} - i\frac{2\Gamma}{A_2 L_1^2} e^{-\frac{\pi\Gamma}{A_2 L_1^2}} (\tau + i\pi/2) + \mathcal{O}_2(\tau + i\pi/2).$$

From these two last expansions, we obtain the residue for $a_0^- = -i\pi/2$.

Now we compute the other two residues. We compute them at the same time. Note that for $i = 1, 2$,

$$\sinh a_i^- = -i\frac{\chi}{\sqrt{1 + \chi^2}}$$

and

$$\cosh a_1^- = \frac{1}{\sqrt{1 + \chi^2}}, \quad \cosh a_2^- = -\frac{1}{\sqrt{1 + \chi^2}}.$$

Therefore

$$\begin{aligned} 7 + 6 \sinh^2 \tau &= \frac{7 + \chi^2}{1 + \chi^2} + \mathcal{O}_1(\tau - i\pi/2) \\ \chi^2 + (1 + \chi^2) \sinh^2 \tau &= -2\chi i(\tau - a_1^-) + \mathcal{O}_2(\tau - a_1^-) \\ \chi^2 + (1 + \chi^2) \sinh^2 \tau &= +2\chi i(\tau - a_2^-) + \mathcal{O}_2(\tau - a_2^-) \end{aligned}$$

and we have also

$$\begin{aligned} e^{-i\frac{2\Gamma}{A_2L_1^2}\tau} &= e^{-\frac{2\Gamma}{A_2L_1^2}\arcsin\frac{\chi}{\sqrt{1+\chi^2}}} + \mathcal{O}_1(\tau - a_1^-) \\ e^{-i\frac{2\Gamma}{A_2L_1^2}\tau} &= e^{-\frac{2\Gamma}{A_2L_1^2}\left(\pi - \arcsin\frac{\chi}{\sqrt{1+\chi^2}}\right)} + \mathcal{O}_1(\tau - a_2^-). \end{aligned}$$

With all these expansions, one can easily deduce the last two residues. \square

Analogously, the function \mathcal{L}_2 introduced in (35), satisfies

$$\mathcal{L}_2 = -\frac{C_2}{A_2} \frac{2\pi i}{1 + e^{-\frac{2\Gamma\pi}{A_2L_1^2}}} \sum \text{Residues},$$

where the sum stands over all residues in $-\pi < \tau < 0$ of

$$f_2(\tau) = \frac{e^{-i\frac{2\Gamma}{A_2L_1^2}\tau}}{\cosh \tau} \left[\frac{21}{5} - 5\frac{\Gamma^2}{L_1^2} + \frac{\sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau} \left(\frac{\Gamma^2}{L_1^2 \chi^2} \frac{9\chi^2 + \left(\frac{12}{5} + 9\chi^2\right) \sinh^2 \tau}{\cosh^2 \tau} - \frac{3}{5} \left(14 - 11\frac{\Gamma^2}{L_1^2} \right) \right) \right].$$

Lemma 5.5. *In the region $-\pi < \text{Im } \tau < 0$, the function f_2 has the same singularities as f_1 , given in Lemma 5.4. Moreover, the associated residues are given by*

$$\begin{cases} \text{Res}(f_2, a_0^-) = -\frac{i}{5} \left(21 - 8\frac{\Gamma^2}{L_1^2} + 24\frac{\Gamma^4}{A_2^2 L_1^6 \chi^2} \right) e^{-\frac{\pi\Gamma}{A_2L_1^2}} \\ \text{Res}(f_2, a_1^-) = -i\frac{3}{5} \frac{\chi}{(1+\chi^2)^{3/2}} \left(11\frac{\Gamma^2}{L_1^2} - 7 \right) e^{-\frac{2\Gamma}{A_2} \arcsin \frac{\chi}{\sqrt{1+\chi^2}}} \\ \text{Res}(f_2, a_2^-) = -i\frac{3}{5} \frac{\chi}{(1+\chi^2)^{3/2}} \left(11\frac{\Gamma^2}{L_1^2} - 7 \right) e^{-\frac{2\Gamma}{A_2} \left(\pi - \arcsin \frac{\chi}{\sqrt{1+\chi^2}} \right)} \end{cases}$$

Hence, the function \mathcal{L}_2 introduced in (35) satisfies

$$\mathcal{L}_2 = -\frac{C_2}{A_2} \frac{2\pi i}{1 + e^{-\frac{2\Gamma\pi}{A_2L_1^2}}} \left(\begin{aligned} &-\frac{i}{5} \left(21 - 8\frac{\Gamma^2}{L_1^2} + 24\frac{\Gamma^4}{A_2^2 L_1^6 \chi^2} \right) e^{-\frac{\pi\Gamma}{A_2} +} \\ &-i\frac{3}{5} \frac{\chi}{(1+\chi^2)^{3/2}} \left(11\frac{\Gamma^2}{L_1^2} - 7 \right) e^{-\frac{2\Gamma}{A_2} \arcsin \frac{\chi}{\sqrt{1+\chi^2}} +} \\ &-i\frac{3}{5} \frac{\chi}{(1+\chi^2)^{3/2}} \left(11\frac{\Gamma^2}{L_1^2} - 7 \right) e^{-\frac{2\Gamma}{A_2} \left(\pi - \arcsin \frac{\chi}{\sqrt{1+\chi^2}} \right)} \end{aligned} \right).$$

Proof of Lemma 5.5. We use the expansions obtained in the proof of Lemma 5.4. For the last two residues, it is enough to use also that

$$\frac{\Gamma^2}{L_1^2 \chi^2} \frac{9\chi^2 + \left(\frac{12}{5} + 9\chi^2\right) \sinh^2 \tau}{\cosh^2 \tau} = \frac{33}{5} \frac{\Gamma^2}{L_1^2} + \mathcal{O}(\tau - a_i^-)$$

for $i = 1, 2$. Then, using the expansions of Lemma 5.4, we can deduce the two last residues.

Now we compute the residue at $\tau = a_0$. We split f_2 into three parts $f_2 = h_1 + h_2 + h_3$ with

$$\begin{cases} h_1(\tau) = \left(\frac{21}{5} - 5\frac{\Gamma^2}{L_1^2} \right) \frac{e^{-i\frac{2\Gamma}{A_2L_1^2}\tau}}{\cosh \tau} \\ h_2(\tau) = -\frac{3}{5} \left(14 - 11\frac{\Gamma^2}{L_1^2} \right) \frac{\sinh^2 \tau}{\chi^2 + (1+\chi^2) \sinh^2 \tau} \frac{e^{-i\frac{2\Gamma}{A_2L_1^2}\tau}}{\cosh \tau} \\ h_3(\tau) = \frac{\Gamma^2}{L_1^2 \chi^2} \frac{9\chi^2 + \left(\frac{12}{5} + 9\chi^2\right) \sinh^2 \tau}{\chi^2 + (1+\chi^2) \sinh^2 \tau} \frac{\sinh^2 \tau}{\cosh^3 \tau} e^{-i\frac{2\Gamma}{A_2L_1^2}\tau}. \end{cases}$$

Functions h_1 and h_2 have a pole of order one at $\tau = a_0^-$ so one can proceed as for the others singularities. However, h_3 has a pole of order 3 and therefore, we need to compute the expansion up to order 3 of each term in h_2 .

We use the expansions

$$\begin{cases} \sinh \tau &= -i - \frac{i}{2}y^2 + \mathcal{O}_4(y) \\ \cosh \tau &= -iy - \frac{i}{6}y^3 + \mathcal{O}_5(y) \end{cases}$$

where $y = \tau + i\frac{\pi}{2}$. We use this notation throughout the proof.

We start with h_1 and h_2 . It can be easily seen that

$$\begin{cases} \text{Res}(h_1, a_0^-) &= i \left(\frac{21}{5} - 5 \frac{\Gamma^2}{L_1^2} \right) e^{-\frac{\pi\Gamma}{A_2 L_1^2}} \\ \text{Res}(h_2, a_0^-) &= -i \frac{3}{5} \left(14 - 11 \frac{\Gamma^2}{L_1^2} \right) e^{-\frac{\pi\Gamma}{A_2 L_1^2}} \end{cases}$$

For h_3 we use the following expansions

$$\begin{aligned} \frac{9\chi^2 + \left(\frac{12}{5} + 9\chi^2\right) \sinh^2 \tau}{\chi^2 + (1 + \chi^2) \sinh^2 \tau} \sinh^2 \tau &= -\frac{12}{5} - \left(\frac{12}{5} + \frac{33}{5}\chi^2 \right) y^2 + \mathcal{O}_4(y) \\ \frac{1}{\cosh^3 \tau} &= -\frac{i}{y^3} \left(1 - \frac{y^2}{2} + \mathcal{O}_4(y) \right) \\ e^{-i\frac{2\Gamma}{A_2 L_1^2} \tau} &= e^{-\frac{\pi\Gamma}{A_2 L_1^2}} \left(1 - i \frac{2\Gamma}{A_2 L_1^2} y - \frac{2\Gamma^2}{A_2^2 L_1^4} y^2 + \mathcal{O}_4(y) \right). \end{aligned}$$

Putting together all these expansions one can deduce the residue

$$\text{Res}(h_3, a_0^-) = i \frac{\Gamma^2}{L_1^2 \chi^2} \left(\frac{33}{5} \chi^2 - \frac{24}{5} \frac{\Gamma^2}{A_2^2 L_1^4} \right) e^{-\frac{\pi\Gamma}{A_2 L_1^2}}$$

Now it only remains to add the three residues to obtain the residue of f_2 at $\tau = a_0^-$. \square

Gathering what precedes, and letting

$$\alpha = \frac{\pi\Gamma}{A_2 L_1^2}, \quad \hat{\Gamma} = \frac{\Gamma}{L_1}, \quad \hat{\chi} = \frac{\chi}{\sqrt{1 + \chi^2}} \quad \text{and} \quad \beta = \frac{\alpha}{\pi} \arcsin \hat{\chi},$$

we obtain the following analytic expression:

$$\mathcal{L}^+ = -\frac{2\pi i}{A_2(1 + e^{-2\alpha})} \left(\begin{array}{l} \tilde{C}_1 \left(\begin{array}{l} -\frac{2\alpha}{\pi} e^{-\alpha} + \\ + \frac{7 + \chi^2}{2\chi} \sin \frac{\beta\pi}{\alpha} e^{-2\beta} (1 - e^{-2\alpha}) \end{array} \right) \\ + C_2 \left(\begin{array}{l} \frac{1}{5} \left(21 - 8\hat{\Gamma}^2 + 24 \frac{\hat{\Gamma}^2 \alpha^2}{\pi^2 \chi^2} \right) e^{-\alpha} + \\ + \frac{3}{5} \frac{\chi}{(1 + \chi^2)^{3/2}} \left(11\hat{\Gamma}^2 - 7 \right) e^{-2\beta} + \\ + \frac{3}{5} \frac{\chi}{(1 + \chi^2)^{3/2}} \left(11\hat{\Gamma}^2 - 7 \right) e^{-2(\alpha - \beta)} \end{array} \right) \end{array} \right). \quad (38)$$

In order to check that \mathcal{L}^+ is not constant, notice that, asymptotically when Γ tends to 0 (and L_1 is kept constant), $\alpha = O(\Gamma^{-2})$ while $\beta = O(\Gamma^{-1})$, so that the term in $\tilde{C}_1 e^{-2\beta}$ dominates the others: for some $C_3(L_1)$ and $C_4(L_1)$ independent of Γ ,

$$\mathcal{L}^+ \sim \frac{C_3(L_1)}{\Gamma^2} \exp(-2C_4(L_1)\Gamma + O(\Gamma^{-2})).$$

Since the dominant term of the right hand side is a non constant function of Γ , Proposition 4.1 is proved.

Funding: J. F. is partially supported by the French ANR (projet ANR-10-BLAN 0102 DynPDE). M. G. is partially supported by the Spanish MINECO-FEDER Grant MTM2012-31714 and the Catalan Grant 2014SGR504.

Conflict of Interest: The authors declare that they have no conflict of interest.

References

- [Ale69] V. M. Alekseev. Quasirandom dynamical systems. *Mat. Zametki*, 6:489–498, 1969.
- [BKZ11] P. Bernard, V. Kaloshin, and K. Zhang. Arnold diffusion in arbitrary degrees of freedom and crumpled 3-dimensional normally hyperbolic invariant cylinders. Preprint available at <http://arxiv.org/abs/1112.2773>, 2011.
- [BL10] G. Boué and J. Laskar. A collisionless scenario for Uranus tilting. *Astrophysics J. Letters*, 712(1):L44, 2010.
- [BLF12] G. Boué, J. Laskar, and F. Farago. A simple model of the chaotic eccentricity of mercury. *Astron. & Astrophysics*, 2012.
- [Bol06] S. Bolotin. Symbolic dynamics of almost collision orbits and skew products of symplectic maps. *Nonlinearity*, 19(9):2041–2063, 2006.
- [DdlLS00] A. Delshams, R. de la Llave, and T.M. Seara. A geometric approach to the existence of orbits with unbounded energy in generic periodic perturbations by a potential of generic geodesic flows of \mathbb{T}^2 . *Comm. Math. Phys.*, 209(2):353–392, 2000.
- [DdlLS06] A. Delshams, R. de la Llave, and T.M. Seara. A geometric mechanism for diffusion in hamiltonian systems overcoming the large gap problem: heuristics and rigorous verification on a model. *Mem. Amer. Math. Soc.*, 2006.
- [DdlLS08] A. Delshams, R. de la Llave, and T. M. Seara. Geometric properties of the scattering map of a normally hyperbolic invariant manifold. *Adv. Math.*, 217(3):1096–1153, 2008.
- [DG00] A. Delshams and P. Gutiérrez. Splitting potential and the Poincaré-Melnikov method for whiskered tori in Hamiltonian systems. *J. Nonlinear Sci.*, 10(4):433–476, 2000.
- [DKdlRS14] A. Delshams, V. Kaloshin, A. de la Rosa, and T. Seara. Parabolic orbits in the restricted three body problem. Preprint, available at <http://arxiv.org/abs/1501.01214>, 2014.
- [Féj02] J. Féjóz. Quasiperiodic motions in the planar three-body problem. *J. Differential Equations*, 183(2):303–341, 2002.
- [Fen72] N. Fenichel. Persistence and smoothness of invariant manifolds for flows. *Indiana Univ. Math. J.*, 21:193–226, 1971/1972.

- [FGKR15] J. Féjóz, M. Guardia, V. Kaloshin, and P. Roldán. Kirkwood gaps and diffusion along mean motion resonances in the restricted planar three-body problem. *to appear, J. Eur. Math. Soc.*, 2015.
- [FS89] C. Froeschle and H. Scholl. The three principal secular resonances $\nu(5)$, $\nu(6)$, and $\nu(16)$ in the asteroidal belt. *Celestial Mechanics and Dynamical Astronomy*, 46:231–251, September 1989.
- [GdlLT15] M. Guardia, R. de la Llave, and R. Treviño. Normally hyperbolic invariant manifolds in hilbert spaces. Work in progress, 2015.
- [GMS13] M. Guardia, P. Martín, and T. M. Seara. Homoclinic solutions to infinity and oscillatory motions in the restricted planar circular three body problem. In *Progress and challenges in dynamical systems*, volume 54 of *Springer Proc. Math. Stat.*, pages 265–280. Springer, Heidelberg, 2013.
- [JM66] W. H. Jefferys and J. Moser. Quasi-periodic solutions for the three-body problem. *Astronom. J.*, 71:568–578, 1966.
- [Las08] J. Laskar. Chaotic diffusion in the Solar System. *Icarus*, 196(1):1–15, 2008.
- [LR93] J. Laskar and Ph. Robutel. The chaotic obliquity of the planets. *Nature*, 361(6413):608–612, 1993.
- [LS80a] J. Llibre and C. Simó. Oscillatory solutions in the planar restricted three-body problem. *Math. Ann.*, 248(2):153–184, 1980.
- [LS80b] J. Llibre and C. Simó. Some homoclinic phenomena in the three-body problem. *J. Differential Equations*, 37(3):444–465, 1980.
- [Mel63] V. K. Melnikov. On the stability of the center for time periodic perturbations. *Trans. Moscow Math. Soc.*, 12:1–57, 1963.
- [Moe89] R. Moeckel. Chaotic dynamics near triple collision. *Arch. Rational Mech. Anal.*, 107(1):37–69, 1989.
- [Mos01] J. Moser. *Stable and random motions in dynamical systems*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 2001. With special emphasis on celestial mechanics, Reprint of the 1973 original, With a foreword by Philip J. Holmes.
- [Nie96] L. Niederman. Stability over exponentially long times in the planetary problem. *Nonlinearity*, 9(6):1703–1751, 1996.
- [Sit60] K. Sitnikov. The existence of oscillatory motions in the three-body problems. *Soviet Physics. Dokl.*, 5:647–650, 1960.
- [Yan09] D. Yang. An invariant manifold for odes and its applications. Preprint available at <http://arxiv.org/abs/0909.1103v1>, 2009.
- [Zha14] L. Zhao. Quasi-periodic almost-collision orbits in the spatial three-body problem. *Com. Pure App. Math.*, 2014.